

Review Article



Breast Cancer: A Global Concern, Diagnostic and Therapeutic Perspectives, Mechanistic Targets in Drug Development

Gul-e-Saba Chaudhry^{1*}, Rehmat Jan², Abdah Akim³, Muhammad Naveed Zafar⁴, Yeong Yik Sung¹, Tengku Sifzizul Tengku Muhammad¹

¹Institute of Marine Biotechnology, University Malaysia Terengganu, 21030 Kuala Terengganu, Malaysia.

²Department of Environmental Sciences, Fatima Jinnah University, Rawalpindi, Pakistan.

³Department of Biomedical Sciences, Universiti Putra Malaysia, Seri Kembangan, Selangor, Malaysia.

⁴Department of Chemistry, Quaid-i-Azam University, Islamabad, 45320, Pakistan.

Article info

Article History:

Received: 27 Feb. 2020

Revised: 10 Aug. 2020

Accepted: 14 Oct. 2020

epublished: 14 Oct. 2020

Keywords:

- Apoptosis
- Breast cancer
- Cell cycle arrest
- Drug development
- Natural products
- Mechanism of action
- Phytochemicals
- Plant anticancer drugs

Abstract

Cancer is a complex multifactorial process, unchecked and abrupt division, and cell growth—conventional chemotherapy, along with radiotherapy, is used to treat breast cancer. Due to reduce efficacy and less survival rate, there is a particular need for the discovery of new active anticancer agents. Natural resources such as terrestrial/marine plants or organisms are a promising source for the generation of new therapeutics with improving efficacy. The screening of natural plant extracts and fractions, isolations of phytochemicals, and mechanistic study of those potential compounds play a remarkable role in the development of new therapeutic drugs with increased efficacy. Cancer is a multistage disease with complex signaling cascades. The initial study of screening whole extracts or fractions and later the isolation of secondary compounds and their mechanism of action study gives a clue of potential therapeutic agents for future drug development. The phytochemicals present in extracts/fractions produce remarkable effects due to synergistically targeting multiple signals. In this review, the molecular targets of extracts/ fractions and isolated compounds highlighted. The therapeutic agent's mechanistic targets in drug development focused involves; i) Induction of Apoptosis, ii) modulating cell cycle arrest, iii) Inhibition or suppression of invasion and metastasis and iv) various other pro-survival signaling pathways. The phytochemicals and their modified analogs identified as future potential candidates for anticancer chemotherapy.

Introduction

Breast cancer is the second leading cause of death (11.6% of the total cancer deaths), followed by colorectal and lung cancer.¹ Cancer is a complex disease concerning pathology and biochemistry. It begins when cells in the body start to grow in an uncontrolled and abnormal manner, which may also cause disturbances and alter the structure of surrounding tissues.² The evasion of apoptosis, limitless replicative potential, evading growth suppressors, sustaining proliferative signaling, inducing angiogenesis and activating tissue invasion and metastasis are critical features of cancer which contribute towards tumor development.^{3,4} Alterations in cellular DNA and transcriptional/translational processes causes irregularity in the gene expressions and results in cancer cell proliferation. Primary entities involved in carcinogenesis are oncogenes and tumor suppressor genes. Defects in tumor suppressor genes and mutations in the proto-oncogenes results in uncontrolled multiplication of cells leading to cancer.^{5,6} Breast cancer, lung cancer, and colorectal cancer are frequently occurring cancer in both men and women.⁷

Breast cancer: a global concern

Breast cancer triggers due to the uncontrolled multiplication of cells. It is the most frequently occurring cancer type and the leading cause of death in women over the last few years.⁸ The cancer trigger due to mutations in genes responsible for the production of pro-apoptotic/ anti-apoptotic proteins, tumor suppressors proteins, and growth factors. According to the United States, cancer statistics report 2018, an estimate of about 268 670 new breast cancer cases and a total of 41 400 deaths cases due to breast cancer in the United States in 2018.⁹ Breast cancer broadly categorized into two types, invasive breast cancer, and non-invasive breast cancer. However, other types of breast cancer include medullary and tubular carcinoma, inflammatory breast cancer, Paget's disease (PD), and phyllodes tumor (PT). Generally, in invasive breast cancer, cells are not only confined to ducts and lobular walls but also spread to surrounding areas of breast (connective and fatty tissues). The infiltrating-lobular-carcinoma and infiltrating-ductal-carcinoma are frequently occurring invasive-breast cancer. The lobular carcinoma is initiated

*Corresponding Author: Gul-e-Saba Chaudhry, Tel: +609-6683810, Fax: +609-6683810, Email: gul.saba@umt.edu.my

© 2021 The Author (s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

in the milk glands, while ductal carcinoma began in the breast's milk ducts. Medullary breast carcinoma and tubular carcinomas are the subtypes of invasive breast carcinoma.¹⁰ Furthermore, inflammatory breast cancer type is characterized by inflamed breasts with indentation and thick ridges. Only 1% to 2% of all invasive breast cancers and 1% of all breast cancers are inflammatory breast cancer with low survival rates at all stages. However, non-invasive breast cancer cells restricted to ducts only (do not penetrate surrounding tissues) of the breast. Ductal carcinoma in situ and lobular carcinoma in situ are the two forms of non-invasive breast cancer.¹⁰⁻¹³ PD pharmaceutically described by the infiltration of the nipple epidermis by destructive breast epithelial cells. PD of the breast defined as a skin alteration in the nipple-areola region. It is less common and generally linked with *in-situ* or invasive carcinoma.¹⁴ Breast PT, is a rare tumor, and shows different behavior, as it could be benign (non-cancerous) or malignant (cancerous).¹⁵ PT can cause uncommon fibroepithelial lesions to account for around 0.3% to 0.5% of breast tumors diagnosed in women and has an occurrence of about 2.1 per million.¹⁶

Breast cancer usually classified as two types i) estrogen receptor-positive (ER+) and ii) estrogen receptor-negative (ER-) breast cancer. Estrogen receptor-positive cell lines include MCF-7 and T-47D, while MDA-MB-231, MDA-MB-453 and MDA-MB-468 are estrogen negative receptor cell lines. Which, further characterized as luminal A (ER+, PR+, HER2-), luminal B (ER+, PR+, HER2+), HER2-enriched, basal-like, and normal-like based on progesterone receptor (PR) and human epidermal growth factor receptor-2 (HER2) status.¹⁷ The MCF-7 and T-47D cell line together with other breast cancer cell type MDA-MB-231, have been studied in above two-third of the total publications in Medline.¹⁸

Risk factors, diagnostic and therapeutic perspective for breast cancer

Common risk factors associated, such as age, gender, family history, breast density, radiation exposure, reproductive factors, genetic mutations, and diabetes.¹⁹ Early screening, detection, and diagnosis, significantly affect the occurrence and survival rate of breast cancer. Several diagnostic approaches include mammograms, ultrasound, magnetic resonance imaging, breast self-examination, positron emission tomography scan, computerized tomography, bone scintigraphy, chest X-ray, and biopsy.²⁰ However, due to some limitations of these approaches, such as high cost, time consumption, and age restriction, the development of highly sensitive and early-stage diagnostic techniques required. Different biomarkers such as proteomic biomarkers, gene biomarkers, and various imaging techniques are a useful analytical tool for fast and economic early-stage breast cancer diagnosis.²¹ Breast cancer conventional treatment approach involves; (i) surgical removal of cancer cells. (ii) use of chemotherapy

coupled with hormonal therapy and gene therapy; and (iii) radiation therapy.²² Surgery is considered as the earliest method and used for most of the solid tumors.²² The surgical treatment depends on the stage and tumor form; involves removal of the only lump (lumpectomy) or surgical removal of the entire breast (mastectomy). Breast-conserving surgery includes lumpectomy (removal of lump only or a small number of surrounding tissues), wide excision (partial mastectomy), and quadrantectomy (removal of about one-quarter of the breast).¹² Currently, sentinel lymph node dissection has become a well-known suitable technique as it necessitates the excision of very few lymph nodes, causing very few or no side effects. Over the past decade, advances in sentinel lymph node mapping have enhanced the precision of detecting sentinel lymph nodes from 80% to 92%-98% using different combined modalities.^{12,22}

Chemotherapy is the most conventional therapy available for malignant cancers.²² In chemotherapy, anticancer drugs, orally or intravenously given to patients, might cause severe side effects due to non-specific killing of cancer cells. However, radiotherapy is a conventional approach used in the treatment of cancer, along with chemotherapy.²² For the treatment of HER2-neu positive tumors, trastuzumab, in combination with radiotherapy, is needed.²³ Hormonal therapy studied for the treatment of ER+ breast cancer for several decades. The anticancer drug tamoxifen behaves as an antagonist in the breast, causing a delay in the transcription of estrogen-regulated genes and interrupting in the proliferative effects of estrogen in the breast. Similarly, fulvestrant acts as tamoxifen, but it causes degradation of the ER protein and loss of estrogen and progesterone receptor expression.²⁴ Menopausal hormone therapy usually restrained from breast cancer survivors because of the risk of reoccurrence. Menopausal hormone therapy provides adequate assistance from climacteric symptoms, but few are associated with enhanced risk of stroke and also breast, ovarian, and endometrial cancers.²⁵ Gene therapies have developed as promising new treatments for breast cancer. Proto-oncogene and tumor suppressor genes have shown accelerated improvement in gene therapy approaches.¹² Various clinical trials are ongoing to deliver p53 to cancer cells. The viral vectors have employed to transfer a breast cancer gene BRCA1, as a mutation in BRCA genes is also responsible for breast cancer cases. Also, the use of antisense strategies in clinical trials considered the most common approach. Adenoviral gene E1A that interferes with the transcription of erbB-2 can use to inhibit the transcription of overexpressed oncogenes in the treatment of ovarian and breast cancer.¹²

Plant-derived anti-breast cancer therapeutic agents

Natural products played a remarkable role in the prevention and treatment of cancer and remained a focus of research in drug discovery.²⁶ Over 3000 plant species reported having anticancer properties.²⁷ Plant-derived

natural products have significant efficacy in cancer treatment due to reduced adverse side effects as compared to conventional chemotherapy.²⁸ This review article's primary objective was to study the therapeutic potential of natural products in whole plant extracts/fractions or isolated secondary metabolites in breast cancer treatment. The initial stage in drug discovery is to screen the potential extracts and fractions, which gives the clue of the presence of novel phytochemicals. The knowledge provides a better understanding of the presence of various phytochemicals and their synergistic approach, which nowadays played a remarkable role in combination drug therapy. The extracts/fractions screening reduces the cost of isolation of phytochemicals by revealing a potential agent's presence. The isolated secondary metabolites from various natural sources, mechanistic study (in-vitro), give better molecular fundamental knowledge of the future therapeutic agent. Figure 1 describes the preparation of phytochemical compounds isolated from plants and their use in breast cancer therapy.

Anti-breast cancer therapeutic agents and their molecular mechanistic targets

Several crude extracts/fractions possessing potential natural products have tested on a variety of breast cancer cell lines. The potential phytochemicals induced cytotoxicity on breast cancer through several mechanisms. Such as via induction of apoptosis, cell cycle arrest in cancer cells, inhibition of metastatic potential, obstructing the process of angiogenesis, pro-survival signaling, and autophagy activation. Figures 2 and 3 provide detailed information about potential therapeutic approaches of several anti-breast cancer plant species and their mechanism of action, respectively.

Induction of apoptosis

Apoptosis, tightly regulated mechanism of cell death as a result of signal cascades involved during healthy development and morphogenesis.^{3,29} The enzymatic proteins caspases are prominent initiators and executioners in the process of apoptosis. Along with caspases, various pro and anti-apoptotic proteins such as Bcl-2 family proteins, tumor suppressor proteins (p53), cytochrome c release from mitochondria, activation of several death receptors involved in the trigger of apoptosis. Besides, various apoptosis proteins (IAPs) play a vital role in the induction and regulation of apoptosis.^{30,31} Apoptosis can occur via both the extrinsic pathway (death receptor-mediated pathway) and the intrinsic pathway (mitochondrial-mediated), and these pathways converge at the execution pathway of apoptosis.³ Death receptors, DR4, DR5 trigger the extrinsic pathway of apoptosis, up-regulation of pro-apoptotic protein (Bax) and down-regulation of an anti-apoptotic member of Bcl-2, which is essential for the activation of the intrinsic pathway of apoptosis.^{3,32} Extract of *Phaseolus vulgaris* (family Fabaceae) induces apoptosis in MCF-7 and MDA-MB-231 via up-regulating pro-apoptotic protein (Bax) and down-regulated anti-apoptotic protein (Bcl-2, Bcl-xL).³³ Similarly, fruit extract of *Momordica cochinchinensis* causes the induction of apoptosis in breast cancer (MCF-7) cells via the up-regulation of Bax and enhanced caspase 6, 8, and 9 activity.³⁴ Also, *Fragaria ananassa* (Strawberry) methanolic extract induced apoptosis by an intrinsic pathway in T-47D by the up-regulation of Bax, Bid, p73, and down-regulation of BCL-xL.³⁵ The aqueous extract fraction of *Oenocarpus bacaba* also induced apoptosis in MCF-7 cells by both extrinsic and intrinsic pathways through activation of caspases-6, -8, and -9.³⁶ Moreover,




Figure 1. Schematic illustration of plant-derived anti-breast cancer therapeutic agent




Figure 2. Anti-breast cancer plants and their therapeutic approaches

the methanolic fractions of *Scrophularia oxysepala* cause caspase-dependent apoptosis in MCF-7 cells.³⁷ The up-regulation of Bax induces apoptosis in MDA-MB-468 cells, treated with acetone and methanolic extracts of *Vatica diospyroides*.³⁸

Modulating cell cycle arrest

Cell cycle, remarkable role in cellular genomic integrity, and timely progression of cells.³⁹ Different phases such as (i) G1-phase (gap 1), (ii) S-phase (DNA synthesis), (iii) G2-phase (gap 2), and (iv) M-phase (mitosis) involve in the cell cycle. In S-phase, DNA synthesis and genome replication occur, required for the transmission of genetic information between generations. The M-phase causes segregation of genetic information, sister chromatids, and cell division. G1 is the gap between M and S phase, while G2 is the gap between S and M phase. These intervals (G1 and G2), essential to ensure that each phase is complete before moving to the next phase.^{40,41} Activation of cell cycle check-points usually occurs as a response to replication stress and DNA damage. The activation and inactivation of cyclin-dependent kinases and cyclins play a vital role in the cellular progression and cell cycle regulation.^{39,41} The methanolic extract prepared from *Allium atroviolaceum* (family Amaryllidaceae) induces apoptosis by modulating cell cycle arrest in caspase-dependent and p53-independent pathway in the breast cancer cell (MCF-7, MDA-MB-231).⁴² Similarly, ethanol extract of *Ganoderma lucidum* chipped fruiting bodies causes cell cycle arrest in MCF-7 cells by up-regulation of p21/Waf1 and down-regulation of cyclin D1.⁴³ The crude extracts of *Echinophora platyloba*, *Vernonia amygdalina*, *Morinda Citrifolia* induces apoptosis in MCF-7 and MDA-

MB-231 cell lines via G0/G1/S phase cell cycle arrest.^{12,44,45}

The diethyl-ether extract of *Artocarpus altilis* and hexane and methylene chloride fractions from roots of *Salvia chloroleuca* induced apoptosis and sub-G1 peak in T-47D and MCF-7 cells respectively.^{46,47} Also, ethyl acetate fraction from *Phaleria macrocarpa* (fruit) induce G0/G1 and G2/M cell cycle arrest in MDA-MB-231.⁴⁸

Inhibition of invasion and metastasis suppression

The conventional therapeutic approaches quite challenging, especially in metastasized cancer. The mechanism of metastasis involves invasion, intravasation, and extravasation. The process of invasion characterizes by the spreading of cancer cells to distant sites via the circulatory system. However, extravasation requires the penetration of cancer cells to the endothelium and the basement membrane. At the point of extravasation, cancer cells can grow at secondary focus.^{49,50} The matrix metalloproteinases (MMPs), critical proteins involved in metastasis of tumor cells. The inhibition or blocking of MMPs is an essential target in the suppression of metastatic potential. Other than MMPs, metastasis suppressor genes, MKK4 (mitogen-activated protein kinase 4), BRMS1 (breast cancer metastasis suppressor 1) and NM23 (non-metastatic gene 23) also play a remarkable role in the inhibition of metastasis.^{50,51} Similarly, modulation of uPA, uPAR, and TIMP expression also plays a vital role in the suppression of metastasis.⁵² The crude extracts of *Catharanthus roseus*, *Origanum majorana*, and *Brassica oleracea* possess anti-invasive and anti-metastatic activities in breast cancer cell line, MDA-MB-231. Anti-invasive and anti-metastatic activities via suppression of MMPs (MMP-2 and MMP-9) activities.⁵³⁻⁵⁵ Similarly, ethanol




Figure 3. Therapeutic targets of plant derived anti-breast cancer agents on breast cancer cells

extract of *Smilax china* causes suppression of metastasis via modulation of uPA, uPAR, and TIMP expression in MDA MB 231 cells.⁵⁶ Also, different fractions from stem of *Bauhinia unguolata* anti-metastatic decrease the activity of potential target of metastasis MMP-2.⁵⁷

Pro-survival signaling pathway

Several pro-survival signaling pathways were determining the fate of a cancer cell and mainly transduced by a complex net of signaling molecule cascade. Pro-survival signaling cascades, IP3K-PKB/Akt, and MAPK, activated by several cytokines and growth factors. The nuclear factor- κ B (NF κ B) plays an essential role in the regulation of inflammation and immune responses.⁵⁸ Blocking of these pro-survival signaling pathways has been widely studied, crucial for the treatment of breast cancer. The previous study shows that methanol extract of *Ixeris dentata* induced apoptosis in T-47D, MCF-7, SK-BR-3, and MDA-MB-231 via inhibiting Akt and NF- κ B signaling pathway.⁵⁹ Similarly, ethyl acetate fraction of *Euphorbia humifusa* causes inhibition of NF- κ B activity in MDA-MB-231 cell line.⁶⁰ Water- ethanol extract of *Astragalus membranaceus* induced apoptosis in MCF-7, SK-BR-3, and MDA-MB-231 through inhibition of PI3K, Akt and mTOR signaling pathways.⁶¹ Also, ethyl acetate extract from Roots of *Dillenia suffruticosa* induces apoptosis in MCF-7 via inhibition of AKT and ERK, and activation of JNK.⁶²

Other potential pathways

Various signal cascades induce cytotoxicity of breast

cancer cell lines via regulation of angiogenesis, autophagy, suppression of ERA expression, down-regulation of intracellular ROS generation, and mitochondrial membrane potential activated. The ethanol crude extract of *Salvia triloba* possesses angiogenesis activities in MCF-7 that is mediated by the inhibition of VEGF expression at both mRNA and protein levels.⁶³ Similarly, ethyl acetate fractions of *Eugenia jambolana* and *Musa paradisiaca* causes suppression of VEGF-induced angiogenesis in MCF-7 and MDA-MB-231 cells.⁶⁴ The extract of *Buxus sempervirens* induces autophagic cell death in MCF7, T47D, MCF10CA1a, and BT-20.⁶⁵ However, ROS mediated apoptosis in MCF-7 and MDA-MB-231 noticeable after treatment with chloroform fraction of *Tinospora cordifolia*.⁶⁶ Similarly, the hexane and methylene chloride fractions of *Salvia chloroleuca* also induce ROS-mediated pathway in MCF-7 cells.⁴⁶ Also, *Morinda citrifolia* (ethyl-acetate) extract downregulates intracellular ROS generation and mitochondrial membrane potential in MCF-7, and MDA-MB-231.⁴⁵ The *Acanthopanax sessiliflorus* (hexane fraction) causes mitochondria associated with both ROS dependent and independent pathways in MDA-MB-231 and MCF-7.⁶⁷

Phytochemicals in anti-breast cancer drug development

Plants possess different phytochemical compounds, and they classified based on the functional group, structures, and biosynthetic origins. Phytochemicals in medicinal plants include phenolics, flavonoids, alkaloids, terpenoids, carotenoids, saponins, steroids, and antioxidants induces cell death in MCF-7 cell lines.⁶⁸⁻⁸⁶ Among all

phytochemicals, phenolics are the most structurally diverse.⁸⁶ Here, we discuss potential phytochemicals as future anti-breast cancer therapeutic agents and drug development.

Phenolics

Phenolic compounds, widely occurring secondary metabolites isolated from plants and are most structurally diverse among all phytochemicals.⁸⁷ Plant-derived phenolic compounds classified as; (i) simple phenols, (ii) flavonoids, (iii) lignins, (iv) lignans, (v) tannins, (vi) xanthenes, and (vii) coumarins. Previous studies show that various phenolic compounds inhibit the initiation and progression of a variety of cancers by inducing cell cycle arrest, angiogenesis and apoptosis, modulating ROS levels and inhibiting oncogenic signaling cascades controlling cell proliferation.⁸⁸ The quercetin induces apoptosis in MCF-7, T47D, MDA-MB-453 and MDA-MB-231 cell line by up-regulation of Bax, down-regulation of Bcl-2 and activation of caspase-3. Similarly, quercetin also results in cell cycle arrest via modulation of Foxo3a activity in breast cancer.⁸⁹⁻⁹⁴ Interestingly, luteolin causes cytotoxicity in breast cancer cell line, MDA-MB-231 via suppression of epidermal growth factor receptor-mediated pathway IGF-1 pathway-dependent ER α .⁹⁵ Moreover, different phenolic acids such as ferulic acid, caffeic acid, and gallic acid also induces apoptosis in ER⁺ and ER⁻ breast cancer cell lines.^{88, 96,97}

Alkaloids

Plant-derived alkaloids possess oncogenesis suppression via modulating critical signaling pathways in human cancer. The paclitaxel possesses anticancer activity against breast cancer, ovarian cancer, prostate cancer, and lung cancer and is in clinical use.⁹⁸ Similarly, vinca alkaloids clinically used to treat human cancers. The vinca alkaloids (VA), from the *Madagascar periwinkle* plant (*Catharanthus roseus* G. Don), possess hypoglycaemic and cytotoxic properties. The VA considered as cancer fighters, second-most-certified class of cancer drugs. The four major vinca alkaloids include; (i) vincristine, (ii) vinblastine, (iii) vinorelbine, and (iv) vindesine are in clinical use. The vinflunine, a new synthetic vinca alkaloid, is used in treatment for carcinomas and other malignancies.⁹⁹ The vinca alkaloids interaction with tubulin protein, interfere with the assembly of microtubules, leads to cell division arrest in metaphase.

Similarly, vinflunine, a potent inhibitor of tubulin, causes hindrance in microtubule assembly and induces apoptosis. Moreover, vinflunine apoptosis mechanism involved activation of caspases 3 and 7 and c-Jun N-terminal kinase 1.¹⁰⁰ Other than vinca alkaloids; another alkaloid compound berberine induces cell cycle arrest and mitochondrial or intrinsic pathway in MCF-7 and MDA-MB-231 cells.^{101,102} Similarly, noscapine induces apoptosis in breast cancer cells via intrinsic and extrinsic

pathways by upregulation of Bax, downregulation of Bcl-2 and activation of caspases.¹⁰³⁻¹⁰⁵ The hirsutine causes cell death in MDA-MB-231 cells by activating the intrinsic pathway of apoptosis and targeting NF- κ B signaling pathway.^{106,107} Moreover, the treatment of MCF-7 cells by procaine decreases DNA methylation and RAR β 2 promoter methylation.¹⁰⁸

Terpenes

Terpenes or terpenoids classified based on the number of C5 units or cyclic structures present in the molecule.¹⁰⁹ The terpenoids can exert a broad spectrum of biological activities such as antioxidation, anti-inflammation, and anticancer activities. Numerous terpenoid compounds are known to possess anticancer potential in a variety of human cancers by causing inhibition of cancer cell proliferation and inducing apoptosis. Monoterpenoids such as D-limonene, have demonstrated antitumor and anticancer activities against breast cancer.¹¹⁰ Several diterpenoids also possess anticancer activity against breast cancers and are involved in the induction of apoptosis. These include triptolide, oridonin, and ponocidin.¹¹¹ The triptolide also possesses antiproliferative activity and down-regulates the expression of ER α in different breast cancer cell lines.¹¹²⁻¹¹⁴ The triterpenoids are close to steroids in structure and evoke apoptosis in a variety of cancer such as prostate and breast cancer. Different triterpenoids, like cucurbitacins, dammaranes, friedelanes, limonoids, lanostanes, lupanes, oleananes, tirucallanes, and ursanes, have been isolated from plants and studied for anticancer efficacy in breast cancer cells.¹¹⁵ Ursolic acid, a triterpene acid causes DNA fragmentation induced apoptosis in MCF-7 cells by downregulation of Bcl-2 and activation of caspase -3.^{116,117} Tetraterpenes also was known as carotenoids broadly categorized as acyclic tetraterpenoids and bicyclic tetraterpenoids. Carotenoids or tetraterpenoids such as lycopene and lutein are also known to possess anticancer activities in breast cancer cell lines.¹¹¹

Saponins

Saponins are natural glycosides widely distributed in plants classified into; triterpenoid, saponins, and steroid saponins.¹¹⁸ Saponins possess potential biological activities includes; anti-inflammatory, antiproliferative, immunomodulatory, and anticancer activities.¹¹⁹ Several saponins possess anticancer activities against various cancer cell lines.¹²⁰ For example, Avicin D, a triterpenoid glycoside compound, induces apoptosis in cutaneous T-cell lymphoma cells via downregulation of p-STAT-3 and bcl-2.¹²¹ Similarly, tubeimoside-1 exhibits anticancer effects via mitochondrial dysfunction and endoplasmic reticulum stress pathways in HeLa cells.¹²² The steroid saponins, degalactotigonin, and Polyphyllin D, possess cytotoxicity activity in ER⁺ human breast cancer cell line, MCF-7.^{123,124} Moreover, the triterpene saponins such as gummiferaoside B and C possess antiproliferative activity

in MDA-MB-435 cells.¹²⁵ Also, Avicins D, G induce apoptosis and cell cycle arrest in MDA-MB-435 cell line.¹²⁶ The phenolics, alkaloids, terpenoids and saponins derived from other sources also possess anticancer activity in different breast cancer cell lines.¹²⁷⁻¹⁴² Moreover, plant extracts, phytochemicals, and their potential mechanism of action against breast cancer are enlisted in Tables 1 and 2, respectively. Also, our various studies show the induction of cell death (apoptosis) in other cell lines.¹⁴³ The induction of cell death mainly due to presence of potential phytochemicals such as phenolics, saponins, terpenoids. The more screening and mechanistic studies

need to be done to fully explore the potent phytochemicals in field of cancer therapeutics.

Conclusion

Cancer is a complex disease, leading cause of death worldwide. Despite the development of many synthetic anticancer drugs, toxicity remains the main problem, which reduces the survival rate. Therefore, there is an increase in demand for alternative treatments. Amongst the alternative approaches, the natural product derived anticancer agents are a practical choice. The secondary metabolites, as potential anticancer agents

Table 1. Plant extracts and their potential mechanism of action against breast cancer

Plant name	Extract / Fraction	Part used	Target cell lines	Mechanism of cell death	References
<i>Allium atroviolaceum</i>	Methanolic Extract	Flower	MCF-7, MDA-MB-231	- Induces apoptosis - Modulating cell cycle arrest - Caspase-dependent and p53-independent Pathway	42
<i>Phaseolus vulgaris</i> (black turtle bean)	Extract	Seeds	MCF-7 and MDA-MB231	- Upregulation of Bax and downregulation of Bcl-2 and Bcl-xL - Activation of caspase -3/7	33
<i>Ganoderma lucidum</i>	Ethanol extract	Chipped fruiting bodies	MCF-7	- Induces cell cycle arrest and apoptosis - Up-regulation of p21/Waf1 and down-regulation of cyclin D1	43
<i>Echinophora Platyloba</i>	Methanol Extract	Leaves	MDA-MB-231	- Up-regulation of pro-apoptotic Bax protein - Induces apoptosis and cell cycle arrest at S-phase - Up-regulation of bax and p27 - Down-regulation of bcl-2	44
<i>Momordica cochinchinensis</i>	Aril Extract	Fruit	MCF-7	- Induces apoptosis - Increased bax enhanced caspase 6, 8 and 9 activity	34
<i>Morinda Citrifolia</i>	Ethyl-acetate extract	Fruit	MCF-7, MDA-MB-231	- Arrested the cell cycle in the G1/S phase in MCF-7 and G0/G1 phase in MDA-MB-231 cells - Downregulation of intracellular ROS generation and mitochondrial membrane potential - Cleavage of MCL-1 - downregulation of BCL-xL	45
<i>Fragaria ananassa</i> Strawberry	Methanolic extract	Fruit	T-47D	- Upregulation of expression of proapoptotic proteins such as BAX and BID - Upregulation of p73 - Activation of CASPASE 3 and CASPASE 9	35
<i>Vatica diospyroides</i>	Acetone and methanolic extracts	Fruit	MDA-MB-468	- Induces apoptosis - Up-regulation of Bax	38
<i>Oenocarpus bacaba</i>	Phenolic extract	Fruit	MCF-7	- Induces apoptosis - Caspases-6, -8 and -9 activated	36
<i>Averrhoa Bilimbi</i>	Methanolic extract	Fruit, Leaves	MCF-7	- Anticancer activity	68
<i>Carica papaya</i> L	Aqueous Extract	Leaves	MCF-7	- Anti-proliferation and Apoptosis induction	69
<i>Mimosa caesalpiniiifolia</i>	Ethanol extract	Leaf	MCF-7	- Induces apoptosis - DNA fragmentation	70
<i>Annona muricata</i>	Aqueous extract	Leaves	MCF-7, MDA-MB-231	- Induces apoptosis	71
<i>Acanthopanax sessiliflorus</i>	Hexane fraction	Stem bark	MDA-MB-231 and MCF-7	- Non-apoptotic cell death via mitochondria associated with both ROS dependent and independent pathways - Induce G0/G1 and G2/M cell cycle arrest - Activation of caspase -8,9 and 3	67
<i>Phaleria macrocarpa</i>	Ethyl acetate fraction	Fruit	MDA-MB-231	- Upregulation of Bax, Bid - cytochrome c, p21, p27, p53 and SMAC - Downregulation of Bcl-2, Bcl-w, XIAP and survivin	48
<i>Stryphnodendron adstringen</i>	Aqueous extract fraction	Leaves	MCF-7, MDA-MB-435	- Upregulation of Bax, caspase-9, active caspase-3, caspase-8, LC-3, and beclin-1 - Downregulation of Bcl-2	72
<i>Avicennia Marina</i>	Crude methanol extract and fraction	Leaves	MDA-MB 231	- DNA fragmentation - Decreased mRNA expression level of Bcl-2 and increased p53	73

Table 1. Continued

Plant name	Extract / Fraction	Part used	Target cell lines	Mechanism of cell death	References
<i>Salvia chloroleuca</i>	Hexane and methylene chloride fractions	Roots	MCF-7	- Induced a sub-G1 peak - DNA fragmentation - ROS-mediated pathway	46
<i>Scrophularia oxypetala</i>	Methanolic subfractions	Aerial parts	MCF-7	- Activation of caspase-3 - Downregulation of Bcl-2	36
<i>Artocarpus altilis</i>	Diethyl ether extract	Wood	T-47D	- Induced apoptosis and sub-G1 phase formation	47
<i>Piper crocatum</i>	Methanol extract	Leaves	T-47D	- Inhibition of p44/p42 phosphorylation	74
<i>Pistacia atlantica sub kurdica</i>	Methanol fraction	Fruits skin	T-47D	- Activation of caspase 3 - Poly ADP ribose polymerase (PARP) cleavage	75
<i>Vitex rotundifolia</i>	fraction	leave	MCF-7	- extrinsic and intrinsic pathway	76
<i>Vitex rotundifolia</i>	fraction	leave	T47D	- extrinsic and intrinsic pathway	77
<i>Aptos sp., marine</i>	fraction	whole	MCF-7	- DNA fragmentation	78
<i>Marine sponges</i>	Methanol extract	whole	MCF-7	- DNA fragmentation	79
<i>Vitex negundo</i>	Aqueous and Ethanolic extract	Leaves	MCF-7	- Induced apoptosis	80
<i>Jatropha curcas</i>	Ethanol extract	Root bark	MCF-7	- Inducing anoikis	81
<i>Vernonia amygdalina</i>	Ethanol extract	Leaves	MCF-7 and MDA-MB-231	- Induced apoptosis - G1/S phase cell cycle arrest - Caspase-dependent	45
<i>Strobilanthes crispa</i>	Hexane extract	Stem	MDA-MB-231	- Induced apoptosis	82
<i>Ixeris dentata</i>	Methanol extract	-	T-47D, MCF-7, SK-BR-3, and MDA-MB-231	- Induced apoptosis - via Akt-NF- κ B signaling	59
<i>Tinospora cordifolia</i>	Chloroform fraction	Stems	MCF-7 and MDA-MB-231	- ROS mediated apoptosis	66
<i>Smilax china</i>	Ethanol extract	Bark	MDA-MB-231	- Suppression of metastasis Modulation of uPA, uPAR and TIMP expression	56
<i>Bauhinia unguolata</i>	Different fractions	Stem	4T1	- Anti-tumor - Antimetastatic - decreasing the MMP-2 activity	57
<i>Nicotiana glauca</i>	Dichloromethane fraction	Stem	MCF-7	- Anti-Metastatic	83
<i>Euphorbia humifusa</i>	Ethyl acetate fraction	Whole plant	MDA-MB-231	- Inhibition of NF- κ B activity - Induced matrix metalloproteinase (MMP)-9 mRNA expression	60
<i>Withania coagulans</i>	Ethyl acetate	Aerial with fruit	MCF-7, MDA-MB-231	- Inhibited TNF- α induced NF κ B activity	84
<i>Astragalus membranaceus</i>	Water- ethanol extract	Roots	MCF-7, SK-BR-3 and MDA-MB-231	- Anti-proliferative - Induced apoptosis - Inhibition of PI3K/AKT/mTOR signaling pathway	61
<i>Dillenia suffruticosa</i>	Ethyl acetate extract	Roots	MCF-7	- Induces apoptosis via inhibition of AKT and ERK, and activation of JNK	62
<i>Catharanthus roseus</i>	Methanol extract	Leaves	MDA-MB-231	- Anti-invasive - Suppressed the MMP-2 and MMP-9 activity	53
<i>Forsythia koreana</i>	Methanol extract	Fruit and leaves	MDA-MB-231	- Suppressed invasion and MMPs activities - Inhibited the receptor activator of nuclear factor kappa-B	85
<i>Origanum majorana</i>	Ethanolic extract	Leaves	MDA-MB-231	- Anti-invasive and anti-metastatic - Downregulates the phosphorylation of I κ B, nuclear level of NF κ B and Nitric Oxide (NO) production	54
<i>Brassica oleracea</i>	Extract	-	MDA-MB-231	- Anti-invasive - Suppressed TPA-induced MMP-9 activity	55
<i>Salvia triloba</i>	Ethanolic crude extracts	Whole plant	MCF 7	- Antiangiogenesis - Inhibited the expression of VEGF at the mRNA and protein level	63
<i>Eugenia jambolana</i>	Ethyl acetate fractions	Seeds	MCF-7 and MDA-MB-231	- Suppression of VEGF-induced angiogenesis	64
<i>Musa paradisiaca</i>	Ethyl acetate fractions	Roots	MCF-7 and MDA-MB-231	- Suppression of VEGF-induced angiogenesis	64
<i>Buxus sempervirens</i>	Acetonic extract	Leaves and flowers	MCF7, T47D, MCF10CA1a and BT-20	- Induces apoptosis, Cell cycle arrest, autophagy	65

Table 2. Plant derived phytochemicals and their potential mechanism of action against breast cancer

Phytochemicals	Compound name, type	Target cell lines	Mechanism of cell death	References
Phenolics	Quercetin (Flavonoid)	MCF-7, T47D, MDA-MB-453, MDA-MB-231	- Induces apoptosis - Through suppression of Twist via p38 MAPK pathway - Increased Bax expression and decreased Bcl-2 expression - Increased cleaved caspase-3 and PARP expression - Cell cycle arrest through modulation of Foxo3a activity	89-94
	Casticin (flavonoid)	MCF-7, MDA-MB-231	- Induces apoptosis - Inhibiting the expression of forkhead box protein M1	127
	Luteolin (flavonoid)	MDA-MB-231, MCF-7	- Suppression of epidermal growth factor receptor-mediated pathway - IGF-1 pathway dependent ER α	95,128
	Ferulic acid (phenolic acid)	MDA-MB-231, T47D, MCF-7	- Induces apoptosis - Suppression of metastatic potential - Anti-proliferative	96,97,129
	Caffeic acid (phenolic acid)	T47D, MCF-7	- Anti-proliferative, - Induces apoptosis - Inhibition of NF κ B and activation of Fas	97,130
	Gallic acid (Phenolic acid)	MDA-MB-231	- Induces apoptosis	88
Alkaloids	Berberine	MCF-7, MDA-MB-231	- Inducing cell cycle arrest - Increasing levels of cytoplasmic cytochrome c, caspase-9, p53 and p27 - Cleavage of PARP - Decreasing levels of Bcl-2	101,102,131
	Noscapine	MCF-7, MDA-MB-231, T47D	- Activation of caspase-8 and caspase-9 - Upregulation of Bax, downregulation of Bcl-2 - Anti-neoplastic	103-105
	Pretazettine	MCF-7	- Anti-tumor activity	132
	Piperlongumine	MDA-MB-453, MCF-7, T-47D	- STAT3 Inhibitor	133
	Hirsutine	MDA-MB-453, MDA-MB-231, 4T1	- DNA damage response - NF- κ B and Akt pathways - Activation of the p38 MAPK pathway - Upregulation of Bax, downregulation of Bcl-2 - Activating caspase 9 and caspase 3	106-107, 134
	Procaine	MCF-7	- Decrease global DNA methylation - Decrease RAR β 2 promoter methylation	108
Terpenoid	Benzyl Isothiocyanate	MDA-MB-231, MCF-7, MDA-MB-468, BT-474, and BRI-JM04	- FoxO1-mediated autophagic cell death	135
	D-Limonene (Monoterpene)			
	Triptolide (Diterpene)	MDA-MB-435, MDA-MB-231, MCF7	- Anti-proliferative - Suppression of phospholipase D expression - Down-regulate the expression of ER α	112-114
	Ursolic acid (triterpene acid)	MCF-7	- Triggers apoptosis - DNA fragmentation - Downregulation of Bcl-2 - Activation of caspase -3	116-117
	Betulinic acid (triterpene)	MDAMB- 231, MDL13E, BT438, BT474, BT549, T47D	- Exhibited cytotoxicity - Induces apoptosis - Down-regulation of Bcl-2 and cyclin D1	136-137
	Lupeol (triterpene)	MDA-MB-231	- Suppressed the proliferation	138
Terpenoid	Lycopene (tetraterpenoids, carotenoids)	MCF 7, MDA-MB-231	- Trigger G2/M arrest and suppress Bcl-2 expression - Induce apoptosis	139
	Parthenolide	ZR-75-1, MDA-MB-231	- Inhibits HDAC1 increases global H3 acetylation, - Induces p21	140-141

Table 2. Continued

Phytochemicals	Compound name, type	Target cell lines	Mechanism of cell death	References
Saponins	Gummiferaoside B, C (triterpene saponins)	MDA-MB-435	- Anti-proliferative	125
	Degalactotigonin (steroid saponins)	MCF-7	- Cytotoxic	123
	Polyphyllin D (Steroid saponins)	MCF-7	- Cytotoxic	124
	Avicins D, G (triterpenoid saponins)	MDA-MB-435	- Apoptosis, - Cell cycle (G1) arrest	126
	Ginsenoside Rh ₂ (dammarane-type saponins)	MCF7	- Cytotoxic	142

with understandable anticancer mechanisms of action, leads to the development of novel therapeutic drugs. Additionally, the plant extracts are an excellent source of lead compounds. The isolated lead compound can either used or undergoes some structural modifications to increase the effectiveness in term of their pharmacological potential.

Ethical Issues

Not applicable.

Conflict of Interest

The authors have no conflict of interest.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68(6):394-424. doi: 10.3322/caac.21492
- Nithya M, Ambikapathy V, Panneerselvam A, Thajuddin N. Anti tumour activity of different extracts of *Ganoderma lucidum* (Curt.: Fr.) P. karst. *World J Pharm Res* 2014;3(4): 2204-14.
- Jan R, Chaudhry GE. Understanding apoptosis and apoptotic pathways targeted cancer therapeutics. *Adv Pharm Bull* 2019;9(2):205-18. doi: 10.15171/apb.2019.024
- Tonini T, Rossi F, Claudio PP. Molecular basis of angiogenesis and cancer. *Oncogene* 2003;22(42):6549-56. doi: 10.1038/sj.onc.1206816
- Rice H, Bryant S, Handley C, Hall M. Oncogenes and tumor suppressor genes: an essential building block of cancer. *Chemist* 2014;87(2):15-8.
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66(1):7-30. doi: 10.3322/caac.21332
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA Cancer J Clin* 2017;67(1):7-30. doi: 10.3322/caac.21387
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68(1):7-30. doi: 10.3322/caac.21442
- Arpino G, Bardou VJ, Clark GM, Elledge RM. Infiltrating lobular carcinoma of the breast: tumor characteristics and clinical outcome. *Breast Cancer Res* 2004;6(3):R149-56. doi: 10.1186/bcr767
- Raghav K, Morris V, Tang C, Morelli P, Amin HM, Chen K, et al. MET amplification in metastatic colorectal cancer: an acquired response to EGFR inhibition, not a de novo phenomenon. *Oncotarget* 2016;7(34):54627-31. doi: 10.18632/oncotarget.10559
- Karakas C. Paget's disease of the breast. *J Carcinog* 2011;10:31. doi: 10.4103/1477-3163.90676
- Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, et al. A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 2010;141(1):69-80. doi: 10.1016/j.cell.2010.02.027
- Virnig BA, Tuttle TM, Shamlivan T, Kane RL. Ductal carcinoma in situ of the breast: a systematic review of incidence, treatment, and outcomes. *J Natl Cancer Inst* 2010;102(3):170-8. doi: 10.1093/jnci/djp482
- Roberts N, Runk DM. Aggressive malignant phyllodes tumor. *Int J Surg Case Rep* 2015;8c:161-5. doi: 10.1016/j.ijscr.2014.12.041
- Mishra SP, Tiwary SK, Mishra M, Khanna AK. Phyllodes tumor of breast: a review article. *ISRN Surg* 2013;2013:361469. doi: 10.1155/2013/361469
- Taherian-Fard A, Srihari S, Ragan MA. Breast cancer classification: linking molecular mechanisms to disease prognosis. *Brief Bioinform* 2015;16(3):461-74. doi: 10.1093/bib/bbu020
- Dai X, Cheng H, Bai Z, Li J. Breast cancer cell line classification and its relevance with breast tumor subtyping. *J Cancer* 2017;8(16):3131-41. doi: 10.7150/jca.18457
- Fenga C. Occupational exposure and risk of breast cancer. *Biomed Rep* 2016;4(3):282-92. doi: 10.3892/br.2016.575
- McDonald ES, Clark AS, Tchou J, Zhang P, Freedman GM. Clinical diagnosis and management of breast cancer. *J Nucl Med* 2016;57 Suppl 1:9S-16S. doi: 10.2967/jnumed.115.157834
- Wang L. Early diagnosis of breast cancer. *Sensors (Basel)* 2017;17(7):1572. doi: 10.3390/s17071572
- Roy PS, Saikia BJ. Cancer and cure: a critical analysis. *Indian J Cancer* 2016;53(3):441-2. doi: 10.4103/0019-509x.200658
- van Uden DJ, van Laarhoven HW, Westenberg AH, de Wilt JH, Blanken-Peeters CF. Inflammatory breast cancer: an overview. *Crit Rev Oncol Hematol* 2015;93(2):116-26. doi: 10.1016/j.critrevonc.2014.09.003
- Puhalla S, Bhattacharya S, Davidson NE. Hormonal therapy in breast cancer: a model disease for the personalization of

- cancer care. *Mol Oncol* 2012;6(2):222-36. doi: 10.1016/j.molonc.2012.02.003
24. Jones ME, Schoemaker MJ, Wright L, McFadden E, Griffin J, Thomas D, et al. Menopausal hormone therapy and breast cancer: what is the true size of the increased risk? *Br J Cancer* 2016;115(5):607-15. doi: 10.1038/bjc.2016.231
 25. Valli M, Pivatto M, Danuello A, Castro-Gamboa I, Silva DH, Cavaleiro AJ, et al. Tropical biodiversity: has it been a potential source of secondary metabolites useful for medicinal chemistry? *Quím Nova* 2012;35(11):2278-87. doi: 10.1590/s0100-40422012001100036
 26. Habli Z, Toumeh G, Fatfat M, Rahal ON, Gali-Muhtasib H. Emerging cytotoxic alkaloids in the battle against cancer: overview of molecular mechanisms. *Molecules* 2017;22(2):250. doi: 10.3390/molecules22020250
 27. Desai AG, Qazi GN, Ganju RK, El-Tamer M, Singh J, Saxena AK, et al. Medicinal plants and cancer chemoprevention. *Curr Drug Metab* 2008;9(7):581-91. doi: 10.2174/138920008785821657
 28. Nikolettou V, Markaki M, Palikaras K, Tavernarakis N. Crosstalk between apoptosis, necrosis and autophagy. *Biochim Biophys Acta* 2013;1833(12):3448-59. doi: 10.1016/j.bbamcr.2013.06.001
 29. Proskuryakov SY, Gabai VL, Konoplyannikov AG. Necrosis is an active and controlled form of programmed cell death. *Biochemistry (Mosc)* 2002;67(4):387-408. doi: 10.1023/a:1015289521275
 30. Ouyang W, Liao W, Luo CT, Yin N, Huse M, Kim MV, et al. Novel Foxo1-dependent transcriptional programs control T(reg) cell function. *Nature* 2012;491(7425):554-9. doi: 10.1038/nature11581
 31. Elmore S. Apoptosis: a review of programmed cell death. *Toxicol Pathol* 2007;35(4):495-516. doi: 10.1080/01926230701320337
 32. Kumar S, Sharma VK, Yadav S, Dey S. Antiproliferative and apoptotic effects of black turtle bean extracts on human breast cancer cell line through extrinsic and intrinsic pathway. *Chem Cent J* 2017;11(1):56. doi: 10.1186/s13065-017-0281-5
 33. Petchsak P, Sripanidkulchai B. *Momordica cochinchinensis* aril extract induced apoptosis in human MCF-7 breast cancer cells. *Asian Pac J Cancer Prev* 2015;16(13):5507-13. doi: 10.7314/apjcp.2015.16.13.5507
 34. Somasagara RR, Hegde M, Chiruvella KK, Musini A, Choudhary B, Raghavan SC. Extracts of strawberry fruits induce intrinsic pathway of apoptosis in breast cancer cells and inhibits tumor progression in mice. *PLoS One* 2012;7(10):e47021. doi: 10.1371/journal.pone.0047021
 35. Abadio Finco FDB, Kloss L, Graeve L. Bacaba (*Oenocarpus bacaba*) phenolic extract induces apoptosis in the MCF-7 breast cancer cell line via the mitochondria-dependent pathway. *NFS J* 2016;5:5-15. doi: 10.1016/j.nfs.2016.11.001
 36. Orangi M, Pasdaran A, Shanebandi D, Kazemi T, Yousefi B, Hosseini BA, et al. Cytotoxic and apoptotic activities of methanolic subfractions of *Scrophularia oxyssepala* against human breast cancer cell line. *Evid Based Complement Alternat Med* 2016;2016:8540640. doi: 10.1155/2016/8540640
 37. Srisawat T, Sukpondma Y, Chimplee S, Kanokwiroon K, Tedasen A, Graidist P. Extracts from *Vatica diospyroides* type SS fruit show low dose activity against MDA-MB-468 breast cancer cell-line via apoptotic action. *Biomed Res Int* 2014;2014:479602. doi: 10.1155/2014/479602
 38. Azzopardi M, Farrugia G, Balzan R. Cell-cycle involvement in autophagy and apoptosis in yeast. *Mech Ageing Dev* 2017;161(Pt B):211-24. doi: 10.1016/j.mad.2016.07.006
 39. Israels ED, Israels LG. The cell cycle. *Oncologist* 2000;5(6):510-3. doi: 10.1634/theoncologist.5-6-510
 40. Pucci B, Kasten M, Giordano A. Cell cycle and apoptosis. *Neoplasia* 2000;2(4):291-9. doi: 10.1038/sj.neo.7900101
 41. Khazaei S, Abdul Hamid R, Ramachandran V, Mohd Esa N, Pandurangan AK, Danazadeh F, et al. Cytotoxicity and proapoptotic effects of *Allium atroviolaceum* flower extract by modulating cell cycle arrest and caspase-dependent and p53-independent pathway in breast cancer cell lines. *Evid Based Complement Alternat Med* 2017;2017:1468957. doi: 10.1155/2017/1468957
 42. Hu H, Ahn NS, Yang X, Lee YS, Kang KS. *Ganoderma lucidum* extract induces cell cycle arrest and apoptosis in MCF-7 human breast cancer cell. *Int J Cancer* 2002;102(3):250-3. doi: 10.1002/ijc.10707
 43. Birjandian E, Motamed N, Yassa N. Crude methanol extract of *Echinophora platyloba* induces apoptosis and cell cycle arrest at S-phase in human breast cancer cells. *Iran J Pharm Res* 2018;17(1):307-16.
 44. Sharma K, Pachauri SD, Khandelwal K, Ahmad H, Arya A, Biala P, et al. Anticancer effects of extracts from the fruit of *Morinda citrifolia* (Noni) in breast cancer cell lines. *Drug Res (Stuttg)* 2016;66(3):141-7. doi: 10.1055/s-0035-1555804
 45. Wong FC, Woo CC, Hsu A, Tan BK. The anti-cancer activities of *Vernonia amygdalina* extract in human breast cancer cell lines are mediated through caspase-dependent and p53-independent pathways. *PLoS One* 2013;8(10):e78021. doi: 10.1371/journal.pone.0078021
 46. Tayarani-Najaran Z, Asili J, Aioubi E, Emami SA. Growth inhibition and apoptosis induction of *Salvia chloroleuca* on MCF-7 breast cancer cell line. *Iran J Pharm Res* 2013;12(4):789-99.
 47. Arung ET, Wicaksono BD, Handoko YA, Kusuma IW, Yulia D, Sandra F. Anti-cancer properties of diethylether extract of wood from sukun (*Artocarpus altilis*) in human breast cancer (T47D) cells. *Trop J Pharm Res* 2009;8(4):317-24. doi: 10.4314/tjpr.v8i4.45223
 48. Kavitha N, Ein Oon C, Chen Y, Kanwar JR, Sasidharan S. *Phaleria macrocarpa* (Boerl.) fruit induce G(0)/G(1) and G(2)/M cell cycle arrest and apoptosis through mitochondria-mediated pathway in MDA-MB-231 human breast cancer cell. *J Ethnopharmacol* 2017;201:42-55. doi: 10.1016/j.jep.2017.02.041
 49. Carr I, Orr FW. Invasion and metastasis. *Can Med Assoc J* 1983;128(10):1164-7.
 50. Martin TA, Ye L, Sanders AJ, Lane J, Jiang WG. Cancer invasion and metastasis: molecular and cellular perspective. In: Blagosklonny MV, Pardee AB, eds. *Madame Curie Bioscience Database*. Austin, TX: Landes Bioscience; 2000-2013.
 51. Decock J, Thirkettle S, Wagstaff L, Edwards DR. Matrix metalloproteinases: protective roles in cancer. *J Cell Mol Med* 2011;15(6):1254-65. doi: 10.1111/j.1582-4934.2011.01302.x
 52. Cox G, Steward WP, O'Byrne KJ. The plasmin cascade and matrix metalloproteinases in non-small cell lung cancer. *Thorax* 1999;54(2):169-79. doi: 10.1136/thx.54.2.169

53. Eltayeb NM, Ng SY, Ismail Z, Salhimi SM. Anti-invasive effect of *Catharanthus roseus* extract on highly metastatic human breast cancer MDA-MB-231 cells. *J Teknol* 2016;78(11-3):35-40. doi: 10.11113/jt.v78.9870
54. Al Dhaheri Y, Attoub S, Arafat K, Abuqamar S, Viallet J, Saleh A, et al. Anti-metastatic and anti-tumor growth effects of *Origanum majorana* on highly metastatic human breast cancer cells: inhibition of NF κ B signaling and reduction of nitric oxide production. *PLoS One* 2013;8(7):e68808. doi: 10.1371/journal.pone.0068808
55. Rose P, Huang Q, Ong CN, Whiteman M. Broccoli and watercress suppress matrix metalloproteinase-9 activity and invasiveness of human MDA-MB-231 breast cancer cells. *Toxicol Appl Pharmacol* 2005;209(2):105-13. doi: 10.1016/j.taap.2005.04.010
56. Nho KJ, Chun JM, Kim HK. Anti-metastatic effect of *Smilax china* L. extract on MDA-MB-231 cells. *Mol Med Rep* 2015;11(1):499-502. doi: 10.3892/mmr.2014.2698
57. Santos KM, Gomes INF, Romao W, Ribeiro RIMA, Silva-Oliveira RJ, Pinto FE, et al. *Bauhinia* stem extracts as a possible new treatment for breast cancer and metastasis: inhibition of migration, invasion and of the activity of matrix metalloproteinases. *J Biotechnol Biomater* 2017;7(6 Suppl):59. doi: 10.4172/2155-952x-c1-085
58. Klener P Jr, Andera L, Klener P, Necas E, Zivný J. Cell death signalling pathways in the pathogenesis and therapy of haematologic malignancies: overview of apoptotic pathways. *Folia Biol (Praha)* 2006;52(1-2):34-44.
59. Shin SA, Lee HN, Choo GS, Kim HJ, Che JH, Jung JY. *Ixeris dentata* (Thunb. Ex Thunb.) Nakai extract inhibits proliferation and induces apoptosis in breast cancer cells through Akt/NF- κ B pathways. *Int J Mol Sci* 2017;18(2). doi: 10.3390/ijms18020275
60. Shin WS, Han J, Kumar R, Lee GG, Sessler JL, Kim JH, et al. Programmed activation of cancer cell apoptosis: a tumor-targeted phototherapeutic topoisomerase I inhibitor. *Sci Rep* 2016;6:29018. doi: 10.1038/srep29018
61. Zhou R, Chen H, Chen J, Chen X, Wen Y, Xu L. Extract from *Astragalus membranaceus* inhibit breast cancer cells proliferation via PI3K/AKT/mTOR signaling pathway. *BMC Complement Altern Med* 2018;18(1):83. doi: 10.1186/s12906-018-2148-2
62. Tor YS, Yazan LS, Foo JB, Wibowo A, Ismail N, Cheah YK, et al. Induction of apoptosis in MCF-7 cells via oxidative stress generation, mitochondria-dependent and caspase-independent pathway by ethyl acetate extract of *Dillenia suffruticosa* and its chemical profile. *PLoS One* 2015;10(6):e0127441. doi: 10.1371/journal.pone.0127441
63. Zihlif M, Afifi F, Abu-Dahab R, Abdul Majid AM, Sumrein H, Saleh MM, et al. The antiangiogenic activities of ethanolic crude extracts of four *Salvia* species. *BMC Complement Altern Med* 2013;13:358. doi: 10.1186/1472-6882-13-358
64. Harsha Raj M, Ghosh D, Banerjee R, Salimath BP. Suppression of VEGF-induced angiogenesis and tumor growth by *Eugenia jambolana*, *Musa paradisiaca*, and *Coccinia indica* extracts. *Pharm Biol* 2017;55(1):1489-99. doi: 10.1080/13880209.2017.1307422
65. Ait-Mohamed O, Battisti V, Joliot V, Fritsch L, Pontis J, Medjkane S, et al. Acetonic extract of *Buxus sempervirens* induces cell cycle arrest, apoptosis and autophagy in breast cancer cells. *PLoS One* 2011;6(9):e24537. doi: 10.1371/journal.pone.0024537
66. Ansari JA, Rastogi N, Ahmad MK, Mahdi AA, Khan AR, Thakur R, et al. ROS mediated pro-apoptotic effects of *Tinospora cordifolia* on breast cancer cells. *Front Biosci (Elite Ed)* 2017;9:89-100. doi: 10.2741/e788
67. Thamizhiniyan V, Young-Woong C, Young-Kyoon K. The cytotoxic nature of *Acanthopanax sessiliflorus* stem bark extracts in human breast cancer cells. *Saudi J Biol Sci* 2015;22(6):752-9. doi: 10.1016/j.sjbs.2015.04.004
68. Nair MS, Soren K, Singh V, Boro B. Anticancer activity of fruit and leaf extracts of *Averrhoa bilimbi* on MCF-7 human breast cancer cell lines: a preliminary study. *Austin J Pharmacol Ther* 2016;4(2):1082.
69. Zuhrotun Nisa F, Astuti M, Murdiati A, Mubarika Haryana S. Anti-proliferation and apoptosis induction of aqueous leaf extract of *Carica papaya* L. on human breast cancer cells MCF-7. *Pak J Biol Sci* 2017;20(1):36-41. doi: 10.3923/pjbs.2017.36.41
70. Silva MJD, Carvalho AJS, Rocha CQ, Vilegas W, Silva MA, Gouvêa CMCP. Ethanolic extract of *Mimosa caesalpinhiifolia* leaves: chemical characterization and cytotoxic effect on human breast cancer MCF-7 cell line. *S Afr J Bot* 2014;93:64-9. doi: 10.1016/j.sajb.2014.03.011
71. Kim JY, Dao TTP, Song K, Park SB, Jang H, Park MK, et al. *Annona muricata* leaf extract triggered intrinsic apoptotic pathway to attenuate cancerous features of triple negative breast cancer MDA-MB-231 cells. *Evid Based Complement Alternat Med* 2018;2018:7972916. doi: 10.1155/2018/7972916
72. Sabino APL, Eustáquio LMS, Miranda ACF, Biojone C, Mariosa TN, Gouvêa C. *Stryphnodendron adstringens* ("Barbatimão") leaf fraction: chemical characterization, antioxidant activity, and cytotoxicity towards human breast cancer cell lines. *Appl Biochem Biotechnol* 2018;184(4):1375-89. doi: 10.1007/s12010-017-2632-z
73. Momtazi-Borojeni AA, Behbahani M, Sadeghi-Aliabadi H. Antiproliferative activity and apoptosis induction of crude extract and fractions of *Avicennia marina*. *Iran J Basic Med Sci* 2013;16(11):1203-8.
74. Wicaksono BD, Handoko YA, Arung ET, Kusuma IW, Yulia D, Pancaputra AN, et al. Antiproliferative effect of the methanol extract of *Piper crocatum* Ruiz & Pav leaves on human breast (T47D) cells in-vitro. *Trop J Pharm Res* 2009;8(4):345-52. doi: 10.4314/tjpr.v8i4.45229
75. Rezaei PF, Fouladdel S, Cristofanon S, Ghaffari SM, Amin GR, Azizi E. Comparative cellular and molecular analysis of cytotoxicity and apoptosis induction by doxorubicin and Baneh in human breast cancer T47D cells. *Cytotechnology* 2011;63(5):503-12. doi: 10.1007/s10616-011-9373-6
76. Chaudhry GE, Jan R, Mohammad H, Muhammad TST. *Vitex rotundifolia* fractions induce apoptosis in human breast cancer cell line, MCF-7, via extrinsic and intrinsic pathways. *Res Pharm Sci* 2019;14(3):273-85. doi: 10.4103/1735-5362.258496
77. Chaudhry GE, Jan R, Zafar MN, Mohammad H, Muhammad TST. *Vitex rotundifolia* fractions induced apoptosis in human breast cancer T-47D cell line via activation of extrinsic and intrinsic pathway. *Asian Pac J Cancer Prev* 2019;20(12):3555-62. doi: 10.31557/apjcp.2019.20.12.3555
78. Chaudhry GE, Kassim MNI, Ismail N, Mohammad H, Sung YY, Muhammad TST. Induction of apoptosis by *Aptos* sp.,

- fractions in human breast cancer cell line, MCF-7. *Int J Res Pharm Sci* 2018;9(2):328-37.
79. Hudaya T, Chaudhry GE, Taib M, Ismail N, Mohammad TST. Methanol extracts of four selected marine sponges induce apoptosis in human breast cancer cell line, MCF-7. *Int J Res Pharm Sci* 2017;8(4):667-75.
80. Arulvasu C, Prabhu D, Manikandan R, Srinivasan P, Dinesh D, Babu G, et al. Induction of apoptosis by the aqueous and ethanolic leaf extract of *Vitex negundo* L. in MCF-7 human breast cancer cells. *Int J Drug Discov* 2010;2(1):1-7. doi: 10.9735/0975-4423.2.1.1-7
81. Engel N, Falodun A, Kühn J, Kragl U, Langer P, Nebe B. Pro-apoptotic and anti-adhesive effects of four African plant extracts on the breast cancer cell line MCF-7. *BMC Complement Altern Med* 2014;14:334. doi: 10.1186/1472-6882-14-334
82. Koh RY, Lim FP, Ling LSY, Ng CPL, Liew SF, Yew MY, et al. Anticancer mechanisms of *Strobilanthes crisper* Blume hexane extract on liver and breast cancer cell lines. *Oncol Lett* 2017;14(4):4957-64. doi: 10.3892/ol.2017.6821
83. Tabana YM, Dahham SS, Ahmed Hassan LE, Al-Mansoub MA, Taleb-Agha M, Abdul Majid AM. In vitro anti-metastatic and antioxidant activity of *Nicotiana glauca* fraction against breast cancer cells. *Adv Biol Res* 2015;9(2):95-102. doi: 10.5829/idosi.abr.2015.9.2.9521
84. Ihsan-ul-Haq, Mirza B, Kondratyuk TP, Park EJ, Burns BE, Marler LE, et al. Preliminary evaluation for cancer chemopreventive and cytotoxic potential of naturally growing ethnobotanically selected plants of Pakistan. *Pharm Biol* 2013;51(3):316-28. doi: 10.3109/13880209.2012.728612
85. Kim YL, Lee SK, Park KK, Chung WY. The inhibitory effects of *Forsythia koreana* extracts on the metastatic ability of breast cancer cells and bone resorption by osteoclasts. *J Cancer Prev* 2016;21(2):88-94. doi: 10.15430/jcp.2016.21.2.88
86. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *J Pharmacogn Phytochem* 2013;1(6):168-82.
87. Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by-products: antioxidant activity, occurrence, and potential uses. *Food Chem* 2006;99(1):191-203. doi: 10.1016/j.foodchem.2005.07.042
88. Anantharaju PG, Gowda PC, Vimalambike MG, Madhunapantula SV. An overview on the role of dietary phenolics for the treatment of cancers. *Nutr J* 2016;15(1):99. doi: 10.1186/s12937-016-0217-2
89. Somasagara RR, Hegde M, Chiruvella KK, Musini A, Choudhary B, Raghavan SC. Extracts of strawberry fruits induce intrinsic pathway of apoptosis in breast cancer cells and inhibits tumor progression in mice. *PLoS One* 2012;7(10):e47021. doi: 10.1371/journal.pone.0047021
90. Duo J, Ying GG, Wang GW, Zhang L. Quercetin inhibits human breast cancer cell proliferation and induces apoptosis via Bcl-2 and Bax regulation. *Mol Med Rep* 2012;5(6):1453-6. doi: 10.3892/mmr.2012.845
91. Rahimifard M, Sadeghi F, Asadi-Samani M, Nejati-Koshki K. Effect of quercetin on secretion and gene expression of leptin in breast cancer. *J Tradit Chin Med* 2017;37(3):321-5.
92. Choi EJ, Bae SM, Ahn WS. Antiproliferative effects of quercetin through cell cycle arrest and apoptosis in human breast cancer MDA-MB-453 cells. *Arch Pharm Res* 2008;31(10):1281-5. doi: 10.1007/s12272-001-2107-0
93. Chien SY, Wu YC, Chung JG, Yang JS, Lu HF, Tsou MF, et al. Quercetin-induced apoptosis acts through mitochondrial- and caspase-3-dependent pathways in human breast cancer MDA-MB-231 cells. *Hum Exp Toxicol* 2009;28(8):493-503. doi: 10.1177/0960327109107002
94. Nguyen LT, Lee YH, Sharma AR, Park JB, Jagga S, Sharma G, et al. Quercetin induces apoptosis and cell cycle arrest in triple-negative breast cancer cells through modulation of Foxo3a activity. *Korean J Physiol Pharmacol* 2017;21(2):205-13. doi: 10.4196/kjpp.2017.21.2.205
95. Lee EJ, Oh SY, Sung MK. Luteolin exerts anti-tumor activity through the suppression of epidermal growth factor receptor-mediated pathway in MDA-MB-231 ER-negative breast cancer cells. *Food Chem Toxicol* 2012;50(11):4136-43. doi: 10.1016/j.fct.2012.08.025
96. Zhang X, Lin D, Jiang R, Li H, Wan J, Li H. Ferulic acid exerts antitumor activity and inhibits metastasis in breast cancer cells by regulating epithelial to mesenchymal transition. *Oncol Rep* 2016;36(1):271-8. doi: 10.3892/or.2016.4804
97. Kampa M, Alexaki VI, Notas G, Nifli AP, Nistikaki A, Hatzoglou A, et al. Antiproliferative and apoptotic effects of selective phenolic acids on T47D human breast cancer cells: potential mechanisms of action. *Breast Cancer Res* 2004;6(2):R63-74. doi: 10.1186/bcr752
98. Habli Z, Toumeh G, Fatfat M, Rahal ON, Gali-Muhtasib H. Emerging cytotoxic alkaloids in the battle against cancer: overview of molecular mechanisms. *Molecules* 2017;22(2):250. doi: 10.3390/molecules22020250
99. Moudi M, Go R, Yien CY, Nazre M. Vinca alkaloids. *Int J Prev Med* 2013;4(11):1231-5.
100. Ali R, Mirza Z, Ashraf GM, Kamal MA, Ansari SA, Damanhour GA, et al. New anticancer agents: recent developments in tumor therapy. *Anticancer Res* 2012;32(7):2999-3005.
101. Kim JB, Yu JH, Ko E, Lee KW, Song AK, Park SY, et al. The alkaloid Berberine inhibits the growth of Anoikis-resistant MCF-7 and MDA-MB-231 breast cancer cell lines by inducing cell cycle arrest. *Phytomedicine* 2010;17(6):436-40. doi: 10.1016/j.phymed.2009.08.012
102. Patil JB, Kim J, Jayaprakasha GK. Berberine induces apoptosis in breast cancer cells (MCF-7) through mitochondrial-dependent pathway. *Eur J Pharmacol* 2010;645(1-3):70-8. doi: 10.1016/j.ejphar.2010.07.037
103. Quisbert-Valenzuela EO, Calaf GM. Apoptotic effect of noscapiene in breast cancer cell lines. *Int J Oncol* 2016;48(6):2666-74. doi: 10.3892/ijo.2016.3476
104. Afzali M, Ghaeli P, Khanavi M, Parsa M, Montazeri H, Ghahremani MH, et al. Non-addictive opium alkaloids selectively induce apoptosis in cancer cells compared to normal cells. *Daru* 2015;23(1):16. doi: 10.1186/s40199-015-0101-1
105. DeBono A, Capuano B, Scammells PJ. Progress toward the development of noscapiene and derivatives as anticancer agents. *J Med Chem* 2015;58(15):5699-727. doi: 10.1021/jm501180v
106. Huang QW, Zhai NN, Huang T, Li DM. [Hirsutine induces apoptosis of human breast cancer MDA-MB-231 cells through mitochondrial pathway]. *Sheng Li Xue Bao* 2018;70(1):40-6.

107. Lou C, Takahashi K, Irimura T, Saiki I, Hayakawa Y. Identification of Hirsutine as an anti-metastatic phytochemical by targeting NF- κ B activation. *Int J Oncol* 2014;45(5):2085-91. doi: 10.3892/ijo.2014.2624
108. Villar-Garea A, Fraga MF, Espada J, Esteller M. Procaine is a DNA-demethylating agent with growth-inhibitory effects in human cancer cells. *Cancer Res* 2003;63(16):4984-9.
109. Rajeshwari CU, Shobha RI, Andallu B. Phytochemicals in diet and human health with special reference to polyphenols. *Ann Phytomed* 2014;3(2):4-25.
110. Crowell PL. Monoterpenes in breast cancer chemoprevention. *Breast Cancer Res Treat* 1997;46(2-3):191-7. doi: 10.1023/a:1005939806591
111. Yang H, Dou QP. Targeting apoptosis pathway with natural terpenoids: implications for treatment of breast and prostate cancer. *Curr Drug Targets* 2010;11(6):733-44. doi: 10.2174/138945010791170842
112. Yang S, Chen J, Guo Z, Xu XM, Wang L, Pei XF, et al. Triptolide inhibits the growth and metastasis of solid tumors. *Mol Cancer Ther* 2003;2(1):65-72.
113. Kang DW, Lee JY, Oh DH, Park SY, Woo TM, Kim MK, et al. Triptolide-induced suppression of phospholipase D expression inhibits proliferation of MDA-MB-231 breast cancer cells. *Exp Mol Med* 2009;41(9):678-85. doi: 10.3858/emmm.2009.41.9.074
114. Liang M, Fu J. Triptolide inhibits interferon-gamma-induced programmed death-1-ligand 1 surface expression in breast cancer cells. *Cancer Lett* 2008;270(2):337-41. doi: 10.1016/j.canlet.2008.05.025
115. Bishayee A, Ahmed S, Brankov N, Perloff M. Triterpenoids as potential agents for the chemoprevention and therapy of breast cancer. *Front Biosci (Landmark Ed)* 2011;16:980-96. doi: 10.2741/3730
116. Zhang GP, Lu YY, Lv JC, Ou HJ. [Effect of ursolic acid on caspase-3 and PARP expression of human MCF-7 cells]. *Zhongguo Zhong Yao Za Zhi* 2006;31(2):141-4.
117. Kassi E, Sourlingas TG, Spiliotiaki M, Papoutsis Z, Pratsinis H, Aliogiannis N, et al. Ursolic acid triggers apoptosis and Bcl-2 downregulation in MCF-7 breast cancer cells. *Cancer Invest* 2009;27(7):723-33. doi: 10.1080/07357900802672712
118. Cheok CY, Salman HA, Sulaiman R. Extraction and quantification of saponins: a review. *Food Res Int* 2014;59:16-40. doi: 10.1016/j.foodres.2014.01.057
119. Podolak I, Galanty A, Sobolewska D. Saponins as cytotoxic agents: a review. *Phytochem Rev* 2010;9(3):425-74. doi: 10.1007/s11101-010-9183-z
120. Yokosuka A, Jitsuno M, Yui S, Yamazaki M, Mimaki Y. Steroidal glycosides from *Agave utahensis* and their cytotoxic activity. *J Nat Prod* 2009;72(8):1399-404. doi: 10.1021/np900168d
121. Zhang C, Li B, Gaikwad AS, Haridas V, Xu Z, Guterman JU, et al. Avicin D selectively induces apoptosis and downregulates p-STAT-3, bcl-2, and survivin in cutaneous T-cell lymphoma cells. *J Invest Dermatol* 2008;128(11):2728-35. doi: 10.1038/jid.2008.138
122. Xu Y, Chiu JF, He QY, Chen F. Tubeimoside-1 exerts cytotoxicity in HeLa cells through mitochondrial dysfunction and endoplasmic reticulum stress pathways. *J Proteome Res* 2009;8(3):1585-93. doi: 10.1021/pr801001j
123. Zhou X, He X, Wang G, Gao H, Zhou G, Ye W, et al. Steroidal saponins from *Solanum nigrum*. *J Nat Prod* 2006;69(8):1158-63. doi: 10.1021/np060091z
124. Siu FM, Ma DL, Cheung YW, Lok CN, Yan K, Yang Z, et al. Proteomic and transcriptomic study on the action of a cytotoxic saponin (Polyphyllin D): induction of endoplasmic reticulum stress and mitochondria-mediated apoptotic pathways. *Proteomics* 2008;8(15):3105-17. doi: 10.1002/pmic.200700829
125. Cao S, Norris A, Miller JS, Ratovoson F, Razafitsalama J, Andriantsiferana R, et al. Cytotoxic triterpenoid saponins of *Albizia gummifera* from the Madagascar rain forest. *J Nat Prod* 2007;70(3):361-6. doi: 10.1021/np060506g
126. Mujoo K, Haridas V, Hoffmann JJ, Wächter GA, Hutter LK, Lu Y, et al. Triterpenoid saponins from *Acacia victoriae* (Bentham) decrease tumor cell proliferation and induce apoptosis. *Cancer Res* 2001;61(14):5486-90.
127. Liu LP, Cao XC, Liu F, Quan CF, Sheng XF, Ren KQ. Casticin induces breast cancer cell apoptosis by inhibiting the expression of forkhead box protein M1. *Oncol Lett* 2014;7(5):1711-7. doi: 10.3892/ol.2014.1911
128. Wang LM, Xie KP, Huo HN, Shang F, Zou W, Xie MJ. Luteolin inhibits proliferation induced by IGF-1 pathway dependent ER α in human breast cancer MCF-7 cells. *Asian Pac J Cancer Prev* 2012;13(4):1431-7. doi: 10.7314/apjcp.2012.13.4.1431
129. Vashisth P, Kumar N, Sharma M, Pruthi V. Biomedical applications of ferulic acid encapsulated electrospun nanofibers. *Biotechnol Rep (Amst)* 2015;8:36-44. doi: 10.1016/j.btre.2015.08.008
130. Watabe M, Hishikawa K, Takayanagi A, Shimizu N, Nakaki T. Caffeic acid phenethyl ester induces apoptosis by inhibition of NF κ B and activation of Fas in human breast cancer MCF-7 cells. *J Biol Chem* 2004;279(7):6017-26. doi: 10.1074/jbc.M306040200
131. Xie J, Xu Y, Huang X, Chen Y, Fu J, Xi M, et al. Berberine-induced apoptosis in human breast cancer cells is mediated by reactive oxygen species generation and mitochondrial-related apoptotic pathway. *Tumour Biol* 2015;36(2):1279-88. doi: 10.1007/s13277-014-2754-7
132. Zupkó I, Réthy B, Hohmann J, Molnár J, Ocsosvzki I, Falkay G. Antitumor activity of alkaloids derived from Amaryllidaceae species. *In Vivo* 2009;23(1):41-8.
133. Bharadwaj U, Eckols TK, Kolosov M, Kasembeli MM, Adam A, Torres D, et al. Drug-repositioning screening identified piperlongumine as a direct STAT3 inhibitor with potent activity against breast cancer. *Oncogene* 2015;34(11):1341-53. doi: 10.1038/onc.2014.72
134. Lou C, Yokoyama S, Saiki I, Hayakawa Y. Selective anticancer activity of hirsutine against HER2-positive breast cancer cells by inducing DNA damage. *Oncol Rep* 2015;33(4):2072-6. doi: 10.3892/or.2015.3796
135. Xiao D, Bommareddy A, Kim SH, Sehwat A, Hahm ER, Singh SV. Benzyl isothiocyanate causes FoxO1-mediated autophagic death in human breast cancer cells. *PLoS One* 2012;7(3):e32597. doi: 10.1371/journal.pone.0032597
136. Kessler JH, Mullauer FB, de Roo GM, Medema JP. Broad in vitro efficacy of plant-derived betulinic acid against cell lines derived from the most prevalent human cancer types. *Cancer Lett* 2007;251(1):132-45. doi: 10.1016/j.canlet.2006.11.003
137. Hordyjewska A, Ostapiuk A, Horecka A, Kurzepa J. Betulin and betulinic acid: triterpenoids derivatives with a powerful

- biological potential. *Phytochem Rev* 2019;18(3):929-51. doi: 10.1007/s11101-019-09623-1
138. Lambertini E, Lampronti I, Penolazzi L, Khan MT, Ather A, Giorgi G, et al. Expression of estrogen receptor alpha gene in breast cancer cells treated with transcription factor decoy is modulated by Bangladeshi natural plant extracts. *Oncol Res* 2005;15(2):69-79.
139. Li Z, Wang Y, Mo B. [The effects of carotenoids on the proliferation of human breast cancer cell and gene expression of bcl-2]. *Zhonghua Yu Fang Yi Xue Za Zhi* 2002;36(4):254-7.
140. Gopal YN, Arora TS, Van Dyke MW. Parthenolide specifically depletes histone deacetylase 1 protein and induces cell death through ataxia telangiectasia mutated. *Chem Biol* 2007;14(7):813-23. doi: 10.1016/j.chembiol.2007.06.007
141. Salisbury CM, Cravatt BF. Optimization of activity-based probes for proteomic profiling of histone deacetylase complexes. *J Am Chem Soc* 2008;130(7):2184-94. doi: 10.1021/ja074138u
142. Qiu YK, Dou DQ, Cai LP, Jiang HP, Kang TG, Yang BY, et al. Dammarane-type saponins from *Panax quinquefolium* and their inhibition activity on human breast cancer MCF-7 cells. *Fitoterapia* 2009;80(4):219-22. doi: 10.1016/j.fitote.2009.01.011
143. Chaudhry GE, Kassim MNI, Zafar MN, Mohammad H, Andriani Y, Ismail N, et al. Induction of apoptosis by *Stichopus chloronotus* and *Holothuria nobilis* fractions in human cervical cancer cell line, HeLa. *Int J Res Pharm Sci* 2020;11(1):1238-47. doi: 10.26452/ijrps.v11i1.1964