

- Urtica dioica L on lipid peroxidation, antioxidant enzyme systems and liver enzymes in carbon tetrachloride-treated rats. *World J Gastroenterol* 2005;11(42):6684-8. doi: 10.3748/wjg.v11.i42.6684
9. Schinella GR, Giner RM, Recio MC, Mordujovich de Buschiazzo P, Rios JL, Manez S. Anti-inflammatory Effects of South American *Tanacetum vulgare*. *J Pharm Pharmacol* 1998;50(9):1069-74. doi: 10.1111/j.2042-7158.1998.tb06924.x
 10. Baghaei A, Esmaily H, Abdolghaffari AH, Baeri M, Gharibdoost F, Abdollahi M. Efficacy of Setarud (IMOD®), a novel drug with potent anti-toxic stress potential in rat inflammatory bowel disease and comparison with dexamethasone and infliximab. *Indian J Biochem Biophys* 2010;47(4):219-26.
 11. Khairandish P, Mohraz M, Farzamfar B, Abdollahi M, Shahhosseiny M, Madani H, et al. Preclinical and phase I clinical safety of Setarud (IMOD™), a novel immunomodulator. *DARU* 2009;17(3):148-56.
 12. Khorram Khorshid HR, Novitsky YA, Abdollahi M, Shahhosseiny MH, Sadeghi B, Madani H, et al. Studies on potential mutagenic and genotoxic activity of Setarud. *DARU* 2008;16(4):223-8.
 13. Look M, Rockstroh JK, Rao GS, Barton S, Lemoch H, Kaiser R, et al. Sodium selenite and N-acetylcysteine in antiretroviral-naive HIV-1-infected patients: a randomized, controlled pilot study. *Eur J Clin Invest* 1998;28(5):389-97. doi: 10.1046/j.1365-2362.1998.00301.x
 14. Mahmoodpoor A, Eslami K, Mojtahedzadeh M, Najafi A, Ahmadi A, Dehnadi-Moghadam A, et al. Examination of Setarud (IMOD™) in the management of patients with severe sepsis. *DARU* 2010;18(1):23-8.
 15. Mohseni-Salehi-Monfared SS, Habibollahzadeh E, Sadeghi H, Baeri M, Abdollahi M. Efficacy of Setarud (IMOD™), a novel electro-magnetically-treated multi-herbal compound, in mouse immunogenic type-1 diabetes. *Arch Med Sci* 2010;6(5):663-9. doi: 10.1114/achm.2010.17078
 16. Ogunro PS, Ogungbemisigbe TO, Elemie PO, Egbewale BE, Arewale SA. Plasma selenium concentration and glutathione peroxidase activity in HIV-1/AIDS infected patients: a correlation with the disease progression. *Niger Postgrad Med J* 2006;13(1):1-5.
 17. Paydary K, Lalamzadeh-Fard S, Khorram Khorshid HR, Kamali K, SeyedAlinaghi S, Mohraz M. Safety and efficacy of Setarud (IMOD™) among people living with HIV/AIDS: a review. *Recent Pat Antiinfect Drug Discov* 2012;7(1):66-72. doi: 10.2174/157489112799829756
 18. Daneshmand P, Saliminejad K, Dehghan Shasaltaneh M, Kamali K, Riazi GH, Nazari R, et al. Neuroprotective Effects of Herbal Extract (*Rosa canina*, *Tanacetum vulgare* and *Urtica dioica*) on Rat Model of Sporadic Alzheimer's Disease. *Avicenna J Med Biotechnol* 2016;8(3):120-5.
 19. Ghanbari S, Yonessi M, Mohammadirad A, Gholami M, Baeri M, Khorram-Khorshid HR, et al. Effects of IMOD™ and Angipars™ on mouse D-galactose-induced model of aging. *DARU* 2012;20(1):68. doi: 10.1186/2008-2231-20-68
 20. Morris RG. Morris water maze. *Scholarpedia* 2008;3(8):6315. doi: 10.4249/scholarpedia.6315
 21. Paxinos G, Watson C. *The Rat Nervous System: The New Coronal Set*. New York: Elsevier; 2004.
 22. National Research Council (US) Committee. *Guide for the Care and Use of Laboratory Animals*. 8th ed. Washington (DC): National Academies Press (US); 2011.
 23. Moura AC, Pazzari VM, Agnes G, Almeida S, Giovannardi M, Veiga AB. Transcriptional expression study in the central nervous system of rats: what gene should be used as internal control? *Einstein (Sao Paulo)* 2014;12(3):336-41. doi: 10.1590/s1679-45022014ao3042
 24. Silver N, Cotroneo E, Proctor G, Osailan S, Paterson KL, Carpenter GH. Selection of housekeeping genes for gene expression studies in the adult rat submandibular gland under normal, inflamed, atrophic and regenerative states. *BMC Mol Biol* 2008;9:64. doi: 10.1186/1471-2199-9-64
 25. Mateo I, Llorca J, Infante J, Rodríguez-Rodríguez E, Fernández-Viadero C, Pena N, et al. Low serum VEGF levels are associated with Alzheimer's disease. *Acta Neurol Scand* 2007;116(1):56-8. doi: 10.1111/j.1600-0404.2006.00775.x
 26. Tang H, Mao X, Xie L, Greenberg DA, Jin K. Expression level of vascular endothelial growth factor in hippocampus is associated with cognitive impairment in patients with Alzheimer's disease. *Neurobiol Aging* 2013;34(5):1412-5. doi: 10.1016/j.neurobiolaging.2012.10.029