

How to cite this article: Ebadi sharafabad B, Abdoli A, Panahi M, Abdolmohammadi Khiav L, Jamur P, Abedi Jafari F, Dilmaghani A. Anti-tumor effects of cisplatin synergist in combined treatment with *Clostridium novyi*-NT spores against hypoxic microenvironments in a mouse model of cervical cancer caused by TC-1 cell line. *Advanced Pharmaceutical Bulletin*, doi: 10.34172/apb.2023.084

doi: [10.34172/apb.2023.084](https://doi.org/10.34172/apb.2023.084)

Anti-tumor effects of cisplatin synergist in combined treatment with *Clostridium novyi*-NT spores against hypoxic microenvironments in a mouse model of cervical cancer caused by TC-1 cell line

Behrouz Ebadi sharafabad¹, Asghar Abdoli², Mohammad Panahi², Lida Abdolmohammadi Khiav³, Parisa Jamur², Fatemeh Abedi Jafari⁴, Azita Dilmaghani^{1,5*}

1. Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.
2. Department of Hepatitis and HIV, Pasteur Institute of Iran (IPI), Tehran, Iran
3. Department of Anaerobic Vaccine Research and Production, Specialized Clostridia Research Laboratory, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization, Karaj, Iran
4. Department of Microbiology and Microbial Biotechnology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, Tehran, Iran
5. Infectious and Tropical Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

* **Correspond:** Azita Dilmaghani

Email: dilmaghani.a@gmail.com

Fax: 04133344798

Tel: +989141035048

ORCID number: 0000-0001-6556-8332

Abstract

Purpose: Despite the development of anti-HPV vaccines, cervical Cancer is still a common disease in women, especially in developing countries. The presence of a hypoxic microenvironment causes traditional treatments to fail. In this study, we presented a combined treatment method based on the chemotherapeutic agent cisplatin and *Clostridium novyi*- NT spores to treat Normoxic and Hypoxic areas of the tumor.

Methods: TC -1 Cell line capable of expressing HPV-16 E6/7 oncoproteins was subcutaneously transplanted into female 6-8 week old C57/BL6 mice. The tumor-bearing mice were randomly divided into four groups and treated with different methods after selecting a control group. Group 1: Control without treatment (0.1 ml sterile PBS intratumorally), Group: *C. novyi*- NT (10^7 *C. novyi*- NT). Group 3: Receives cisplatin intraperitoneally (10 mg/kg). Fourth group: Intratumoral administration of *C. novyi*- NT spores + intraperitoneal cisplatin. Western blot analysis was used to examine the effects of anti-hypoxia treatment and expression of HIF-1 and VEGF proteins.

Results: The results clearly showed that combined treatment based on *C. novyi*-NT and cisplatin significantly reduced the expression of HIF-1 alpha and VEGF proteins compared to cisplatin alone. At the same time, the amount of necrosis of tumor cells in the combined treatment increased significantly compared to the single treatment and the control. At the same time, the mitotic count decreased significantly.

Conclusion: Our research showed that developing a combined treatment method based on *C. novyi*-NT and cisplatin against HPV-positive cervical Cancer could overcome the treatment limitations caused by the existence of hypoxic areas of the tumor.

Keywords

Clostridium novyi-NT, Cisplatin, TC-1 Cell line, cervical cancer, Tumor Hypoxia

1- Introduction

It is well known that the human papillomavirus (HPV) is the leading cause of more than 90% of cervical cancers¹. The E6 and E7 oncoproteins of high-risk HPV types, particularly HPV16, alter routine host anti-tumor pathways, such as intrinsic apoptosis, and then induce cancer in host cells by generating genomic instability by targeting two host tumor suppressors, p53 and the retinoblastoma protein (pRb)^{2,3}. Despite developing prevention methods such as vaccination of adolescent girls against high-risk strains of HPV and modern diagnostic procedures in developed countries, in poor and less developed countries, 500,000 new cases of cervical Cancer and more than 300,000 deaths are reported due to this disease⁴⁻⁶. Today, various treatment regimens based on surgery, chemotherapy, and Radiotherapy are combined to treat cervical Cancer⁷. However, there are still challenges to treatment success⁸. An in-depth understanding of HPV-related malignancies and treatment failure factors is essential to provide successful treatment regimens.

The fact that solid cervical tumors have hypoxic areas with oxygen concentrations below 1.5% has been proven⁹. Hypoxia is an adverse prognostic factor that suppresses the host's immune system and severely reduces the effectiveness of various treatments, including Radiotherapy and chemotherapy^{10,11}. It has recently been shown that under hypoxic conditions, HPV-infected tumor cells, by being in a reversible proliferative stasis state, drastically reduce E6 / E7 expression and cause the cell to escape immunity¹². If tumor cells are in the proper oxygen delivery state, the expression of E6 / E7 oncoproteins resumes rapidly and, after treatment, acts as an abundant source of oncoprotein expression^{12,13}. This complex interaction, which can eventually cause the tumor cell to escape from the senescent, is mediated through the Phosphoinositide 3-kinase (PI3K) / AKT / mammalian target of rapamycin (mTOR) signaling pathway in hypoxic conditions, despite decreased E6 / E7 oncoprotein expression¹³⁻¹⁶. HIF-1 α (Hypoxia-inducible factor 1-alpha) acts as a significant protein in the adaptation of cancer cells to hypoxic conditions, and its role in the angiogenesis and development of various cancers has been discussed^{17,18}. By stabilizing HIF-1 α in hypoxic conditions in cervical cancer, regulated in development and DNA damage responses 1 (REDD1) is activated and inhibits mTORC1^{15,19,20}. As a result, despite AKT-mediated inhibition of E6 / E7, hypoxic HPV-positive cancer cells survive the aging process¹³. Thus, targeting tumor cells in hypoxic areas can be vital to a successful treatment regimen.

Cisplatin is one of the most common chemotherapy drugs prescribed as a first-line treatment for many cancers²¹. This substance induces intrinsic cell death by creating irreversible bindings to the DNA of tumor cells²². However, research has shown that the presence of a hypoxic

microenvironment in the tumor, through decreased expression of anti-tumor protein p53²³, Increased cancer stemness²⁴, Regulation of non-coding RNAs²⁵, Generation of reactive oxygen species (ROS)²⁶, Increased exosome secretion²⁷, Loss of mismatch Repair²⁸, Cyclophilin overexpression²⁹, Surviving overexpression³⁰, and increased Glucose transporter 1 (GLUT-1) expression³¹ may increase the resistance of hypoxic tumor cells to this treatment. Therefore, offering complementary therapies based on anti-hypoxia agents, which target the expression and stability of the HIF-1 α axial protein that plays a crucial role in inducing this resistance³², is considered a promising factor in increasing therapeutic efficacy.

Today, various strategies have been developed to treat all types of solid tumors, including Surgery, Chemotherapy, Radiotherapy, Targeted therapy, Immunotherapy, Combination therapy, and Extracellular Vesicle (EV) therapy³³⁻³⁵. More than half of cancer patients consider Radiotherapy as part of their treatment plan. Despite the benefits of this treatment modality, the development of various physical complications and, most importantly, the susceptibility of the patient's normal cells to carcinogenesis from radiation exposure appears to be a significant challenge³⁶. In the context of immunotherapy, immune checkpoint inhibition is considered standard treatment, but the most significant concern is for cold cancers, which are unlikely to respond to such therapy³⁷. Some medicines are inefficient because not every tumor can be removed by surgery or because it is resistant to chemotherapy and radiotherapy. Chemotherapy decreased drug penetration into neighboring tumor cells in the weakly vascularized area, environmental toxicity and reduced immunity, which faces some clinical problems after treatment³⁸. The existence of these challenges in treatment, driven mainly by the disturbances in tumor physiology and the presence of hypoxic areas in all types of solid tumors, led researchers to revive the method of using anaerobic bacteria at the beginning of the 20th century, because after injecting these bacteria into the tumor, they grow selectively in the hypoxic areas of the tumor and can cause its destruction³⁹. The existence of these challenges in treatment, caused mainly by the disturbances in tumor physiology and the presence of hypoxic areas in all types of solid tumors, led researchers to revive the method of using anaerobic bacteria at the beginning of the 20th century, because after injecting these bacteria into the tumor, they grow selectively in the hypoxic areas of the tumor and can cause its destruction³⁹.

One of the most attractive ways to overcome the resistance of cancer cells in hypoxic areas of solid tumors to standard therapies, which have been vigorously pursued over the last two decades, is to use the spores of a non-lethal type of *Clostridium novyi*⁴⁰. After injection into the tumor, spores of *Clostridium novyi*-NT migrate to hypoxic areas of the cancer, and due to the hypoxic nature of these areas, begin to germinate and cause cells to lysis in this area⁴¹. The nature of the therapeutic function of this agent is not fully understood⁴², but it appears that this bacterium induces a robust immune response from the host⁴³. On the other hand, by secreting several extracellular proteins lysing the lipid structure of the cell membrane, including phospholipase C (PLC) (NT01CX0979), it causes the destruction of tumor cells in the hypoxic areas⁴⁴.

In this study, for the first time, a combined treatment strategy based on cisplatin and *Clostridium neuvi*-NT spores is presented in a mouse model for HPV-related cervical cancer and the effects of this type of treatment on resistance factors in hypoxic tumor areas are shown. Understanding these effects could pave the way for combination therapies to make cisplatin more effective against HPV-related cancers.

2- Materials and Methods

Preparation of Cell culture and *Clostridium novyi*-NT spores

To cause HPV-Associated cervical Cancer in a mouse model, the TC-1 cell line with the ability to express HPV-16 E6 / 7 oncoproteins⁴⁵ was prepared by the National Cell bank of Iran (NCBI) affiliated with Pasteur Institute (Tehran, Iran). TC-1 cell line was suspended in RPMI-1640 (Sigma, USA) medium with 10% fetal bovine serum (FBS) (Gibco, USA), 1% penicillin-streptomycin (Sigma, USA), 25 mM HEPES (Sigma, USA), 1% glutamine (Sigma, USA) and incubated at 37 ° C and 5% CO₂. We have previously been able to generate spores without lethal genes from the wild *C. novyi* type B strain⁴⁶. After processing, *Clostridium novyi*-NT spores were packaged in lyophilized tablets containing 10⁷ and stored at -20 ° C for future use.

Animal study design and creation of a mouse model of HPV-Associated Cervical Cancer

This study was reviewed and approved by the Research and Technology Vice-Chancellor of Tabriz University of Medical Sciences (IR.TBZMED.VCR.REC.1398.434). Also, all animal work was under the standards of the Iran National Committee for Ethics in Biomedical Research and was performed under the supervision of the Ethics Committee in Animal Research of the Pasteur Institute of Iran. Thirty female 6-8 weeks old C57 / BL6 mice weighing 18 to 22 grams were obtained from Pasteur Institute (Iran, Karaj) and fed for seven days according to laboratory standards. The animals were then randomly divided into five groups and six mice in each group (group= 5 and n=6). One group was randomly chosen as a non-tumor and healthy group, and the other four groups were selected to challenge the tumor. To cause HPV-Associated cervical cancer, one million TC-1 cells were suspended in 0.2 ml of PBS (Sigma, USA) and injected subcutaneously into the right flank of each mouse. During this period, the mice were monitored daily, and with the appearance of palpable tumors, the size of the tumors was measured using a caliper. The volume of the tumors was determined using the standard formula (longest diameter of the tumor) × (shortest diameter)² × 0.5.

Treatment with *Clostridium novyi*-NT spores and cisplatin

Cisplatin (CAS 15663-27-1) was purchased from sigma Aldrich (Sigma, USA). To evaluate the therapeutic effects of cisplatin chemotherapy alone or in combination with the anti-tumor effects of *Clostridium novyi*-NT spores, after reaching the volume of mouse tumors in the range of 300 to 500 mm³, the following was performed; Group 1: They did not receive any treatment and were injected with 0.1 ml of sterile PBS intratumoral. Group 2: A lyophilized tablet containing 10⁷ *Clostridium novyi*-NT spores was suspended in 0.1 ml of sterile PBS and injected intra-tumor into different tumor parts. Group 3: 10 mg/kg cisplatin⁴⁷ was dissolved in 0.1 ml sterile PBS and injected intraperitoneally into each mouse. Group 4 - This group is a combination therapy group and received 10⁷ *Clostridium novyi*-NT spores suspended in 0.1 ml PBS on day 0. Subsequently, 8 hours later, they received 10 mg/kg cisplatin dissolved in 0.1 ml PBS intraperitoneally. According to our previous study⁴⁶, the 15th day after the start of treatment was chosen as the day of the end of treatment; on this day, all animals received 30 µl of anesthetic solution with the following characteristics: Ketamine 10% (100 mg/ml; Medistar, Ascheberg, Germany) and xylazine 2% (20 mg/ml; Riemsers, Greifswald, Germany) were combined in a single insulin syringe (2 parts ketamine and 1 part xylazine)⁴⁸ and After ensuring deep anesthesia of the animals, a blood sample was taken through the heart muscle and maintained at 10% using EDTA (ethylene diamine tetraacetic acid) as an anticoagulant. The animals were then sacrificed through the displacement of the neck vertebra. Tumor tissue was then carefully isolated and kept at -70 ° C for gene analysis and protein expression and in 10% formalin solution for histopathological examination.

RNA Extraction and cDNA synthesis

Total RNA was extracted from 100 mg of cervical cancer tissue isolated from each mouse and used RNA Extraction kit (Thermo Fisher Scientific, USA) according to the manufacturer's instructions. The purification of total RNA was evaluated by NanoDrop ND-1000 (NanoDrop, USA) spectrophotometer. Extracted product was tested on a 2% agarose gel to check RNA integrity. 500 µg of extracted RNA was used for cDNA synthesis by random hexamer primer using a reverse transcription kit (Biotech Rabbit, Germany) according to the manufacturer's instructions.

Real-time polymerase chain reaction (RT-PCR) analysis

The expression level of the GLUT -1 and PLC genes was determined by RT-PCR. 2 µl of syntonies cDNA was subjected to PCR cycle with SYBR-Green 2x Master Mix (Amplicon, Denmark). PCR conditions included pre-denaturation at 95°C for 15 minutes, denaturation at 95°C for 30 seconds, and denaturation at 60°C for 45 seconds for 40 cycles. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as a reaction internal reference. All the samples were determined 3 times. The special primers were as follows: GLUT-1, forward: 5'-GAGAACCGGGCCAAGAGTG-3' and reverse 5'- TTCTTCTCCCGCATCATCTG-3'; PLC, forward 5'- GGAGCATCAAGTAAAGCGTA-3' and reverse 5'- CATTTCGGATCATAATCAGGA-3'; GAPDH, forward 5'-GCCAAAGGGTCATCATCTC-3' and reverse 5'-GTAGAGGCAGGGATGATGTT-3'. All primers were designed by Oligo 7 software and synthesized by Metabion Company (Germany). The target mRNA value was measured by comparison with the control sample, then the comparison period threshold ($\Delta\Delta Ct$) method was used for calculation.

Western blot analysis

Western blot analyses were performed as previously described with some modifications⁴⁹. The lysates were removed by centrifugation at 14,000 rpm for 20 min at 4°C. According to the manufacturer's instructions, the BCA Protein Quantification kit determined the protein concentration of exosome lysates. The exosome lysates were mixed with a 2X Laemmli sample buffer equal to volume. Lysates (15 µg) were then subjected to SDS-PAGE after a 5 min boiling and subsequently transferred to a 0.2 µm immune-Blot™ polyvinylidene difluoride (PVDF) membrane (Bio-Rad Laboratories, CA, USA). The membranes were then blocked with 5% BSA (Cat No: A-7888; Sigma Aldrich, MO, USA) in 0.1% Tween 20 for 1 h. Then, the membranes were incubated with Anti-HIF-1 alpha (Abcam), Anti-VEGF (Abcam), and anti-beta actin-loading control antibodies (Abcam) for one h at room temperature. Subsequently, membranes were washed thrice with TBST and incubated with goat anti-rabbit IgG H&L (HRP) (Abcam) secondary antibodies. The membranes were then incubated with enhanced chemiluminescence (ECL) for 1–2 min. Protein expression was normalized to β-actin. Densitometry of protein bands was performed using the gel analyzer Version 2010a software (NIH, USA), such that the percentage area under the curve of each band was divided by the percentage area under the curve of its corresponding actin band, and then calculated values were compared between groups as described previously.⁴⁹

Reactive oxygen species (ROS) analysis

ROS1 / ROS ELISA Kit (LifeSpan BioSciences, USA) was used to measure the number of reactive oxygen species based on the Sandwich ELISA method. Tumor tissues were washed in PBS to remove excess blood. Next, the tissues were weighed before homogenization. The tissues were minced and homogenized in 10 ml PBS with a glass homogenizer on ice. All reagents, working standards, and samples were prepared according to the kit instructions and placed at room temperature for 20 minutes. 100 µl of sample and standard were added to each

well and incubated for 2 hours at 37°C. The plate was emptied, and 100 µl of Detection Reagent A was added and incubated for 1 hour at 37°C. The plate was opened and washed thrice with 400 µl of the wash solution in the plate kit. 100 µl of Detection Reagent B was added and incubated for 30 minutes at 37°C. The plate was emptied and washed thrice with 400 microliters of the wash solution in the plate kit. 90 ml of substrate solution was added to each well and incubated for 20 minutes in the dark at 37°C. Finally, 50 µl of Stop solution was added to each well. Using an ELISA device, the amount of light absorption was read at 450 nm, and based on the amount of light absorption, the standard curve was drawn, and based on the slope of the line and the width of the beginning, and the amount of ROS concentration in each well was measured.

Histopathological studies

Tumor tissue was fixed in 10% formalin and dehydrated through graded ethanol (70%, 90%, 96%, and 100%). Then, the tissues were paraffin-embedded, and 5µm sections were taken from the tissue by microtome. So, the slides were stained with hematoxylin and eosin dyes and Giemsa staining according to routine laboratory protocols. And then, for the histology of tumors, conventional hematoxylin and eosin (H&E) staining and light microscopy were used. Formula calculated the relative necrotic area (%) of tumor tissues: Relative necrotic area (%) = Necrotic area in tumor section/Total area of tumor section × 100⁵⁰.

Statistical analysis

All data for this experiment have been presented as mean ± SD. We used GraphPad Prism 9 software for statistical analysis, including a one-way analysis of variance (ANOVA) and a t-test, and selected 0.05 as statistically significant.

3- Results and Discussion

***Clostridium novyi*-NT spores combined with cisplatin showed a synergistic anti-tumor effect in TC-1 mice models**

The tumor tissues stained by the H&E technique were carefully examined to understand the double anti-tumor function of the combined treatment based on *Clostridium novyi*-NT spores and the cisplatin (Figure1-a). The detailed examination showed that the percentage of necrosis in the treatment group that received cisplatin and bacteria compared to the group that received only cisplatin increased statistically significantly (P< 0.0001) (Figure1-c). Also, detailed microscopic examinations showed that the mitotic count, as an indicator of tumor malignancy, in the group receiving combined treatment was statistically significantly lower compared to the control group and the group treated only with cisplatin (P< 0.0001) (Figure1-d).

Effects of *Clostridium novyi*-NT spores and cisplatin on GLUT 1 mRNA expression level in mice with cervical Cancer and qPCR for detecting *Clostridium novyi*-NT

Glucose transporter-1 (GLUT1) has been suggested as a prognostic factor in various cancers associated with treatment resistance and immune evasion⁵¹. In a study by Byoung Hyuck Kim et al., with the aim of intending to investigate the predictive effect of GLUT1 in cervical Cancer, they analyzed the data of 298 patients. They showed that high expression of GLUT1 with old age, squamous cell carcinoma, high tumor stage, metastasis to pelvic lymph nodes, and low hysterectomy rates are associated. Finally, they concluded that GLUT1 expression and

HPV16 subtype might have independent prognostic value in cervical Cancer. GLUT1-mediated immune modulation may be one of the crucial reasons for treatment failure, especially in HPV16 positive group⁵². As a result, it can be concluded that treatments that modulate the expression of glucose transporter-1 (GLUT1) will have promising efficacy. The qPCR technique was employed to study the effect of *Clostridium novyi*-NT spores and cisplatin treating cervical cancer in mice. The GAPDH gene was applied as a housekeeping gene (Figure 2), showing the expression level of GLUT 1 and *Clostridium novyi*-NT Phospholipase C (PLC, Gene: NT01CX_0979) in different groups of mice. Results show that the GLUT 1 expression was significantly reduced in *Clostridium novyi*-NT spores compared to control ($P = 0.0001$), the combination of *C. novyi*-NT spores and cisplatin compared to *Clostridium novyi*-NT spores ($P < 0.05$), cisplatin compared to control and variety of *Clostridium novyi*-NT spores and cisplatin compared to control ($P < 0.0001$). *Clostridium novyi*-NT Phospholipase C (PLC, Gene: NT01CX_0979) was considered a factor confirming the presence and germination of *Clostridium novyi*-NT spores. As expected, the expression level of this gene was zero in the control group and the group that received only cisplatin, as the PLC expression levels were significantly increased in cisplatin compared to control and cisplatin compared to *Clostridium novyi*-NT spores ($P < 0.005$) and was seen the reduced level of expression in the combination of *Clostridium novyi*-NT spores and cisplatin compared to cisplatin ($P < 0.05$) (Figure 2).

Effects of *Clostridium novyi*-NT spores and cisplatin on HIF-1 α and VEGF proteins expression level in mice with cervical Cancer

Hypoxia-inducible factors (HIFs) have been identified in the hypoxic tumor microenvironment as essential transcription factors that regulate the expression of many of genes related to angiogenesis, metastasis, cell proliferation, and resistance to chemotherapy and Radiotherapy. HIF-1 α promotes cancer cell proliferation, and VEGF induces vascular endothelial cell division to promote tumor growth. High expression of HIF-1 α and VEGF in cervical cancer tissues is associated with clinical stage, pathological grade and lymph node metastasis⁵³. The effect of *Clostridium novyi*-NT spores and cisplatin on protein expression level was examined by western blot and used anti-beta actin-loading by control antibodies. There were statistically significant differences in the protein expression levels of HIF-1 α and VEGF in different groups of mice. Figure 2 shows protein expression levels in all groups of mice. Results show that the HIF-1 α protein expression significantly reduces in the combination of *Clostridium novyi*-NT spores and cisplatin group compared to control ($P < 0.05$), the combination of *Clostridium novyi*-NT spores and cisplatin group compared to cisplatin ($P < 0.05$). The VEGF protein expression significantly reduces in the combination of *Clostridium novyi*-NT spores and cisplatin group compared to control ($P < 0.05$), the combination of *Clostridium novyi*-NT spores and cisplatin group compared to *Clostridium novyi*-NT spores ($P < 0.05$).

Effects of *Clostridium novyi*-NT spores and cisplatin on Reactive oxygen species (ROS) in mice with cervical cancer

In several types of cancer, reactive oxygen species (ROS) are widely associated with carcinogenesis and cancer progression⁵⁴. To investigate the effect of *Clostridium novyi*-NT spores and cisplatin on the changes related to Reactive oxygen species (ROS), it was performed according to the protocol of ROS1 / ROS ELISA Kit (LifeSpan BioSciences, USA). The results showed that the number of reactive oxygen species (ROS) was significantly reduced in the group receiving cisplatin along with *Clostridium novyi*-NT spores compared to the group

receiving cisplatin and *Clostridium novyi*-NT spores alone and compared to the control group ($P = 0.0001$). (Figure 4)

It has been well established that the most important cause of cervical cancer is a chronic infection caused by high-risk types of human papillomavirus⁵⁵. These viruses cause genomic instability and cancer by deactivating suppressor proteins p53 and pRb through oncoproteins E6 and E7. Despite the development of prevention methods for this malignancy based on vaccination against high-risk types of HPV, the infection rate is still high in poor and developing countries, and as a result, there is a need to develop new treatment approaches⁵⁶. One of the essential factors that strongly affect the success of conventional treatments of this disease is the presence of hypoxic micro-regions in this type of tumor. Therefore, it can be said that hypoxia is the most critical factor in the failure of conventional treatments based on chemotherapy and radiotherapy⁵⁷.

The most important factor that causes cancer cells to adapt to hypoxic conditions is HIF-1 α . This transcription factor, as the central axis of cell adaptation to hypoxic conditions, by regulating more than 100 genes, mediates the requirements for the continued abnormal proliferation of tumor cells, abnormal metabolisms, metastasis, and resistance to therapeutic agents^{53,58}. As a result of the stabilization of this protein, Vascular endothelial growth factor (VEGF) expression is induced, which is a potential factor in the induction of tumor angiogenesis and also causes the expression of the glucose transporter gene and thus increases glycolysis⁵⁹.

One of the compounds widely used in treating various malignancies, including cervical Cancer, is cisplatin⁶⁰. This combination has significant effects on normoxic tumors, but the hypoxic condition of the tumor is the main reason for the failure of this treatment²¹. Research has shown that cisplatin causes the death of tumor cells through upregulating of the p53 suppressor protein²². However, in the hypoxic conditions, HIF-1 α strongly suppresses the level of p53⁶¹. As a result, the resistance to this treatment increases. At the same time, as oxygen decreases in the hypoxic conditions of the tumor, the leakage of electrons from the electron transport chain occurs, which is the reason for the reduction of the electron flow through the mitochondrial complex of the electron transport chain. This process produces reactive oxygen species that can enhance mitochondrial fragmentation and further reduce the expression of p-Drp1 and Mfn1, resulting in increased resistance to cisplatin. Therefore, providing combined treatment regimens based on cisplatin and tumor anti-hypoxic agents can promise to overcome the therapeutic limitations of cisplatin and significantly strengthen its effectiveness in hypoxic conditions^{21,62}.

The hypoxic condition of the tumor creates a suitable environment for the germination and growth of *Clostridium novyi*-NT spores^{63,64}. We and others have shown that this species, simultaneously germination and growing in hypoxic regions of tumor types, has caused extensive regression in various tumors in rodent models^{46,65,66}. Due to the unclear nature of this function, we tried to present a combined treatment method based on cisplatin and *Clostridium novyi*-NT spores in the HPV-related cervical cancer model. Our goal was to increase the therapeutic efficiency and overcome the resistance of cisplatin treatment in hypoxic conditions.

This study showed that the expression level of HIF-1 α and VEGF proteins decreased in the group of tumor mice that received the combined treatment, in a statistically significant way, compared to the control group. By examining the aberrant expression patterns of 3352 differentially expressed genes in 306 cervical cancer samples, Xiaofeng Xu et al. concluded that the HIF-1 signaling pathway related to TFRC might play an important role in cervical

cancer⁶⁷. Conversely, LIU Kai et al. showed that HIF-1 α and VEGF could be considered a parameter in evaluating the progress, metastasis, and prognosis of HPV-related cancers⁶⁸.

Various studies have shown that VEGF plays a significant role in angiogenesis and cancer development. On the other hand, abnormal expression of GLUT 1, as a downstream gene of HIF-1 α , in HPV-positive head and neck cancer tumors increases the probability of invasion and metastasis. Therefore, it can be said that the decrease in the expression of this gene is a positive sign of the effectiveness of various therapeutic strategies. Fortunately, our recent study also showed that the expression level of GLUT 1 was significantly reduced in the group that received the combined treatment of *Clostridium novyi* spores-NT and cisplatin.

Transmission electron microscopy and atomic force microscopy revealed that *Clostridium novyi*-NT spores are surrounded by an amorphous layer interwoven with parasporal honeycomb layers sequentially dissolved during germination. Vegetative cells Most of these spore-specific genes encode spore coat proteins or proteins with redox activity, which could aid germination by scavenging reactive oxygen species^{44,69}. As mentioned earlier, in hypoxic tumors, the reduction in oxygen consumption causes a decrease in electron flow through the mitochondrial complex of the electron transport chain²⁶. This leads to the leakage of electrons from the electron transport chain, resulting in excessive production of ROS, which in turn increases cisplatin resistance. In various tumors, the hypoxic tumor microenvironment induces ROS production, which increases mitochondrial fission and, thus, cisplatin resistance by downregulating the expression of p-Drp1 (Ser637) and Mfn1⁶². Our study showed that the number of reactive oxygen species (ROS) in the group receiving combined treatment was significantly reduced compared to the control group.

The molecular mechanisms involved in the anti-tumor function of *Clostridium novyi*-NT have not been fully established. However, it seems to be related to the destructive properties of the enzymes it secures the induction of host immunity by the secretion of large amounts of cytokines^{40,70}. In 2006, Bettogowda et al. determined the genomic sequence of *Clostridium novyi*-NT spores. They showed that *Clostridium novyi*-NT could affect the structure of the lipid layers of the host cell wall by secreting various proteins such as phospholipase C (PLC) (NT01CX0979), which, in addition to having a direct effect on tumor cell lysis, Activation of host antitumor immune responses is also stimulated⁴⁴.

In this study, we measured the germination of *Clostridium novyi*-NT spores by measuring the amplification of the phospholipase C (PLC) gene (NT01CX0979), along with the use of Giemsa stain. The results showed that, as expected, the expression level of this gene was zero in the control group and the group that received only cisplatin. As the PLC expression levels were significantly increased in cisplatin compared to control and cisplatin compared to *Clostridium novyi*-NT spores and seen the reduced level of expression in the combination of *Clostridium novyi*-NT spores and cisplatin compared to cisplatin.

The results clearly showed that using a combined treatment regimen based on *Clostridium novyi*-NT spores and cisplatin can overcome the therapeutic limitations of cisplatin chemotherapy in tumors with hypoxic areas in cervical Cancer. It also increases treatment efficiency and is considered a favorable option in developing combined treatment methods for solid hypoxic tumors.

4- Conclusion

Our research showed that providing a combined treatment method based on *Clostridium novyi*-NT spores and cisplatin can overcome the limitations and therapeutic resistance caused by hypoxic microenvironments in HPV-positive cervical cancers. However, the mechanism of this

phenomenon is not completely clear, but by overcoming the existing limitations, this method has a positive perspective in developing effective anti-tumor hypoxia treatment methods.

Abbreviation List

HPV: Human papillomavirus
pRb: retinoblastoma protein
PI3K: Phosphoinositide 3-kinase
mTOR: mammalian target of rapamycin
HIF-1 α : Hypoxia-inducible factor 1-alpha
REDD1: regulated in development and DNA damage responses 1
GLUT-1: Glucose transporter 1
ROS: reactive oxygen species
PLC: phospholipase C
NCBI: National Cell bank of Iran
FBS: fetal bovine serum
EDTA: ethylene diamine tetraacetic acid
GAPDH: Glyceraldehyde 3-phosphate dehydrogenase
PVDF: polyvinylidene difluoride
RT-PCR: Real-time polymerase chain reaction
H&E: hematoxylin and eosin
VEGF: Vascular endothelial growth factor

Competing interests

The authors declare that they have no direct or indirect conflict of interest.

Ethics approval and consent to participate

This study was reviewed and approved by the Research and Technology Vice-Chancellor of Tabriz University of Medical Sciences (IR.TBZMED.VCR.REC.1398.434). Also, all animal work was under the standards of the Iran National Committee for Ethics in Biomedical Research and was performed under the supervision of the Ethics Committee in Animal Research of the Pasteur Institute of Iran.

Funding

This project was financially supported by Iran National Science Foundation (INSF) (Grant Number: 99025014)

Acknowledgments

We sincerely thank Iran National Science Foundation (INSF) for its financial support. We thank Tabriz University of Medical Sciences and the Pasteur Institute of Iran for supporting this research.

Disclosure of potential conflicts of interest

The authors declare that they have no direct or indirect conflict of interest.

Informed consent

All information obtained during this study is published in this article.

References

1. Okunade KS. Human papillomavirus and cervical cancer. *Journal of Obstetrics and Gynaecology* 2020;40(5):602-8. doi: 10.1080/01443615.2019.1634030
2. Wardak S. Human Papillomavirus (HPV) and cervical cancer. *Medycyna doświadczalna i mikrobiologia* 2016;68(1):73-84.
3. Mittal S, Banks L. Molecular mechanisms underlying human papillomavirus E6 and E7 oncoprotein-induced cell transformation. *Mutation Research/Reviews in Mutation Research* 2017;772:23-35. doi: 10.1016/j.mrrev.2016.08.001
4. Spayne J, Hesketh T. Estimate of global human papillomavirus vaccination coverage: analysis of country-level indicators. *BMJ open* 2021;11(9):e052016. doi: 10.1136/bmjopen-2021-052016
5. Colzani E, Johansen K, Johnson H, Celentano LP. Human papillomavirus vaccination in the European Union/European Economic Area and globally: a moral dilemma. *Eurosurveillance* 2021;26(50):2001659. doi: 10.2807/1560-7917.ES.2021.26.50.2001659
6. Zhang X, Zeng Q, Cai W, Ruan W. Trends of cervical cancer at global, regional, and national level: Data from the Global Burden of Disease study 2019. *BMC public health* 2021;21(1):1-10. doi: 10.1186/s12889-021-10907-5
7. Cohen PA, Jhingran A, Oaknin A, Denny L. Cervical cancer. *The Lancet* 2019;393(10167):169-82. doi: 10.1016/S0140-6736(18)32470-X
8. Davies-Oliveira J, Smith M, Grover S, Canfell K, Crosbie E. Eliminating cervical cancer: progress and challenges for high-income countries. *Clinical Oncology* 2021;33(9):550-9. doi: 10.1016/j.clon.2021.06.013
9. Vaupel P. Pathophysiology of solid tumors. The impact of tumor biology on cancer treatment and multidisciplinary strategies: Springer; 2009. p. 51-92.
10. Chouaib S, Noman M, Kosmatopoulos K, Curran M. Hypoxic stress: obstacles and opportunities for innovative immunotherapy of cancer. *Oncogene* 2017;36(4):439-45. doi: 10.1038/onc.2016.225
11. Noman MZ, Hasmmim M, Messai Y, Terry S, Kieda C, Janji B, et al. Hypoxia: a key player in antitumor immune response. A review in the theme: cellular responses to hypoxia. *American Journal of Physiology-Cell Physiology* 2015;309(9):C569-C79. doi: 10.1152/ajpcell.00207.2015
12. Hoppe-Seyler K, Mändl J, Adrian S, Kuhn BJ, Hoppe-Seyler F. Virus/host cell crosstalk in hypoxic HPV-positive cancer cells. *Viruses* 2017;9(7):174. doi: 10.3390/v9070174
13. Hoppe-Seyler K, Bossler F, Lohrey C, Bulkescher J, Rösl F, Jansen L, et al. Induction of dormancy in hypoxic human papillomavirus-positive cancer cells. *Proceedings of the National Academy of Sciences* 2017;114(6):E990-E8. doi: 10.1073/pnas.1615758114
14. Bossler F, Hoppe-Seyler K, Hoppe-Seyler F. PI3K/AKT/mTOR Signaling Regulates the Virus/Host Cell Crosstalk in HPV-Positive Cervical Cancer Cells. *International Journal of Molecular Sciences* 2019;20(9):2188. doi: 10.3390/ijms20092188
15. Bossler F, Kuhn BJ, Günther T, Kraemer SJ, Khalkar P, Adrian S, et al. Repression of human papillomavirus oncogene expression under hypoxia is mediated by PI3K/mTORC2/AKT signaling. *MBio* 2019;10(1):e02323-18. doi: 10.1128/mBio.02323-18
16. Li X, Zhao J, Yan T, Mu J, Lin Y, Chen J, et al. Cyanidin-3-O-glucoside and cisplatin inhibit proliferation and downregulate the PI3K/AKT/mTOR pathway in cervical cancer cells. *Journal of Food Science* 2021;86(6):2700-12. doi: 10.1111/1750-3841.15740

17. Zeng X, Wan L, Wang Y, Xue J, Yang H, Zhu Y. Effect of low dose of berberine on the radioresistance of cervical cancer cells via a PI3K/HIF-1 pathway under nutrient-deprived conditions. *International Journal of Radiation Biology* 2020;96(8):1060-7. doi: 10.1080/09553002.2020.1770358
18. Pezzuto A, Carico E. Role of HIF-1 in cancer progression: novel insights. A review. *Current molecular medicine* 2018;18(6):343-51. doi: 10.2174/1566524018666181109121849
19. Jia L, Hu Y, Yang G, Li P. Puerarin suppresses cell growth and migration in HPV-positive cervical cancer cells by inhibiting the PI3K/mTOR signaling pathway. *Experimental and Therapeutic Medicine* 2019;18(1):543-9. doi: 10.3892/etm.2019.7589
20. Marquard FE, Jücker M. PI3K/AKT/mTOR signaling as a molecular target in head and neck cancer. *Biochemical pharmacology* 2020;172:113729. doi: 10.1016/j.bcp.2019.113729
21. Devarajan N, Manjunathan R, Ganesan SK. Tumor hypoxia: the major culprit behind cisplatin resistance in cancer patients. *Critical Reviews in Oncology/Hematology* 2021;162:103327. doi: 10.1016/j.critrevonc.2021.103327
22. Dasari S, Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of action. *European journal of pharmacology* 2014;740:364-78. doi: 10.1016/j.ejphar.2014.07.025
23. Kim CW, Lu JN, Go S-I, Jung JH, Yi SM, Jeong J-H, et al. p53 restoration can overcome cisplatin resistance through inhibition of Akt as well as induction of Bax. *International journal of oncology* 2013;43(5):1495-502. doi: 10.3892/ijo.2013.2070
24. Zhang F, Duan S, Tsai Y, Keng PC, Chen Y, Lee SO, et al. Cisplatin treatment increases stemness through upregulation of hypoxia-inducible factors by interleukin-6 in non-small cell lung cancer. *Cancer science* 2016;107(6):746-54. doi: 10.1111/cas.12937
25. Chen X, Liu M, Meng F, Sun B, Jin X, Jia C. The long noncoding RNA HIF1A-AS2 facilitates cisplatin resistance in bladder cancer. *Journal of cellular biochemistry* 2019;120(1):243-52. doi: 10.1002/jcb.27327
26. Guzy RD, Hoyos B, Robin E, Chen H, Liu L, Mansfield KD, et al. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell metabolism* 2005;1(6):401-8.
27. Zonneveld MI, Keulers TG, Rouschop KM. Extracellular vesicles as transmitters of hypoxia tolerance in solid cancers. *Cancers* 2019;11(2):154. doi: 10.3390/cancers11020154
28. Sawant A, Kothandapani A, Zhitkovich A, Sobol RW, Patrick SM. Role of mismatch repair proteins in the processing of cisplatin interstrand cross-links. *DNA repair* 2015;35:126-36. doi: 10.1016/j.dnarep.2015.10.003
29. Lee J, Kim S. An overview of cyclophilins in human cancers. *Journal of International Medical Research* 2010;38(5):1561-74.
30. Li D, Hu C, Li H. Survivin as a novel target protein for reducing the proliferation of cancer cells. *Biomedical Reports* 2018;8(5):399-406. doi: 10.3892/br.2018.1077
31. Shimanishi M, Ogi K, Sogabe Y, Kaneko T, Dehari H, Miyazaki A, et al. Silencing of GLUT-1 inhibits sensitization of oral cancer cells to cisplatin during hypoxia. *Journal of oral pathology & medicine* 2013;42(5):382-8. doi: 10.1111/jop.12028
32. Mahapatra E, Das S, Biswas S, Ghosh A, Sengupta D, Roy M, et al. Insights of Cisplatin Resistance in Cervical Cancer: A Decision Making for Cellular Survival. *Cervical Cancer-A Global Public Health Treatise* 2021.
33. Zhang Y, Zhang Z. The history and advances in cancer immunotherapy: understanding the characteristics of tumor-infiltrating immune cells and their therapeutic implications. *Cellular & molecular immunology* 2020;17(8):807-21. doi: 10.1038/s41423-020-0488-6

34. Najafi M, Majidpoor J, Toolee H, Mortezaee K. The current knowledge concerning solid cancer and therapy. *Journal of biochemical and molecular toxicology* 2021;35(11):e22900. doi: 10.1002/jbt.22900
35. Li JJ, Liang Q, Sun GC. Traditional Chinese medicine for prevention and treatment of hepatocellular carcinoma: A focus on epithelial-mesenchymal transition. *J Integr Med* 2021;19(6):469-77. doi: 10.1016/j.joim.2021.08.004
36. Zhang Y, Huang R, Jiang Y, Shen W, Pei H, Wang G, et al. The role of bacteria and its derived biomaterials in cancer radiotherapy. *Acta Pharmaceutica Sinica B* 2022. doi: 10.1016/j.apsb.2022.10.013
37. Taefehshokr S, Parhizkar A, Hayati S, Mousapour M, Mahmoudpour A, Eleid L, et al. Cancer immunotherapy: Challenges and limitations. *Pathology-Research and Practice* 2022;229:153723. doi: 10.1016/j.prp.2021.153723
38. Sedighi M, Zahedi Bialvaei A, Hamblin MR, Ohadi E, Asadi A, Halajzadeh M, et al. Therapeutic bacteria to combat cancer; current advances, challenges, and opportunities. *Cancer medicine* 2019;8(6):3167-81. doi: 10.1002/cam4.2148
39. Vijayakumar M, Priya K, Tharmalingam N, Bose RJ, Madheswaran T, Ramasamy M. Bacteria therapeutics for cancer oncology: a crossroads for new paradigms. *Drug Discovery Today* 2022. doi: 10.1016/j.drudis.2022.03.007
40. Feng X, He P, Zeng C, Li Y-H, Das SK, Li B, et al. Novel insights into the role of Clostridium novyi-NT related combination bacteriolytic therapy in solid tumors. *Oncology Letters* 2021;21(2):1-. doi: 10.3892/ol.2020.12371
41. Wang L, Wang Q, Tian X, Shi X. Learning from Clostridium novyi-NT: How to defeat cancer. *Journal of cancer research and therapeutics* 2018;14(8):1. doi: 10.4103/0973-1482.204841
42. Patyar S, Joshi R, Byrav D, Prakash A, Medhi B, Das B. Bacteria in cancer therapy: a novel experimental strategy. *Journal of biomedical science* 2010;17(1):1-9. doi: 10.1186/1423-0127-17-21
43. DeClue AE, Axiak-Bechtel SM, Zhang Y, Saha S, Zhang L, Tung D, et al. Immune response to C. novyi-NT immunotherapy. *Vet Res* 2018;49(1):38. doi: 10.1186/s13567-018-0531-0
44. Bettegowda C, Huang X, Lin J, Cheong I, Kohli M, Szabo SA, et al. The genome and transcriptomes of the anti-tumor agent Clostridium novyi-NT. *Nature biotechnology* 2006;24(12):1573-80. doi: 10.1038/nbt1256
45. Kim J-E, Lee J-I, Jin D-H, Lee WJ, Park GB, Kim S, et al. Sequential treatment of HPV E6 and E7-expressing TC-1 cells with bortezomib and celecoxib promotes apoptosis through p-p38 MAPK-mediated downregulation of cyclin D1 and CDK2. *Oncology reports* 2014;31(5):2429-37. doi: /10.3892/or.2014.3082
46. Jafari FA, Abdoli A, Pilehchian R, Soleimani N, Hosseini SM. The oncolytic activity of Clostridium novyi nontoxic spores in breast cancer. *BioImpacts* 2021. doi: 10.34172/bi.2021.25
47. Perše M. Cisplatin Mouse Models: Treatment, Toxicity and Translatability. *Biomedicines* 2021;9(10). doi: 10.3390/biomedicines9101406
48. Schuetze S, Manig A, Ribes S, Nau R. Aged mice show an increased mortality after anesthesia with a standard dose of ketamine/xylazine. *Laboratory Animal Research* 2019;35(1):8. doi: 10.1186/s42826-019-0008-y
49. Babaei H, Alibabrdel M, Asadian S, Siavashi V, Jabarpour M, Nassiri SM. Increased circulation mobilization of endothelial progenitor cells in preterm infants with retinopathy of prematurity. *Journal of Cellular Biochemistry* 2018;119(8):6575-83. doi: 10.1002/jcb.27327

50. Liu X, Yang X, Chen F, Chen D. Combined application of Doxorubicin and Naringin enhances the antitumor efficiency and attenuates the toxicity of Doxorubicin in HeLa cervical cancer cells. *Int J Clin Exp Pathol* 2017;10(7):7303-11.
51. Vu M, Yu J, Awolude OA, Chuang L. Cervical cancer worldwide. *Current problems in cancer* 2018;42(5):457-65. doi: 10.1016/j.currproblcancer.2018.06.003
52. Kim BH, Chang JH. Differential effect of GLUT1 overexpression on survival and tumor immune microenvironment of human papilloma virus type 16-positive and-negative cervical cancer. *Scientific Reports* 2019;9(1):1-7. doi: /10.1038/s41598-019-49928-x
53. Dong D, Fu Y, Chen F, Zhang J, Jia H, Li J, et al. Hyperoxia sensitizes hypoxic HeLa cells to ionizing radiation by downregulating HIF-1 α and VEGF expression. *Molecular Medicine Reports* 2021;23(1):1-. doi: /10.3892/mmr.2020.11700
54. Sarmiento-Salinas FL, Perez-Gonzalez A, Acosta-Casique A, Ix-Ballote A, Diaz A, Treviño S, et al. Reactive oxygen species: Role in carcinogenesis, cancer cell signaling and tumor progression. *Life Sciences* 2021;284:119942. doi: 10.1016/j.lfs.2021.119942
55. Burd EM. Human papillomavirus and cervical cancer. *Clinical microbiology reviews* 2003;16(1):1-17.
56. Jalil AT, Karevskiy A. The cervical cancer (CC) epidemiology and human papillomavirus (HPV) in the middle east. *International Journal of Environment, Engineering and Education* 2020;2(2):7-12. doi: 10.55151/ijeedu.v2i2.29
57. Jing X, Yang F, Shao C, Wei K, Xie M, Shen H, et al. Role of hypoxia in cancer therapy by regulating the tumor microenvironment. *Molecular cancer* 2019;18(1):1-15. doi: 10.1186/s12943-019-1089-9
58. Zhang Y, Chen C, Zhang J. Effects and significance of formononetin on expression levels of HIF-1 α and VEGF in mouse cervical cancer tissue. *Oncology Letters* 2019;18(3):2248-53. doi: 10.3892/ol.2019.10567
59. Xie Y, Shi X, Sheng K, Han G, Li W, Zhao Q, et al. PI3K/Akt signaling transduction pathway, erythropoiesis and glycolysis in hypoxia. *Molecular medicine reports* 2019;19(2):783-91. doi: 10.3892/mmr.2018.9713
60. Federico C, Sun J, Muz B, Alhallak K, Cosper PF, Muhammad N, et al. Localized delivery of cisplatin to cervical cancer improves its therapeutic efficacy and minimizes its side effect profile. *International Journal of Radiation Oncology* Biology* Physics* 2021;109(5):1483-94. doi: 10.1016/j.ijrobp.2020.11.052
61. Rashid M, Zadeh LR, Baradaran B, Molavi O, Ghesmati Z, Sabzichi M, et al. Up-down regulation of HIF-1 α in cancer progression. *Gene* 2021;798:145796. doi: 10.2174/1566524018666181109121849
62. Han Y, Kim B, Cho U, Park IS, Kim SI, Dhanasekaran DN, et al. Mitochondrial fission causes cisplatin resistance under hypoxic conditions via ROS in ovarian cancer cells. *Oncogene* 2019;38(45):7089-105. doi: 10.1038/s41388-019-0949-5
63. Roberts NJ, Zhang L, Janku F, Collins A, Bai R-Y, Staedtke V, et al. Intratumoral injection of Clostridium novyi-NT spores induces antitumor responses. *Science translational medicine* 2014;6(249):249ra111-249ra111.
64. Dang LH, Bettgowda C, Huso DL, Kinzler KW, Vogelstein B. Combination bacteriolytic therapy for the treatment of experimental tumors. *Proceedings of the National Academy of Sciences* 2001;98(26):15155-60. doi: 10.1073/pnas.251543698
65. Zwagerman NT, Friedlander RM, Monaco III EA. Intratumoral Clostridium novyi as a potential treatment for solid necrotic brain tumors. *Neurosurgery* 2014;75(6):N17-N8. doi: 10.1227/01.neu.0000457197.94533.68

66. Zhang S, Chen Y, Wang J, Tang F, Miao T, Li M. Role of nontoxigenic *Clostridium novyi* in solid tumor therapy. *Reviews in Medical Microbiology* 2014;25(3):71-6.
67. Xu X, Liu T, Wu J, Wang Y, Hong Y, Zhou H. Transferrin receptor-involved HIF-1 signaling pathway in cervical cancer. *Cancer Gene Therapy* 2019;26(11):356-65. doi: 10.1038/s41417-019-0078-x
68. Kai L, Ling-ling M, Duo-ming W. Relationship between HPV infection and expression of HIF-1alpha and VEGF in head and neck squamous cell carcinoma and its prognostic value. *JOURNAL OF SHANGHAI JIAOTONG UNIVERSITY (MEDICAL SCIENCE)* 2013;33(6):806.
69. Staedtke V, Roberts NJ, Bai RY, Zhou S. *Clostridium novyi*-NT in cancer therapy. *Genes Dis* 2016;3(2):144-52. doi: 10.1016/j.gendis.2016.01.003
70. Patyar S, Joshi R, Byrav DP, Prakash A, Medhi B, Das B. Bacteria in cancer therapy: a novel experimental strategy. *Journal of biomedical science* 2010;17(1):1-9. doi: 10.1186/1423-0127-17-21




Figure 1: Histopathological observations of the therapeutic combination of *Clostridium novyi*-NT spores and cisplatin in TC-1 tumor-bearing mice: (a) - The results of H&E staining show that extensive necrosis has occurred in the tissue of tumor mice receiving the combination treatment (yellow box: extensive necrosis of tumor tissue). (b) - Giemsa staining, confirming the germination of *Clostridium novyi*-NT bacteria spores in treatment groups receiving spores (yellow arrow: bacterial accumulation). (c) - The comparison of the mitotic count of the tested groups shows that the amount of this factor is statistically significantly lower compared to the control group and the group receiving cisplatin. (d) - Extensive tumor necrosis of the tumor tissue in the group receiving the combined treatment, the amount of which has increased statistically significantly compared to the other groups and the control group. (* $P < 0.05$, ** $P < 0.005$, *** $P = 0.0001$, **** $P < 0.0001$)




Figure 2: Effects of *Clostridium novyi*-NT spores and cisplatin on the expression level of PLC and GLUT 1 in mice with cervical cancer. GAPDH was used as a housekeeping gene. The RT-qPCR results showed statistically significant differences in the mRNA expression levels of (a) GLUT 1 and (b) PLC in different treatment groups of mice compared to the control. Data demonstrate the mean \pm SD of three independent experiments. (* P < 0.05, ** P < 0.005, *** P = 0.0001, **** P < 0.0001)




Figure 3: Western blotting representative of HIF-1 α and VEGF of different treatment groups, β -actin was used as a normalizer. A statistically significant decrease in the expression of both proteins is evident in the group receiving the combined treatment of *Clostridium novyi*-NT spores and cisplatin compared to the control group. All the experiments were performed in triplicates. (* $P < 0.05$)




Figure 4: Comparison of intracellular levels of ROS in different treatment groups: The results showed that the number of reactive oxygen species (ROS) was significantly reduced in the group receiving cisplatin along with *Clostridium novyi*-NT spores compared to the group receiving cisplatin and *Clostridium novyi*- NT spores alone and compared to the control group (P = 0.0001)