

Original Article



Combining Chemotherapy Agents and Autophagy Modulators for Enhanced Breast Cancer Cell Death

Soraya Moomivand^{1,2}, Mohsen Nikbakht^{2,3*}, Ahmad Majd¹, Maryam Bikhof Torbati⁴, Seyyed Asadollah Mousavi²

¹Department of Biology, North Tehran Branch, Islamic Azad University, Tehran, Iran.

²Research Institute for Oncology, Hematology and Cell Therapy Tehran University of Medical Sciences, Tehran, Iran.

³Cell Therapy and Hematopoietic Stem Cell Transplantation Research Center, Tehran, Iran.

⁴Department of Biology, Yadegar-e-Imam Khomeini (RAH) Shahre rey branch, Islamic Azad University, Tehran, Iran.

Article info

Article History:

Received: January 24, 2024

Revised: October 18, 2024

Accepted: October 29, 2024

Published: October 30, 2024

Keywords:

Apoptosis, Arsenic trioxide, Autophagy, Breast cancer, Carboplatin, Cell lines, Cyclophosphamide

Abstract

Purpose: Autophagy, governed by genes with dual roles in cell death and survival, plays a crucial role in cancer persistence. Arsenic trioxide (ATO), carboplatin (CP), and cyclophosphamide (CY) are used to treat various cancers. ATO impedes cell proliferation and triggers apoptosis in cancer cells. CP, a platinum-based drug, damages cancer cell DNA, while CY acts as an alkylating agent, disrupting cell proliferation. This study investigates the combined effects of ATO, CP, and CY on inducing apoptosis and modulating autophagy in triple-negative breast cancer (TNBC) cell lines, BT-20 and MDA-MB-231.

Methods: The cytotoxic effects of ATO, CP, and CY, alone and in combination, were evaluated using the MTT assay on BT-20 and MDA-MB-231 cells. Apoptosis and cell cycle progression were analyzed by annexin-V FITC/PI staining and flow cytometry. Gene expression of autophagy- and apoptosis-related markers, including Beclin 1, LC3, caspase 3, and BCL2, was quantified using RT-PCR. Data were analyzed using GraphPad Prism 4.0 with one-way ANOVA followed by Dunnett's test.

Results: The combination of ATO, CP, and CY significantly reduced cell viability and enhanced apoptosis, evidenced by increased caspase-3 activity and reduced BCL2 expression. Cell cycle arrest in the G1 phase was observed, alongside elevated autophagy markers Beclin 1 and LC3.

Conclusion: The combination of ATO, CP, and CY induces synergistic effects in promoting apoptosis and autophagy in TNBC cell lines. These findings suggest that this combination therapy could be a promising approach to enhancing treatment efficacy in aggressive breast cancers, offering new insights into potential therapeutic strategies.

Introduction

Breast cancer remains the most prevalent form of cancer among women, affecting approximately 1.7 million women annually and accounting for 11.9% of the global cancer burden.¹⁻³ Moreover, breast cancer constitutes 30% of all cancers in women and contributes to 15%-20% of cancer-related mortalities in the female population.⁴ Survival rates for breast cancer exhibit significant global disparities, with the estimated 5-year survival rate reaching 80% in developed countries, compared to less than 40% in developing nations.⁵ Diagnosis of breast cancer involves assessing the status of several key receptors, including the estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2).⁶ Approximately 70-80% of breast cancers are hormone receptor-positive (ER, PR-positive), while 10-15% overexpress HER2, and another 10%-15% fall into the category of triple-negative breast cancers (TNBCs), which lack ER, PR, and HER2 expression.^{7,8}

Cell death is a fundamental physiological process that

governs the life cycle of an organism, including growth, development, aging, and eventual death.⁹ Two primary mechanisms of programmed cell death are autophagy and apoptosis.^{10,11} Autophagy, often described as a self-degradative process, plays a dual role in cancer cells by recycling cellular components to promote survival under stress or, when overly activated, leading to autophagic cell death.¹² Apoptosis, in contrast, is a highly regulated process that systematically eliminates damaged or malignant cells, acting as a safeguard against uncontrolled cell proliferation.^{13,14} Emerging evidence suggests that many anticancer therapies, including chemotherapy, can induce autophagy, which may contribute to both therapy resistance and tumor cell survival. As a result, targeting autophagy has been proposed as a therapeutic strategy to enhance the efficacy of cancer treatments, including in breast cancer.¹⁵

Arsenic, historically known for its toxicity, has emerged as a therapeutic agent in cancer treatment, particularly through arsenic trioxide (ATO), which is widely used for

*Corresponding Author: Mohsen Nikbakht, Email: m-nikbakht@sina.tums.ac.ir

its anti-cancer properties in treating acute promyelocytic leukemia (APL).¹⁶⁻¹⁸ ATO has shown promise not only in hematological malignancies but also in solid tumors due to its ability to induce apoptosis through oxidative stress and various signaling pathways.^{19,20} Alongside ATO, chemotherapeutic agents like carboplatin (CP) and cyclophosphamide (CY) are frequently employed in the management of breast cancer.²¹⁻²³ CP, a platinum-based compound, exerts its effects by cross-linking DNA, causing significant DNA damage that leads to apoptosis in cancer cells.²⁴ In contrast, CY functions as an alkylating agent, interfering with DNA replication and cell division, thereby inhibiting tumor growth.²⁵ The combination of these agents with novel therapeutic strategies, such as autophagy modulation, offers a promising avenue for enhancing treatment efficacy in aggressive forms of breast cancer. Kuo et al have documented the efficacy of ATO as a viable intervention for refractory or relapsed APL.²⁶ Therefore, it is imperative to conduct a thorough investigation into the amalgamation of ATO with other chemotherapeutic agents currently employed in clinical practice, to enhance its therapeutic effectiveness in the management of solid neoplasms in humans.

However, the role of autophagy in breast cancer is still controversial, with its implications varying depending on the context.²⁷ The aim of this study is to evaluate the combined effects of ATO, CP, and CY on the expression of autophagy-related genes such as LC3 and Beclin 1, as well as their impact on apoptosis in two TNBC cell lines: BT-20 and MDA-MB-231. This investigation will focus on determining whether varying concentrations and treatment durations of these agents enhance their cytotoxic effects, potentially offering new insights into therapeutic strategies for TNBC.

Materials and Methods

Chemicals and reagents

Fetal bovine serum (FBS), recombinant DMEM culture medium, Penicillin/streptomycin, and Trypsin-EDTA were obtained from Bio Idea Company (Tehran, Iran). Trypan Blue, chloroform, isopropanol, and DMSO were acquired from Merck (Germany). Hemocytometer slides and MTT powder were supplied by TGI (Germany) and Sigma (Germany), respectively. A manual RNA extraction solution and a cDNA synthesis kit were purchased from Genex Company (Iran). DEPC-treated water was sourced from Cinacoon (Iran). Master Mix for Real-Time PCR (2x) was provided by Amplicon Company (Denmark). All other unspecified reagents and materials were procured from Merck or Sigma.

Cell line and cell culture

The breast cancer cell lines BT-20 and MDA-MB-231 were acquired from the Pasteur Institute (Tehran, Iran). Both cell lines were cultured in DMEM supplemented with 10% FBS and incubated at 37 °C in a humidified atmosphere

containing 5% CO₂.

Analysis of cell viability by MTT assay

BT-20 and MDA-MB-231 cells were seeded into 96-well plates at a density of 5×10^4 cells per well. Cells were treated with ATO and combinations of ATO with other compounds, incubated at 37 °C with 5% CO₂ for 24 and 36 hours. The MTT assay was performed to assess cell proliferation, with results presented as a percentage relative to control cells treated with 0.1% DMSO. Cell viability was determined by measuring optical density (OD) at 570 nm using a plate reader. The absorbance ratio between treated and control cells was used to calculate cell viability using the following equation:

$$\text{cellviability}(\%) = \frac{(OD_{\text{sample}} - OD_{\text{negative control}})}{(OD_{\text{positive control}} - OD_{\text{negative control}})} \times 100$$

The analysis of the cell cycle

Cells were collected and preserved in 70% ethanol. Following preservation, the cells were treated with RNAaseI (10 µL/mL) and Triton X-100 (0.1%) for 30 minutes at 37 °C. Subsequently, cells were stained with propidium iodide (PI; 10 µg /mL), and flow cytometry was employed to assess the cell cycle. Data acquisition was performed using a BD flow cytometer, and analysis was conducted with the FlowJo software. The percentage of apoptotic cells was determined by evaluating the hypodiploid G0/G1 DNA fraction.

Analysis of cell apoptosis by flow cytometry

Cells were seeded into wells at a density of 15×10^5 cells per well and incubated for 36 hours. During this time, cells were treated with ATO and various ATO combinations or maintained as untreated controls. Apoptosis was assessed using a staining assay with Annexin V-FITC, following the manufacturer's instructions. The percentage of apoptotic cells was determined by analyzing Annexin V+/PI-cells using a BD flow cytometer and FlowJo software. The apoptosis analysis included the evaluation of early apoptosis (Annexin V+/PI-), late apoptosis (Annexin V+/PI+), and necrosis (Annexin V-/PI+).

RNA extraction and quantitative RT-PCR

Total RNA was extracted using TRI pure reagent (Roche Applied Science, Germany) according to the manufacturer's protocol. The RNA pellets were dissolved in DEPC-treated water, and RNA quality and concentration were measured using a Nanodrop ND-1000 spectrophotometer at 260 nm. cDNA was synthesized from 1-2 µg of total RNA using a cDNA synthesis kit (Takara Bio Inc., Otsu, Japan) following the manufacturer's instructions. The concentration of cDNA was standardized, and PCR was prepared using Beclin and LC3 primers (Table 1). Real-time PCR amplification was performed using the ABI Step One Plus™ system (Applied Biosystems, USA). The

Table 1. The primers for RT-PCR analysis

Genes	Primers	Reference
Beclin	F: 5'-AGCTGCCGTTACTGTTCTG-3'	28
	R: 5'-ACTGCCTCTGTGTCTTCAATCTT-3'	
LC3	F: 5'-GATGTCCGACTTATTCGAGAGC-3'	29
	R: 5'-TTGAGCTGTAAGCGCCTTCTA-3'	
Caspase 3	F: 5'-GACTCTGGAATATCCCTGGACAACA-3'	30
	R: 5'-AGGTTTGTGTCATCGACATCTG-3'	
BCL2	F: 5'-CTGCACCTGACGCCCTTACC-3'	31
	R: 5'-CACATGACCCACCGAACTCAAAGA-3	
GAPDH	F: 5'-TGAACGGGAAGCTCACTGG-3'	32
	R: 5'-TCCACCACCTGTGCTGTA-3'	

relative expression of selected genes was determined using the comparative Ct method. Five groups were analyzed, and gene quantification was conducted using the Rotor-Gene Q system (QIAGEN, Hilden, Germany) with the primers listed in Table 1.

Statistical analysis

The data were analyzed using GraphPad Prism 4.0 (GraphPad Software, La Jolla, CA). One-way analysis of variance (ANOVA) followed by Dunnett's test was performed to compare group means. Results were expressed as mean values with standard error (SE). Statistical significance was considered at a threshold of $P < 0.05$.

Results

ATO, CP, and CY inhibit cell proliferation

The cytotoxic effects of ATO, CP, and CY on MDA-MB-231 and BT-20 cells were assessed through the MTT assay to determine the impact of these treatments on cellular viability. The cell viability for both MDA-MB-231 and BT-20 cell lines was evaluated at two distinct time points: 24 hours and 36 hours. In the MDA-MB-231 cell line, CP administered at 1250 μM resulted in a substantial reduction in cell viability, declining from 90.67% at 24 hours to 80.29% at 36 hours. When the concentration of CP was increased to 5000 μM , significant cytotoxic effects were observed, with cell viability decreasing to 57.09% at 24 hours and further declining to 49.57% at 36 hours. Other treatments, such as CY at 200 μM and ATO at 2 μM , maintained relatively high levels of viability, with ATO at 2 μM showing minimal reduction from 89.93% at 24 hours to 88.41% at 36 hours.

Notably, combination treatments of ATO 2 μM +CP 1250 μM resulted in a pronounced reduction in cell viability, from 85.07% to 52.75%, underscoring the synergistic cytotoxic effects. Particularly, the combination of ATO 3 μM +CP 5000 μM caused a marked decrease in viability, from 40.67% to 35.94% (Figure 1 and 2). In the BT-20 cell line, similar trends were observed, with CY 500 μM leading to a significant reduction in viability,

from 55.12% at 24 hours to 50.39% at 36 hours. ATO 2 μM alone maintained relatively high viability at 87.35% at 24 hours but exhibited notable declines when combined with other treatments, such as ATO 2 μM +CY 200 μM , where viability decreased from 72.83% to 69.56%. The most substantial reductions were observed in combination treatments, including ATO 3 μM +CY 200 μM , which dropped from 40.37% to 33.19%, and ATO 3 μM +CP 5000 μM , which declined from 37.62% to 33.26% (Figure 1 and 2).

Apoptosis assay

To assess whether the cytotoxic effects of ATO, CP, and CY, both individually and in combination, were mediated through apoptosis or necrosis, flow cytometry analysis was conducted on BT-20 and MDA-MB-231 cell lines. Cells were treated for 36 hours and stained with Annexin-V FITC/PI, enabling differentiation between apoptotic and necrotic populations. As depicted in Figure 3a for the BT-20 cell line, 87.33% of cells in the control group were classified as viable, with a total apoptosis rate of 10.12% and necrotic cells accounting for 2.48%. In the CP treatment group, viability decreased to 63.47%, with a total apoptosis rate of 25.09% and an increase in necrotic cells to 11.77%. The CY treatment group demonstrated similar viability at 65.17%, but with a higher total apoptosis rate of 32.43% and a lower necrotic cell count of 2.35%. ATO treatment preserved a relatively higher viability at 72.33%, with a total apoptosis rate of 27.60% and minimal necrosis at 0.13%. In contrast, the combination treatment of CY and ATO resulted in a pronounced decline in viability to 28.10%, with total apoptosis reaching 57.30% and necrotic cells accounting for 14.60%. Finally, the CP+ATO combination treatment yielded the lowest viability at 25.27%, with a total apoptosis rate of 44.93% and a significant increase in necrotic cells to 29.80%.

For MDA-MB-231 cells, in the control group, viability remained high at 95.1%, with a total apoptosis rate of only 3.28% (Figure 3b). In contrast, treatment with CP caused a substantial reduction in viability to 25.23%, alongside a total apoptosis rate of 43.17%. The CY treatment group exhibited higher viability at 56.50%, with a total apoptosis rate of 33.63%. ATO treatment maintained a relatively higher viability at 67.30%, with a total apoptosis rate of 31.30%. The combination treatment of CY and ATO further reduced viability to 31.13%, with total apoptosis increasing to 48.47%. Lastly, the CP+ATO combination treatment resulted in 28.13% viability and a total apoptosis rate of 52.40%. Based on these data, the selected doses for both cell lines were ATO at 2 μM , CP at 5000 μM , and CY at 200 μM .

Cell cycle assay

The effects of ATO and CP on cell cycle progression were analyzed via flow cytometry (Figures 4a and 4b). In the BT-20 cell line, the impact of ATO combined with CP on




Figure 1. Effect of ATO, CP, and CY on cell viability in BT-20 and MDA-MB-231 cell lines, assessed using the MTT assay after 24 and 36 hours of treatment. Significant differences in cell viability were observed between the 24-hour and 36-hour treatments. (A) Effect of ATO on BT-20 cells; (B) effect of CP on BT-20 cells; (C) effect of CY on BT-20 cells; (D) effect of ATO on MDA-MB-231 cells; (E) effect of CP on MDA-MB-231 cells; and (F) effect of CY on MDA-MB-231 cells. Data are presented as mean \pm standard error (SE) from three independent experiments. Statistical significance was denoted as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ relative to the corresponding control

cell cycle distribution was assessed and compared to the control group. The results revealed significant alterations in the percentage of cells in each phase of the cell cycle following treatment. In the control group, an average of 1.025% of cells were in the G₀ phase. In contrast, the ATO + CP treatment group showed a substantial increase to 7.295%, indicating a marked induction of quiescence or cell cycle arrest. In the G₁ phase, the control group had an average of 34.545%, which increased to 43.525% in the ATO + CP treatment group. This increase suggests that the

combination therapy may enhance the transition of cells into the G₁ phase, potentially delaying their progression to the S phase. The S phase exhibited a decrease in cell population from 24.265% in the control group to 21.11% following ATO + CP treatment, suggesting that the combined treatment may inhibit DNA synthesis or disrupt cell progression through the S phase. Lastly, the G₂ phase showed a decline from 40.155% in the control group to 28.05% in ATO + CP-treated cells.

In MDA-MB-231 cells, the control group exhibited an




Figure 2. Cell viability in BT-20 and MDA-MB-231 cells was assessed using the MTT assay. The effects of ATO + CP and ATO + CY treatments were examined after 24 and 36 hours. (A) The impact of ATO + CP on BT-20 cells; (B) the impact of ATO + CY on BT-20 cells; (C) the impact of ATO + CP on MDA-MB-231 cells; (D) the impact of ATO + CY on MDA-MB-231 cells. Data are presented as mean \pm SE of three independent experiments. Statistical significance was denoted as * P < 0.05, ** P < 0.01, *** P < 0.001, and **** P < 0.0001 relative to the corresponding control

average of 8.405% of cells in the G0 phase, which increased significantly to 14.915% following ATO + CP treatment, indicating a marked induction of quiescence or cell cycle arrest, thereby promoting dormancy in the treated cells. In the G1 phase, the control group had an average of 60.925%, which decreased significantly to 43.295% in the ATO + CP treatment group. This reduction suggests that the combination therapy may disrupt the transition of cells into the G1 phase, potentially resulting in the accumulation of cells in the G0 phase and inhibiting further progression through the cell cycle. The S phase showed a minor decrease, with the control group averaging 22.425%, and the ATO + CP treatment group at 22.265%. This slight reduction indicates that the combined treatment does not substantially inhibit DNA synthesis or significantly disrupt the overall progression of cells through the S phase, suggesting that replication processes remain largely unaffected. However, the G2 phase exhibited a notable increase from an average of 8.225% in the control group to 19.505% in ATO + CP-treated cells. This significant rise suggests that the treatment may impede the progression of cells into mitosis, causing an accumulation of cells in the G2 phase, and potentially delaying cell cycle progression, which could have important implications for the efficacy of the combination treatment.

A real-time assay for polymerase chain reaction

To investigate the mechanisms underlying the observed

effects of ATO, CP, and CY, real-time PCR was employed to assess the expression levels of Beclin 1, Caspase 3, LC3, and BCL2 genes. As shown in Figure 5a and 5b, Beclin 1 expression significantly increased by 1.34-fold and 1.21-fold in the ATO-treated group compared to the control in BT-20 and MDA-MB-231 cells, respectively. In contrast, the CP and CY groups exhibited non-significant changes. The combination of ATO with CP or CY resulted in significant increases in Beclin 1 expression in both cell lines, with the ATO + CP group exhibiting a 1.54-fold increase in BT-20 cells and a 1.44-fold increase in MDA-MB-231 cells.

Figure 5c and 5d show Caspase 3 gene expression in both cell lines. ATO treatment led to a significant increase in Caspase 3 expression by 1.43-fold in BT-20 cells, while CP and CY treatments resulted in non-significant increases of 1.13-fold and 1.26-fold, respectively. The ATO + CP combination significantly increased Caspase 3 expression by 1.65-fold, and ATO + CY resulted in a 1.33-fold increase. In MDA-MB-231 cells, ATO significantly increased Caspase 3 expression by 1.35-fold, with significant increases also observed in the ATO + CP and ATO + CY groups (1.56-fold and 1.41-fold, respectively).

Changes in LC3 gene expression are presented in Figure 5e and 5f. BT-20 cells treated with ATO exhibited a 1.25-fold increase in LC3 expression compared to the control group, with similar increases observed in the CP and CY groups. The ATO + CP group demonstrated a




Figure 3. Assessment of apoptosis and necrosis using flow cytometry in (A) BT-20 cells and (B) MDA-MB-231 cells after 36 hours of treatment. Early apoptotic cells are represented in the lower right quadrant, while post-apoptotic necrotic cells are indicated in the upper right quadrant. Data are presented as mean \pm SE from three independent experiments. Statistical significance was denoted as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ relative to the corresponding control

1.35-fold increase in LC3 expression, while the ATO + CY group showed a 1.25-fold increase. In MDA-MB-231 cells, LC3 expression increased slightly with ATO (1.12-fold), CP (1.15-fold), and CY (1.06-fold) treatments, though these increases were not significant. However, significant increases were observed in the ATO + CP (1.84-fold) and ATO + CY (1.55-fold) groups.

Lastly, the changes in BCL2 gene expression across different treatment groups are shown in Figure 5g and 5h. In BT-20 cells, ATO treatment resulted in a significant 0.64-fold decrease in BCL2 expression, whereas the decreases in the CP (0.84-fold) and CY (0.85-fold) groups were not statistically significant. However, the ATO + CP and ATO + CY combinations led to significant decreases in BCL2 expression (0.64-fold and 0.76-fold, respectively). Similarly, in MDA-MB-231 cells, ATO reduced BCL2 expression by 0.82-fold, while the CP and CY groups showed non-significant decreases. Notably, the combination treatments resulted in significant decreases in BCL2 expression in MDA-MB-231 cells, with the ATO + CP group reducing expression by 0.75-fold and the ATO + CY group by 0.78-fold.

Discussion

Breast cancer development results from complex alterations in gene expression, occurring through genetic and epigenetic changes.³³ TNBC, which lacks targeted hormonal therapies, poses significant challenges due to its aggressive nature. Therefore, investigating the impact of various drugs on cellular pathways is critical for identifying effective treatments.³⁴ In this study, we evaluated the effects of ATO, CP, and CY on the expression of key autophagy-related genes (Beclin 1, caspase 3, LC3, and BCL2) in BT-20 and MDA-MB-231 TNBC cell lines. Autophagy plays a crucial role in maintaining cellular homeostasis by degrading damaged proteins and organelles. Disruption of autophagy has been implicated in cancer progression, invasion, and metastasis, particularly in aggressive cancers such as TNBC.³⁵

ATO, a well-known inducer of both autophagy and apoptosis, has garnered significant attention due to its ability to modulate multiple molecular pathways, including the intrinsic and extrinsic apoptotic pathways, as well as autophagic mechanisms.³⁶ The present study demonstrated that ATO exhibited dose- and time-dependent cytotoxicity




Figure 4. (A) Cell cycle analysis of BT-20 cells following ATO and CP combination treatment, showing alterations in cell cycle distribution. (B) Cell cycle analysis of MDA-MB-231 cells following ATO and CP combination treatment, showing a reduction in the G1 phase. Data are presented as mean \pm SE of three independent experiments

in both TNBC cell lines, a finding consistent with its known mechanism of action. Specifically, ATO promotes oxidative stress and mitochondrial dysfunction, leading to the activation of apoptosis through caspase-dependent pathways.³⁷ Our flow cytometry analysis revealed a marked decrease in viable cell populations and an increase in both early and late apoptotic cells, further confirming the induction of apoptosis by ATO.

Autophagy was also significantly upregulated in response to ATO treatment, as evidenced by increased expression of Beclin 1 and LC3. This finding is consistent with previous studies demonstrating that ATO induces autophagy as part of its cytotoxic mechanism, particularly in cancer cells where autophagy functions as a survival response to cellular stress.^{38,39} Chiu et al showed that ATO could synergistically induce both apoptosis and autophagy, a concept supported by our data.^{40,41} In another study, Kanzawa et al stated that ATO activates multiple intracellular signaling pathways leading to apoptosis, differentiation enhancement, or inhibition of angiogenesis.⁴² However, the role of autophagy in promoting or inhibiting cell death remains controversial, as autophagy can either protect cancer cells from chemotherapy-induced stress or lead to autophagic cell death under certain conditions.

The present study also evaluated the combined effects of ATO with CP and CY on apoptosis and autophagy. Both CP and CY are alkylating agents that induce DNA

cross-linking, leading to DNA damage and the activation of the DNA damage response (DDR) pathway. The DDR subsequently activates p53 and caspase-dependent apoptotic pathways, contributing to cell cycle arrest and apoptosis.⁴³ The potential of CP as an effective anti-cancer drug, particularly for APL, was investigated by Hu et al, who found that combination therapy of high doses of ATO and CP had a synergistic effect and induced apoptosis in various solid tumor cancer cell lines; however, it is crucial to acknowledge that this treatment approach may have notable adverse effects on patients.⁴⁴ Our results demonstrated that the combination of ATO with CP or CY significantly enhanced apoptosis, as reflected by the increased activation of caspase-3 and the downregulation of BCL2.⁴⁵ Caspase-3 is a key executioner caspase in the intrinsic apoptotic pathway, responsible for cleaving various substrates and driving the cell toward programmed cell death.^{46,47} The activation of caspase-3 in both TNBC cell lines highlights the intrinsic apoptotic commitment induced by this drug combination. Moreover, BCL2 is an anti-apoptotic protein that plays a pivotal role in regulating mitochondrial outer membrane permeabilization (MOMP) and preventing cytochrome c release, thus inhibiting apoptosis.⁴⁸ The combination of ATO with CP or CY appears to potentiate these effects, resulting in a robust pro-apoptotic response. Additionally, the observed reduction in BCL2 expression in both cell lines suggests that this combination therapy may sensitize




Figure 5. Effect of ATO, CP, CY, and their combinations on the mRNA expression levels of specific genes in BT-20 and MDA-MB-231 cells. The study assessed the effects of these treatments on the expression of (A, B) Beclin 1, (C, D) caspase 3, (E, F) LC3, and (G, H) BCL2 in BT-20 and MDA-MB-231 cells. Data are presented as mean \pm SE from three independent experiments. Statistical significance was denoted as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ relative to the corresponding control

TNBC cells to chemotherapy-induced DNA damage, further enhancing apoptosis.

In conclusion, the combination of ATO with CP or CY exerts a synergistic effect on apoptosis and autophagy in TNBC cell lines, offering a promising therapeutic approach for treating aggressive breast cancers. The dual induction of autophagy and apoptosis, coupled

with the downregulation of BCL2 and the activation of caspase-3, highlights the potential of this combination therapy to overcome the limitations of current treatments. Future studies should investigate the precise molecular mechanisms underlying these synergistic effects and evaluate their clinical applicability in TNBC and other difficult-to-treat cancers.

Conclusion

This study demonstrated that the combination of ATO, CP, and CY significantly reduced the survival and proliferation of BT-20 and MDA-MB-231 cells in a dose- and time-dependent manner. The therapy enhanced apoptosis by increasing caspase-3 activation and reducing BCL2 expression, key indicators of the intrinsic apoptotic pathway. The addition of CP and CY to ATO amplified these effects, highlighting the synergistic potential of this combination in promoting cancer cell death. The reduction in BCL2 further sensitized the cells to apoptosis, supporting the use of this combination therapy as a promising approach for breast cancer treatment.

Acknowledgments

This work was supported by the Hematology, Oncology and Cell Therapy Tehran University of Medical Sciences, Tehran, Iran.

Author's Contribution

Conceptualization: Mohsen Nikbakht.

Data curation: Soraya Moomivand, Mohsen Nikbakht.

Formal analysis: Soraya Moomivand, Mohsen Nikbakht, Seyed Asadollah Mousavi.

Funding acquisition: Mohsen Nikbakht, Seyed Asadollah Mousavi.

Investigation: Soraya Moomivand.

Methodology: Soraya Moomivand.

Project administration: Mohsen Nikbakht.

Resources: Mohsen Nikbakht, Ahmad Majd, Seyed Asadollah Mousavi, Maryam Bikhof Torbati.

Software: Soraya Moomivand.

Supervision: Mohsen Nikbakht, Ahmad Majd, Maryam Bikhof Torbati, Seyed Asadollah Mousavi.

Validation: Soraya Moomivand.

Visualization: Soraya Moomivand.

Writing—original draft: Soraya Moomivand, Ahmad Majd, Maryam Bikhof Torbati, Seyed Asadollah Mousavi, Mohsen Nikbakht.

Writing—review & editing: Mohsen Nikbakht, Ahmad Majd.

Competing Interests

The authors declare that they have no conflict of interest.

Ethical Approval

Not applicable.

Funding

This work was funded by the Hematology, Oncology and Cell Therapy Research Center Tehran University of Medical Sciences, Tehran, Iran.

References

- Davey MG, Lowery AJ, Miller N, Kerin MJ. MicroRNA expression profiles and breast cancer chemotherapy. *Int J Mol Sci* 2021;22(19):10812. doi: [10.3390/ijms221910812](https://doi.org/10.3390/ijms221910812)
- Niazvand F, Wagh PR, Khazraei E, Borzouyan Dastjerdi M, Patil C, Najari IA. Application of carbon allotropes composites for targeted cancer therapy drugs: a review. *J Compos Compd* 2021;3(7):140-51. doi: [10.52547/jcc.3.2.7](https://doi.org/10.52547/jcc.3.2.7)
- Sabouri P, Bahadori S, Mohsennejadesaniyani A, Hosseini SM, Khodamorovati M, Yazdani E, et al. Recent advances in nanocarrier-based targeted drug delivery: for lung, colon, and breast cancers. *J Compos Compd* 2023;5:74-90. doi: [10.61186/jcc.5.2.3](https://doi.org/10.61186/jcc.5.2.3)
- Łukasiewicz S, Czeczelewski M, Forma A, Baj J, Sitarz R, Stanislawek A. Breast cancer-epidemiology, risk factors, classification, prognostic markers, and current treatment strategies—an updated review. *Cancers (Basel)* 2021;13(17):4287. doi: [10.3390/cancers13174287](https://doi.org/10.3390/cancers13174287)
- Akram M, Iqbal M, Daniyal M, Khan AU. Awareness and current knowledge of breast cancer. *Biol Res* 2017;50(1):33. doi: [10.1186/s40659-017-0140-9](https://doi.org/10.1186/s40659-017-0140-9)
- Vohra P, Buelow B, Chen YY, Serrano M, Vohra MS, Berry A, et al. Estrogen receptor, progesterone receptor, and human epidermal growth factor receptor 2 expression in breast cancer FNA cell blocks and paired histologic specimens: a large retrospective study. *Cancer Cytopathol* 2016;124(11):828-35. doi: [10.1002/cncy.21745](https://doi.org/10.1002/cncy.21745)
- Hoda RS, Brogi E, Pareja F, Nanjangud G, Murray MP, Weigelt B, et al. Secretory carcinoma of the breast: clinicopathologic profile of 14 cases emphasising distant metastatic potential. *Histopathology* 2019;75(2):213-24. doi: [10.1111/his.13879](https://doi.org/10.1111/his.13879)
- Kumar RV, Panwar D, Amirtham U, Premalata CS, Gopal C, Narayana SM, et al. Estrogen receptor, progesterone receptor, and human epidermal growth factor receptor-2 status in breast cancer: a retrospective study of 5436 women from a regional cancer center in South India. *South Asian J Cancer* 2018;7(1):7-10. doi: [10.4103/sajc.sajc_211_17](https://doi.org/10.4103/sajc.sajc_211_17)
- Muñoz-Pinedo C. Signaling pathways that regulate life and cell death: evolution of apoptosis in the context of self-defense. *Adv Exp Med Biol* 2012;738:124-43. doi: [10.1007/978-1-4614-1680-7_8](https://doi.org/10.1007/978-1-4614-1680-7_8)
- Shimizu S, Yoshida T, Tsujioka M, Arakawa S. Autophagic cell death and cancer. *Int J Mol Sci* 2014;15(2):3145-53. doi: [10.3390/ijms15023145](https://doi.org/10.3390/ijms15023145)
- Zhao J, Jiang P, Guo S, Schrodi SJ, He D. Apoptosis, autophagy, NETosis, necroptosis, and pyroptosis mediated programmed cell death as targets for innovative therapy in rheumatoid arthritis. *Front Immunol* 2021;12:809806. doi: [10.3389/fimmu.2021.809806](https://doi.org/10.3389/fimmu.2021.809806)
- Russell RC, Guan KL. The multifaceted role of autophagy in cancer. *EMBO J* 2022;41(13):e110031. doi: [10.15252/emboj.2021110031](https://doi.org/10.15252/emboj.2021110031)
- Braicu C, Zanoaga O, Zimta AA, Tigau AB, Kilpatrick KL, Bishayee A, et al. Natural compounds modulate the crosstalk between apoptosis- and autophagy-regulated signaling pathways: controlling the uncontrolled expansion of tumor cells. *Semin Cancer Biol* 2022;80:218-36. doi: [10.1016/j.semcancer.2020.05.015](https://doi.org/10.1016/j.semcancer.2020.05.015)
- Cocco S, Leone A, Piezzo M, Caputo R, Di Lauro V, Di Rella F, et al. Targeting autophagy in breast cancer. *Int J Mol Sci* 2020;21(21):7836. doi: [10.3390/ijms21217836](https://doi.org/10.3390/ijms21217836)
- Sui X, Chen R, Wang Z, Huang Z, Kong N, Zhang M, et al. Autophagy and chemotherapy resistance: a promising therapeutic target for cancer treatment. *Cell Death Dis* 2013;4(10):e838. doi: [10.1038/cddis.2013.350](https://doi.org/10.1038/cddis.2013.350)
- Ota A, Wahiduzzaman M, Hosokawa Y. Arsenic-based anticancer-combined therapy: novel mechanism inducing apoptosis of cancer cells. In: *Current Understanding of Apoptosis: Programmed Cell Death*. IntechOpen; 2018. p. 33-54. doi: [10.5772/intechopen.74824](https://doi.org/10.5772/intechopen.74824)
- Akhtar A, Xiaoyan Wang S, Ghali L, Bell C, Wen X. Recent advances in arsenic trioxide encapsulated nanoparticles as drug delivery agents to solid cancers. *J Biomed Res* 2017;31(3):177-88. doi: [10.7555/jbr.31.20160059](https://doi.org/10.7555/jbr.31.20160059)
- Haghi A, Salami M, Mohammadi Kian M, Nikbakht M, Mohammadi S, Chahardouli B, et al. Effects of sorafenib and arsenic trioxide on U937 and KG-1 cell lines: apoptosis or autophagy? *Cell J* 2020;22(3):253-62. doi: [10.22074/cellj.2020.6728](https://doi.org/10.22074/cellj.2020.6728)
- Kumar S, Yedjou CG, Tchounwou PB. Arsenic trioxide induces oxidative stress, DNA damage, and mitochondrial pathway of apoptosis in human leukemia (HL-60) cells. *J Exp Clin Cancer*

- Res 2014;33(1):42. doi: [10.1186/1756-9966-33-42](https://doi.org/10.1186/1756-9966-33-42)
20. Ahmad IM, Dafferner AJ, Salloom RJ, Abdalla MY. Heme oxygenase-1 inhibition modulates autophagy and augments arsenic trioxide cytotoxicity in pancreatic cancer cells. *Biomedicines* 2023;11(9):2580. doi: [10.3390/biomedicines11092580](https://doi.org/10.3390/biomedicines11092580)
 21. Jiang L, Lee SC, Ng TC. Pharmacometabonomics analysis reveals serum formate and acetate potentially associated with varying response to gemcitabine-carboplatin chemotherapy in metastatic breast cancer patients. *J Proteome Res* 2018;17(3):1248-57. doi: [10.1021/acs.jproteome.7b00859](https://doi.org/10.1021/acs.jproteome.7b00859)
 22. Suryawanshi RV. Assessment of efficacy and toxicity of cyclophosphamide chemotherapy in canines with malignant mammary tumor: a retrospective study. *Vet Med Int* 2021;2021:5520603. doi: [10.1155/2021/5520603](https://doi.org/10.1155/2021/5520603)
 23. Skverchinskaya E, Levdarovich N, Ivanov A, Mindukshev I, Bukatin A. Anticancer drugs paclitaxel, carboplatin, doxorubicin, and cyclophosphamide alter the biophysical characteristics of red blood cells, in vitro. *Biology (Basel)* 2023;12(2):230. doi: [10.3390/biology12020230](https://doi.org/10.3390/biology12020230)
 24. Szeffler B, Czeleń P, Krawczyk P. The affinity of carboplatin to B-vitamins and nucleobases. *Int J Mol Sci* 2021;22(7):3634. doi: [10.3390/ijms22073634](https://doi.org/10.3390/ijms22073634)
 25. Mills KA, Chess-Williams R, McDermott C. Novel insights into the mechanism of cyclophosphamide-induced bladder toxicity: chloroacetaldehyde's contribution to urothelial dysfunction in vitro. *Arch Toxicol* 2019;93(11):3291-303. doi: [10.1007/s00204-019-02589-1](https://doi.org/10.1007/s00204-019-02589-1)
 26. Kuo CC, Liu TW, Chen LT, Shiah HS, Wu CM, Cheng YT, et al. Combination of arsenic trioxide and BCNU synergistically triggers redox-mediated autophagic cell death in human solid tumors. *Free Radic Biol Med* 2011;51(12):2195-209. doi: [10.1016/j.freeradbiomed.2011.09.023](https://doi.org/10.1016/j.freeradbiomed.2011.09.023)
 27. Janser FA, Adams O, Büttler V, Schläfli AM, Dislich B, Seiler CA, et al. Her2-targeted therapy induces autophagy in esophageal adenocarcinoma cells. *Int J Mol Sci* 2018;19(10):3069. doi: [10.3390/ijms19103069](https://doi.org/10.3390/ijms19103069)
 28. Zhou L, Lv MQ, Ge P, Yang YQ, He DL, Wang HX, et al. The expression of Beclin-1 in testicular tissues of non-obstructive azoospermia patients and its predictive value in sperm retrieval rate. *Transl Androl Urol* 2021;10(8):3267-74. doi: [10.21037/tau-21-320](https://doi.org/10.21037/tau-21-320)
 29. Zhao Y, Yang J, Liao W, Liu X, Zhang H, Wang S, et al. Cytosolic FoxO1 is essential for the induction of autophagy and tumour suppressor activity. *Nat Cell Biol* 2010;12(7):665-75. doi: [10.1038/ncb2069](https://doi.org/10.1038/ncb2069)
 30. Tao S, Ren Z, Yang Z, Duan S, Wan Z, Huang J, et al. Effects of different molecular weight polysaccharides from *Dendrobium officinale* Kimura & Migo on human colorectal cancer and transcriptome analysis of differentially expressed genes. *Front Pharmacol* 2021;12:704486. doi: [10.3389/fphar.2021.704486](https://doi.org/10.3389/fphar.2021.704486)
 31. Cianfrocca R, Tocci P, Semprucci E, Spinella F, Di Castro V, Bagnato A, et al. β -Arrestin 1 is required for endothelin-1-induced NF- κ B activation in ovarian cancer cells. *Life Sci* 2014;118(2):179-84. doi: [10.1016/j.lfs.2014.01.078](https://doi.org/10.1016/j.lfs.2014.01.078)
 32. Kong X, Xu X, Yan Y, Guo F, Li J, Hu Y, et al. Estrogen regulates the tumour suppressor miRNA-30c and its target gene, MTA-1, in endometrial cancer. *PLoS One* 2014;9(3):e90810. doi: [10.1371/journal.pone.0090810](https://doi.org/10.1371/journal.pone.0090810)
 33. Baxter E, Windloch K, Gannon F, Lee JS. Epigenetic regulation in cancer progression. *Cell Biosci* 2014;4:45. doi: [10.1186/2045-3701-4-45](https://doi.org/10.1186/2045-3701-4-45)
 34. Döhner H, Weisdorf DJ, Bloomfield CD. Acute myeloid leukemia. *N Engl J Med* 2015;373(12):1136-52. doi: [10.1056/NEJMra1406184](https://doi.org/10.1056/NEJMra1406184)
 35. Bar-Yosef T, Damri O, Agam G. Dual role of autophagy in diseases of the central nervous system. *Front Cell Neurosci* 2019;13:196. doi: [10.3389/fncel.2019.00196](https://doi.org/10.3389/fncel.2019.00196)
 36. Haghi A, Salemi M, Fakhimamadi A, Mohammadi Kian M, Yousefi H, Rahmati M, et al. Effects of different autophagy inhibitors on sensitizing KG-1 and HL-60 leukemia cells to chemotherapy. *IUBMB Life* 2021;73(1):130-45. doi: [10.1002/iub.2411](https://doi.org/10.1002/iub.2411)
 37. Zhang G, Liu J, Zhang Y, Qu J, Xu L, Zheng H, et al. Cbl-dependent degradation of FLIP(L) is involved in ATO-induced autophagy in leukemic K562 and gastric cancer cells. *FEBS Lett* 2012;586(19):3104-10. doi: [10.1016/j.febslet.2012.07.067](https://doi.org/10.1016/j.febslet.2012.07.067)
 38. Nakaoka T, Ota A, Ono T, Karnan S, Konishi H, Furuhashi A, et al. Combined arsenic trioxide-cisplatin treatment enhances apoptosis in oral squamous cell carcinoma cells. *Cell Oncol (Dordr)* 2014;37(2):119-29. doi: [10.1007/s13402-014-0167-7](https://doi.org/10.1007/s13402-014-0167-7)
 39. Walker AM, Stevens JJ, Ndebele K, Tchounwou PB. Evaluation of arsenic trioxide potential for lung cancer treatment: assessment of apoptotic mechanisms and oxidative damage. *J Cancer Sci Ther* 2016;8(1):1-9. doi: [10.4172/1948-5956.1000379](https://doi.org/10.4172/1948-5956.1000379)
 40. Chiu HW, Ho YS, Wang YJ. Arsenic trioxide induces autophagy and apoptosis in human glioma cells in vitro and in vivo through downregulation of survivin. *J Mol Med (Berl)* 2011;89(9):927-41. doi: [10.1007/s00109-011-0763-1](https://doi.org/10.1007/s00109-011-0763-1)
 41. Chiu HW, Ho SY, Guo HR, Wang YJ. Combination treatment with arsenic trioxide and irradiation enhances autophagic effects in U118-MG cells through increased mitotic arrest and regulation of PI3K/Akt and ERK1/2 signaling pathways. *Autophagy* 2009;5(4):472-83. doi: [10.4161/auto.5.4.7759](https://doi.org/10.4161/auto.5.4.7759)
 42. Kanzawa T, Zhang L, Xiao L, Germano IM, Kondo Y, Kondo S. Arsenic trioxide induces autophagic cell death in malignant glioma cells by upregulation of mitochondrial cell death protein BNIP3. *Oncogene* 2005;24(6):980-91. doi: [10.1038/sj.onc.1208095](https://doi.org/10.1038/sj.onc.1208095)
 43. Türk G, Ceribaşı AO, Sakin F, Sönmez M, Ateşşahin A. Antiperoxidative and anti-apoptotic effects of lycopene and ellagic acid on cyclophosphamide-induced testicular lipid peroxidation and apoptosis. *Reprod Fertil Dev* 2010;22(4):587-96. doi: [10.1071/rd09078](https://doi.org/10.1071/rd09078)
 44. Hu WC, Teo WH, Huang TF, Lee TC, Lo JF. Combinatorial low dose arsenic trioxide and cisplatin exacerbates autophagy via AMPK/STAT3 signaling on targeting head and neck cancer initiating cells. *Front Oncol* 2020;10:463. doi: [10.3389/fonc.2020.00463](https://doi.org/10.3389/fonc.2020.00463)
 45. Singh N, Nigam M, Ranjan V, Sharma R, Balapure AK, Rath SK. Caspase mediated enhanced apoptotic action of cyclophosphamide- and resveratrol-treated MCF-7 cells. *J Pharmacol Sci* 2009;109(4):473-85. doi: [10.1254/jphs.08173fp](https://doi.org/10.1254/jphs.08173fp)
 46. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH. Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol* 2013;14:32. doi: [10.1186/1471-2121-14-32](https://doi.org/10.1186/1471-2121-14-32)
 47. Inoue S, Browne G, Melino G, Cohen GM. Ordering of caspases in cells undergoing apoptosis by the intrinsic pathway. *Cell Death Differ* 2009;16(7):1053-61. doi: [10.1038/cdd.2009.29](https://doi.org/10.1038/cdd.2009.29)
 48. Kalkavan H, Green DR. MOMP, cell suicide as a BCL-2 family business. *Cell Death Differ* 2018;25(1):46-55. doi: [10.1038/cdd.2017.179](https://doi.org/10.1038/cdd.2017.179)