

Review Article



Lipid-Based Nanoparticles as Oral Drug Delivery Systems: Overcoming Poor Gastrointestinal Absorption and Enhancing Bioavailability of Peptide and Protein Therapeutics

Soheil Mehrdadi^{ID}

Department of Pharmaceutical and Pharmacological Sciences, University of Padova, Padua, Italy.

Article info

Article History:

Received: November 7, 2022

Revised: August 9, 2023

Accepted: October 8, 2023

published: October 14, 2023

Keywords:

Peptide and protein therapeutics, Lipid-based drug delivery systems, Solid lipid nanoparticles, Nanostructured lipid carriers, Oral drug delivery

Abstract

Delivery and formulation of oral peptide and protein therapeutics have always been a challenge for the pharmaceutical industry. The oral bioavailability of peptide and protein therapeutics mainly relies on their gastrointestinal solubility and permeability which are affected by their poor membrane penetration, high molecular weight and proteolytic (chemical and enzymatic) degradation resulting in limited delivery and therapeutic efficacy. The present review article highlights the challenges and limitations of oral delivery of peptide and protein therapeutics focusing on the application, potential and importance of solid lipid nanoparticles (SLNs) and nanostructured lipid carriers (NLCs) as lipid-based drug delivery systems (LBDDSs) and their advantages and drawbacks. LBDDSs, due to their lipid-based matrix can encapsulate both lipophilic and hydrophilic drugs, and by reducing the first-pass effect and avoiding proteolytic degradation offer improved drug stability, dissolution rate, absorption, bioavailability and controlled drug release. Furthermore, their small size, high surface area and surface modification increase their mucosal adhesion, tissue-targeted distribution, physiological function and half-life. Properties such as simple preparation, high-scale manufacturing, biodegradability, biocompatibility, prolonged half-life, lower toxicity, lower adverse effects, lipid-based structure, higher drug encapsulation rate and various drug release profile compared to other similar carrier systems makes LBDDSs a promising drug delivery system (DDS). Nevertheless, undesired physicochemical features of peptide and protein drug development and discovery such as plasma stability, membrane permeability and circulation half-life remain a serious challenge which should be addressed in future.

Introduction

In recent decades, high demands for cost-benefit healthcare expenses, efficient therapeutics (e.g. safety and efficacy) and non-stop generic substitution has urged pharmaceutical companies,¹ industry and market to shift to biotechnology-driven peptide and protein therapeutics and biopharmaceuticals. These novel categories of drugs, unlike active pharmaceutical ingredients (APIs, drugs), offer better feedback due to higher potency, selectivity and specificity for their extracellular target.²

Peptides and proteins, as cell products, have various physiological functions in body such as hormones, enzyme substrates and inhibitors, antibiotics, biological regulators, structural components, signaling factors and catalyzers which all implies their importance in body; hence, any abnormality in their amino acid sequence or structural disfunction leads to sever diseases and pathological conditions; diabetes,³ dwarfism,⁴ cystic fibrosis,⁵ thalassemia⁶ or impaired blood clotting,⁷ among many others.^{8,9} as such, due to their biological specificity and efficient affinity and efficacy, peptides and proteins

have been exploited as drugs for treatment of diseases (Figure 1).¹⁰

Stability in proteins is the result of balance among four destabilizing and stabilizing forces: electrostatic interactions, hydrogen bonding, van der Waals forces and hydrophobic interactions, which form the secondary, tertiary and quaternary structures of proteins and any disruption will influence the structural balance and destabilizes that particular protein.^{11,12} Different environmental factors can influence the chemical and physical stability of proteins such as pH, ionic strength, temperature, high pressure, non-aqueous solvents, metal ions, detergents, adsorption, agitation and shearing which all are inevitably part of the manufacturing, sterilization and lyophilization process, and consequently might damage the developing protein resulting in biological inactivation, aggregation, immunogenicity and precipitation.¹³⁻¹⁵

Peptide and protein therapeutics are highly dependent on the production process, yet are biocompatible, cost-benefit, with modifiable in-vivo bioactivity, specific

*Corresponding Author: Soheil Mehrdadi, Email: soheilmehrdadi@gmail.com

© 2024 The Author (s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.




Figure 1. Biotechnology medicines in research and development (R&D) by therapeutic category

targeting, chemical diversity, and easily synthesized by using solid-phase peptide synthesis methodologies (e.g. Merrifield's method) in which the amino acid sequence can be precisely chosen and inserted at the molecular level by modifying the basic units.¹⁶ nevertheless, undesired physicochemical features of protein drug development and discovery, remains a serious challenge for formulation scientists as well as pharmaceutical and biotechnology companies.

Only recently peptides have been considered as therapeutic agents while they were never considered as a potential therapeutic agent¹⁷; mostly due to their protease degradation, metabolic instability, short half-life, manufacturing complications and high expenses, which in long-term administration renders them unfavorable in terms of patient costs and compliance especially with regard to parenteral administration as the majority of peptides (10%) have a very low oral bioavailability.¹⁸ On the other hand, peptides' biodegradability into non- to low-toxicity metabolites,² low drug-drug interactions and immunogenicity,² higher tissue penetration (owing to their small size), higher in-vivo activity (per unit mass), stability and lower expenses favors them over large therapeutic proteins and antibodies for regulatory approval (higher than 20%) which is twice the rate of small molecules.¹⁷

As well as peptides, proteins also gained importance and their application in pharmaceutical science and industry was emphasized due to more advanced analytical methods resulting in the recognition of various peptides and hormones as therapeutic biopharmaceuticals, novel genetic and molecular engineering methods to produce large-scale proteins and recently-defined roles of proteins as regulatory components of numerous diseases.^{19,20} Since then, pharmaceutical industry has developed various large-scale oral delivery technologies for peptides/proteins as active ingredients.²¹

Low oral bioavailability of peptide and protein

therapeutics renders them being formulated as parenteral preparations; large molecular size,²² susceptibility to enzymatic degradation (local or mostly GI tract), poor stability in the gastric acidic environment,²³ poor intestinal penetration, short plasma half-life, immunogenicity and the propensity to aggregation, adsorption, and denaturation.^{24,25} Enzymatic degradation and poor intestinal penetration, among all, have been mainly mentioned for low oral bioavailability and short half-life of protein drugs (<1%) which is claimed to be increased to 30%-50% by pharmaceutical enterprises.^{26,27} Pharmaceutical therapeutic proteins, due to their large molecular size which leads to low blood absorption and diffusion, requires specific epithelial transporters otherwise they cannot enter the general circulation by the ordinary routes of drugs absorption. Furthermore, low pH and protease enzymes of GI aggravate this condition even more.²⁸

Among different administration routes, the most common route for peptide and protein therapeutics is intravenous (I.V.) injections (Table 1), which is not favorable in terms of patient compliance, clearance varies from a few minutes to several days, and might result in undesired deposition and distribution which require repeated injections with higher therapeutic doses to achieve efficacy,^{29,30} subsequently causing severe adverse effects.

The subcutaneous (S.C.) and intramuscular (I.M.) injections are other administration routes (Table 1), the former is the most common one, especially for vaccines. Different factors such as molecular weight, site of injection, local injection site activity and pathological conditions can influence the fate of peptide and protein therapeutics following S.C. injection leading to a bioavailability as high as 100% or much lower.⁴⁶ upon injection, proteins with high molecular weight (<16000 Da) are absorbed either from vessels' endothelial cells to capillaries or reach the

Table 1. Various peptides and proteins' administration routes

Method	Delivery routes	Formulation/Device requirement	Reference
Invasive	Intravenous (I.V.), subcutaneous (S.C.), intramuscular (I.M.), Intracerebral vein (I.C.V.)	Liquid or reconstituted solid (syringe), i.v. injected liposomes	31-35
	Depot system (S.C. or I.M.)	Biodegradable polymers, liposomes, permeable polymers (not degradable) microspheres, implants	31-33,36,37
Non-invasive	Pulmonary	Liquid or powder formulations, nebulizers, metered dose inhalers, dry powder inhalers	31-34,38
	Oral	Solids, emulsions, microparticles, nanoparticles, with or without absorption enhancers	31-33,38,39
	Nasal	Liquid, usually requires permeation enhancers, nanoparticles	31-33,40,41
	Transdermal	Iontophoresis, electroporation, chemical permeation enhancers, prodrugs, sonophoresis, transfersomes	31-33,42,43
	Buccal, rectal, vaginal, ocular	Gels, suppositories, bioadhesives, microparticles	31-33,44,45

local lymphatic system and through thoracic duct join the blood circulation, while proteins with small molecular weight are mainly diffused through the local capillaries.⁴⁶ however, protein transportation through lymphatic way is undesired due to slow time of circulation which might result in protein enzymatic degradation.^{46,47}

Non-invasive routes have increasingly been investigated as well as an alternative to the conventional invasive injectable routes (Table 1). Numerous studies have investigated nasal, ophthalmic, buccal, rectal, vaginal, transdermal and pulmonary routes for peptide and protein delivery.^{28,48-55}

Based on studies, mucosae which so far have been neglected for drug delivery seem to be a promising approach for drug absorption, especially efficient for biomolecules of large size and molecular weight.^{28,56} The advantages of mucosal surfaces (mouth, eye, nose, rectum and vagina) for drug delivery over skin and GI tract can be named as: fewer biological barriers to pass for systemic diffusion, rapid absorption and evading hepatic first-pass effect. However, one practical challenge of mucosae is related to the preparations that are formulated for local long-term treatment.

Despite alternate routes of drug delivery, oral delivery (P.O.) is still the most preferred one; non-invasive, painlessness, easy self-administration, low risk of cross-infection, high patient convenience/compliance, outpatient feasibility,⁵⁷ cost-benefit (no need for sterile manufacturing).⁵⁷ Oral route does not offer the drawbacks of I.V. route; drug extravasation from blood, catheter-related infectious complications, thrombosis and being expensive and invasive, especially for chronic conditions.

However, biological barriers of GI tract with their associated enzymatic and chemical processes hamper the efficiency of oral route for drug delivery. Furthermore, the epithelial cell monolayer membrane of the GI tract even more aggravate the condition of low permeability for many peptide and protein therapeutics with low gastrointestinal solubility which finally results in low bioavailability.⁵⁸

Some active moieties cannot be delivered through oral route.^{59,60} According to the Biopharmaceutic Classification System (BCS),⁶¹ oral bioavailability of each

drug is determined by its solubility along the GI tract and cellular penetration. Most of the potential drug candidates developed with high-throughput screening methods generally have higher molecular weights and tend to be lipophilic in nature.⁶² Other factors contributing to low oral bioavailability of drugs are low stability in the gastrointestinal environment and poor membrane permeability. Most of drugs are substrates to intestinal efflux transporters like p-glycoprotein resulting in poor oral bioavailability.⁶³

Regardless of the administration route, majority of peptide and protein therapeutics, lacking necessary physicochemical requirements, fail to diffuse and be absorbed to their target tissue which implies the importance of drug delivery and tissue-targeting systems for achieving as maximum therapeutic effects as possible. The carrier systems diffuse and distribute the intended therapeutic molecules to their targeted site with as maximum concentration as possible in the affected area and as minimum concentration as possible in the intact tissues to lower the general adverse effects.⁶⁴

Peptide and Protein Drug Delivery

Introduction of novel biotechnological molecules as potential therapeutics, advent of chemical synthesis methods and recombinant DNA technology have all rendered protein synthesis and delivery an important area of research which resulted in the production of numerous large-scales drugs of peptide/protein origin such as monoclonal antibodies, hormones and vaccines. According to The 2018 and 2019 PhRMA reports,⁶⁵ there have been respectively 4751 and 5422 novel biotechnological medicines in research and development (R&D) phase for more than 100 diseases such as cancer, infectious diseases, autoimmune diseases, AIDS/HIV, antiparasitic and related conditions (Figure 1), which have been either in human clinical trials or under review by the Food and Drug Administration (FDA).⁶⁵ nevertheless, abovementioned challenges for their delivery through GI tract⁶⁶⁻⁷⁰ and the blood brain barrier (BBB)(in the case of central nervous system diseases),^{71,72} makes their therapeutic potential and clinical application questionable.

In the last decades, numerous drugs of peptide/protein origin have been in preclinical studies and clinical trials,⁷³ more than 400 recombinant peptides and proteins and 1300 under clinical trials.⁷⁴ The reason could be attributed mostly to the larger size of peptides and proteins comparing to conventional drugs, which provides drug-target interaction with binding pockets that are not normally available to small molecular drugs. These targets could be part of intracellular protein-protein interaction network which have been recognized in numerous diseases. Peptide and protein therapeutics in order to interact with such targets must penetrate cells, however, most of them are known to have extracellular targets,⁷³ and are parenterally administered so cellular penetration is not their ordinary route as it is for mucosal surfaces. Currently, the main obstacle of the oral administration of these novel categories of drugs for their maximum therapeutic effects could be addressed as the penetration through intestinal cellular membranes and target cellular membranes.

Transport mechanisms in the GI tract

To formulate and synthesize drug delivery systems (DDSs) for oral peptide and protein therapeutics, and biopharmaceuticals a throughput understanding of the biological pathways involved for their absorption and diffusion in the GI tract is necessary and worthwhile. Various physicochemical features govern the pathway through which the molecules will be penetrating the intestinal cells; molecular weight, hydrophobicity/hydrophilicity, ionization constants, and pH stability, among all.

Paracellular transport

It has the following features as, space dimension of 10 Å, aqueous pores (epithelial tight junctions) 7–9, 3–4 and 8–9 Å for the jejunum, the ileum and the colon respectively,⁷⁵ to allow the passage of solutes with a specific molecular radius and tight junctions building 0.01% of the total absorption surface area of the intestine.⁷⁶ These data prove the restriction of the paracellular transport toward the passing molecules (Table 2). however, there is an electrical resistance diversity and consequently ionic selectivity. In the latter case, also transcellular pathway's collaboration adjusts rate and selectivity of export of ions and solutes

and overall tissue-specific transport. The tight junction along with ion channels are involved in size and charge selectivity, ion concentration-dependent penetration, competitive-based penetration among different molecules, unordinary mole-fraction effects and pH-sensitivity.⁷⁷ hydrogen bonding capacity and lipophilicity do not influence much the paracellular pathway.

Transcellular transport

It is an endocytic process at apical membrane and the absorbed molecules are released at the basolateral membrane, glucose is also transported with this mechanism (Table 2). The protein-lipid ratio is very insignificant in the basolateral membrane due to its thinner and more permeable structure than the apical membrane. This transport mechanism is governed by various factors: different physicochemical properties of molecules, size, lipophilicity/hydrophobicity, hydrogen bond formation, surface charge, superficial ligands; the physiological condition of the GI system and the animal models studied for transport mechanism.^{89,90} There are mainly two types of primary intestinal epithelial cells for molecules transportation; Enterocytes and M cells, the former lining about 99% of the GI tract and the latter mainly the area of Peyer's patches and the human follicle-associated epithelium (FAE) (antigen-specialized).⁹¹ M cells function as presenting and transporting peptides and proteins to the local lymphoid tissues for immune reactions and a vulnerable and available way for pathogenic organisms.⁹² Due to their great endocytosis and transcytosis capacity for transporting diverse molecules and biomaterials (e.g. nanoparticles),^{93,94} M cells could be used for oral delivery of peptides and proteins and finally through phagocytosis, adsorptive (through clathrin-coated pits and vesicles) and fluid phase endocytosis they adsorb macromolecules and microorganisms.⁹⁵ Some studies demonstrated nanoparticles transportation through intestinal villi and contradicting the recent debates over the rate of particle absorption.^{96,97} There is a consensus on the transportation of the majority of particles in FAE,^{96,98,99} for which there have been studies on the Peyer's patches and M cells involvement on various biomaterials absorbency. The transcellular mechanism however is not a desirable route for low molecular-weight lipophilic drugs. Overall, absorption by this mechanism is reduced in a great extent

Table 2. Transport mechanisms in the GI tract

Transport Mechanisms	Process	Feature of Molecules	Note
Paracellular	Passive diffusion in intercellular spaces between epithelial cells, tissue-specific transport ^{78,79}	Ions, large substances and solutes < 15 Å (3.5 kDa) ⁸⁰	Restricted protein delivery due to tight junctions ⁸¹
Transcellular	Intestinal transcytosis, Enterocytes and M cells ⁸²	Various physicochemical properties	Limited transport of relatively low molecular-weight lipophilic drugs
Carrier-mediated	Across the cell membrane or entire cell ⁸³	Utilized by small hydrophilic molecules ⁸⁴	Small di-/tripeptides monosaccharides, and amino acids are transported transcellularly ⁸⁵
Receptor-mediated	Receptor specific ligand, ⁸⁶ endocytosis ⁸⁷	The physicochemical and metabolic features	Direct delivery of hydrophilic ligands to liver, direct delivery of lipophilic ligands to the vena cava ⁸⁸

in the colon part of the large intestines in comparison to the paracellular mechanisms.¹⁰⁰

Carrier-mediated transport

It is an active and energy-dependent transportation of specific molecules against their concentration gradient through specific membrane receptors, such as β -lactam antibiotics and angiotensin converting enzyme inhibitors, monosaccharides and amino acids (Table 2). In one study using Caco-2 cell monolayers, it was proved that the conjugated insulin is transported 5 to 15 times more through the transferrin receptor than then insulin receptor itself.¹⁰¹

Receptor-mediated transport

It has been investigated to evaluate the oral bioavailability of peptide and protein drugs by modifying receptor specific ligands-drug interaction. This mechanism has functions in different processes such as endocytosis (clathrin-mediated), phagocytosis, pinocytosis and potocytosis (nonclathrin-mediated) (Table 2). The absorption starts with the binding of molecules to their specific receptors and their internalization into endosomes with low acidic pH which might dissociate receptor-ligand bound and accordingly degrades endosomes. The absorbed peptide and protein access into systemic blood circulation with two distinct pathways: hepatic portal vein and intestinal lymphatic vessels, the amount of peptides and proteins absorbed through either of these two pathways depends greatly on the physicochemical features of the formulation. portal vein is the main pathway for the majority of orally administered peptide and protein drugs and through which hydrophilic molecules are absorbed and transported to the blood systemic circulation first through the hepatic portal vein and then by the hepatic artery, and finally they are delivered to their sites of action. But lipophilic molecules penetrating through the same intestinal barriers are transported to the intestinal lymphatic vessels, bypassing the first-pass effect, and directly delivered to the vena cava for blood systemic circulation.

Absorption of oral drugs

Drugs in order to be absorbed through GI tract are required to have high solubility and permeability, however, this is not the case for numerous drugs with low aqueous solubility and consequently low and diverse bioavailability.¹⁰² for such drugs simultaneous presence of high amount of fat through meals can increase their oral bioavailability,¹⁰³⁻¹⁰⁵ via prolonging GI tract passage time, exocrine pancreas secretion stimulation, reduced metabolism, lymphatic-associated absorption, increased intestinal penetration, reduced cellular efflux and liver- and mesenteric-related blood alteration.^{106,107}

Introduction of lipid-based drug delivery systems (LBDDSs) in 1990s provided scaffolds which increase dissolution rate of poor aqueous soluble drugs (i.e.

hydrophobic drugs) by providing a phase in which the drugs can disintegrate and be absorbed and diffused toward its site of action.¹⁰⁸ after the degradation of lipids in the intestine, active mono- and diglycerides are formed on the surface of lipids which later disassociate and transform into micelles and simultaneously drug is also solubilized inside micelles. These micelle-drug mixtures are finally absorbed.¹⁰⁹⁻¹¹²

The absorbed components through intestinal segment of GI system follow two distinct pathways according to their features: blood vessels and lymphatic vessels. The former is the preferred route for most of the oral drugs by which they are absorbed into the systemic circulation via portal vein, and the latter for highly lipophilic drugs ($\log P > 5$) by which drugs are absorbed into systemic circulation via lymphatic vessels.

Overall, the presence of lipid increases absorption of numerous drugs more through lymphatic vessels, especially lipophilic ones and macromolecules of high molecular weight,^{113,114} mostly due to their higher permeability to nanoparticles than blood vessels,¹¹⁵ and most importantly overpassing hepatic first-pass effect.¹¹⁶ Nevertheless, it has been demonstrated that the absorption through lymphatic pathway is affected by the length of fatty acid chains; long-chain triglycerides (14-18 chains) are more favored for absorption than low-chain ones.^{111,117}

Physiological barriers against peptide and protein therapeutics absorption

Gastrointestinal barriers

In-depth studies of molecular pharmacology have provided a better understanding of the biological and molecular processes of the GI system, while its target sites has introduced new insights for targeted delivery of oral peptide and protein therapeutics and biopharmaceuticals.^{118,119} There are two types of proteolytic enzymes with their specific site of action which are responsible partly for the physiological processes of GI system toward peptides and proteins: *endoproteases*, including trypsin, chymotrypsin, and elastase, which hydrolyze the internal bonds of the peptide chain to the amino- and carboxy-terminus, and *exopeptidases*, including carboxypeptidase A and aminopeptidase, which hydrolyze the amino- and carboxy-terminus bonds to the peptide chain (Table 3).¹²⁰ Enzymatic degradation happens at the lumen, brush border, the cytosol of the enterocytes, in the lysosomes and other cell organelles.¹²¹

The secretions of stomach (hydrochloric acid, potassium chloride and sodium chloride) provide an acidic pH of 1.5-3.5 for the proteolysis of peptides and proteins by breaking them down into amino acids, dipeptides and tripeptides for absorption. The digestion of peptides and proteins starts with pepsin in the acidic environment of stomach (pH 2), however, the alkaline environment of the intestines (pH 6) inactivates pepsin. The instantaneous wide change of pH from acidic stomach to alkaline

Table 3. Proteases and their sites of action

Types	Enzymes	Major Site of Action
Gastric proteases	Pepsins (aspartic proteases)	Broad activity, hydrolyzes numerous peptide bonds
Brush border proteases	Aminopeptidase A Aminopeptidase N Aminooligopeptidase Dipeptidylaminopeptidase IV Carboxypeptidase	Aminopeptidases are N-terminopeptidases, degrading mostly 3–10 amino acid residue-dipeptides and amino acids
Cytosolic proteases	Di- and tripeptidase	2-3 Amino peptide amino acids
Intestinal pancreatic proteases	Trypsin (endopeptidase) α -chymotrypsin (endopeptidase) Elastase (endopeptidase) Carboxypeptidases (exopeptidase)	Peptide bonds of basic amino acids/peptides Peptide bonds of hydrophobic amino acids/peptides Peptide bonds of smaller and nonaromatic amino acids/peptides A: C-terminal amino acid B: C-terminal basic amino acid
Brush border proteases	Aminopeptidase A Aminopeptidase N Amino oligopeptidase Dipeptidyl aminopeptidase IV Carboxypeptidase	Aminopeptidases are N-terminopeptidases, degrading mostly 3–10 amino acid residue-dipeptides and amino acids

intestines influences the degradation of ingested peptides and proteins and might contribute to their precipitation that redissolve following pH change.¹²²⁻¹²⁴

The small intestine is the major site of absorption along the GI tract due to the higher enzymatic activity of proteases (mostly in duodenum and jejunum). The brush borders of the intestinal epithelial cells secrete numerous specific enzymes (e.g. sucrose) leading to peptide/protein absorption and degradation.¹²⁰ furthermore, the digestive secretions of the exocrine part of the pancreas also contain endo-/exopeptidases which are released into duodenum to increase pH for the activity of intestinal enzymes (e.g. trypsin). However, in the terminal parts of jejunum and ileum the enzymatic activity of aminopeptidases is decreased to 20–30% where Peyer's patches are located and these areas could be a potential site for peptide/protein drug delivery.^{98,125}

Mucosal barriers

Mucus has a major importance and function by determining the absorption and bioavailability of administered drugs especially through oral route. The mucosal surface of stomach has three components which further hamper the drug diffusion and absorption as an exogenous component: the first one, which is lined by surface epithelial cells and tight junctions, has a role against irritant and unsuitable fluids; the second one has a very unique insoluble protective mucus (the mixture of surface epithelial cells and neck cell) which creates a jelly-like layer throughout the entire surface mucosae of the stomach; and the third one is composed of bicarbonate ions which are secreted by the surface epithelial cells.^{126,127}

There are however other components which further disturb peptide/protein absorption through oral administration, such as glycocalyx located on the surface layer of the stomach epithelial cells with an acidic nature and containing sulfated mucopolysaccharides. Goblet cells of the stomach wall secrete mucus, which covers upper layer of glycocalyx,¹²⁸ and contains mucin glycoproteins,

enzymes, electrolytes and water.¹²⁹ the mucin glycoprotein gives glycocalyx an adhesive feature and functions more as a physical barrier than a chemical one,^{130,131} and in the stomach and colon parts of GI tract has a thick layer while in the small intestine part is thinner and this fact can be justified according to the digestive functions of each segment.¹³² Overall, the abovementioned components altogether are protected by viscoelastic layers. Peptide and proteins first of all are required to pass the outermost layers, mucus and glycocalyx, to reach the cellular membranes which present a viscous barrier to absorption and diffusion.

Nanomedicines – novel drug delivery systems

After administrating drugs, their plasma concentration increases and they affect their site of action effectively followed by a gradual decrease to a point when their plasma concentration is not sufficient enough anymore to elicit their intended therapeutic effect. Hence, Drugs are required to be re-administrated based on the dose and frequency of administration to provide the same concentration which must be neither higher nor lower than their therapeutic concentration level; a concept widely known as “therapeutic window”; higher doses will result in general toxic effects and lower ones cannot elicit any therapeutic effects.¹³³ the conventional drugs do not have prolonged drug delivery features in their “therapeutic window” and this issue implies the importance of novel DDSs to be introduced.

A novel interdisciplinary branch of biomedical science, “nanomedicine”, has been investigating the potential application of biomaterials and nanostructures as DDSs for sustained, controlled and tissue-targeting delivery of drugs and active agents, which due to their pharmaceutical features overcome some of the limitations of conventional drugs.

Studies have proved that “drug discovery” alone does not offer practical solutions to therapeutics as most of the successful in-vitro experiments result in failure in in-vivo

experiments mostly due to: poor drug concentration owing to low absorption, quick metabolism and elimination (peptides and proteins); general blood distribution which results in drug-related adverse effects and toxicity (anti-cancer drugs); low drug solubility of aqueous solutions when administered intravenously; unpredictable bioavailability and high plasma-level variations with oral administration and physiological processes involving a drug's plasma level (e.g. food on cyclosporine).

In DDSs the in-vivo destiny of drugs is directly influenced by a series of factors which can be modified for in-site delivery with the desired therapeutic concentration. Peptide and protein therapeutics are encapsulated in nanomedicines for transportation through GI tract offering benefits such as high stability for storage and administration, and large-scale sterile manufacturing for oral preparations.¹³⁴ Nanomedicines can be formulated and modified with the desired criteria such as size, surface properties and release profile to have tissue-targeted delivery within drug's unique therapeutic window.

Nanomedicines have a size ranging from 1 to 100 nm where the therapeutic molecules can be incorporated in the core, matrix or attached on the surface (in the case of high surface/volume ratio),¹³⁵ which the latter results in longer half-life and systemic circulation and increased mean residence time (MRT).¹³⁶ Since 1990s LBDDSs have been under investigation owing to their advantages; biocompatibility, higher penetration capacity, lipophilicity with no need for surface modification, simple fabrication, cost benefit and industrial-scale production compared to their counterpart DDSs such as polymeric and inorganic nanomedicines.^{137,138}

Based on the fabrication methods and physicochemical properties LBDDSs are classified into the following:

Liposomes in the form of spherical vesicles which are composed of one or multiple lipid bilayer (phospholipid or natural phospholipids) enclosing an aqueous core.^{137,138} Their size varies from 10 to 1000 nm and they are the first generation of LBDDSs that were employed mainly for parenteral route of drug delivery. They possess low antigenicity and toxicity, high drug encapsulation and loading efficiency and sustained and controlled drug release as their benefits. However, their synthesis is complex and has low stability and rapid reticuloendothelial system clearance and scale-up problems. Despite introducing various surface functionalization (antibodies and peptides) and modification (e.g. PEG coating) which enhances their blood half-time, structural stability and therapeutic efficiency, they are still undesirable in terms of industrial scale production.¹³⁹⁻¹⁴¹ Currently, numerous liposome-based formulations are approved for a variety of diseases: Doxorubicin for cancers (Doxil[®], Myocet[®] and Lipodox[®]), Amphotericin B for fungal infections (Ambisome[®]), Cytarabine for lymphomatous meningitis (Depocyt[®]), Morphine sulfate for pain management (DepoDur[®]), inactivated hepatitis A virus (strain RG-SB

for hepatitis A (Epaxal[®]) and inactivated hemagglutinin of Influenza virus strains A and B for influenza (Inflexal[®]).¹⁴²

Lipoplexes are fabricated from liposomes and are designed in multi-lamellar lipoplexes with positive lipid bilayer and distinct negative nucleic acids. The electrostatic process between self-assembly liposomes and nucleic acids produces such scaffolds. Given their similarity to liposomes they possess similar benefits and drawbacks with a few of their own; multiple cations tendency to bind negative nucleic acids which decreases the internal cell transfection process. They have been investigated for brain-focused studies.¹⁴³

Natural body lipoproteins are another lipid-based system which are very similar to liposomes hence sharing the same advantages and disadvantages and carry lipids (mainly cholesterol), proteins, enzymes, and miRNAs. They have been investigated with other nanoparticles (e.g., albumin, PEG-PLGA) for numerous central nervous system diseases.¹⁴⁴

(2) Niosomes are vesicles with a lamellar self-assembled structure which are composed of non-ionic surfactants and cholesterol or its derivatives.¹⁴⁵ They can be encapsulated by lipophilic and hydrophilic substances. They are cheaper in terms of production and more stable than liposomes.¹⁴⁶

(3) Transferosomes which are composed of a lipid bilayer fabricated by a lipid matrix stabilized by various surfactants,¹⁴⁷ and are similar to niosomes and to liposomes.

(4) Solid lipid nanoparticles (SLNs) which are composed of a solid lipid core.¹⁴⁸

(5) Nanostructured lipid carriers (NLCs) which are composed of a liquid lipid phase core within the solid lipid phase.¹⁴⁹

Among all the lipid-based DDSs, this review will focus further on SLNs and NLCs and their unique characteristics and properties. SLNs, comparing to NLCs which are formulated with both solid and liquid oils, are formulated only with solid ones which gives them more controlled drug release due to limited drug mobility in solid lipids and are designed as oral pellets and retard capsule (e.g. Mucosolvan[®]), as microparticles by spray drying,¹⁵⁰ and oral nanopellets.¹⁵¹

Solid lipid nanoparticles and nanostructured lipid carriers

SLNs and NLCs are lipid-based colloidal drug carriers, synthesized from lipids (solid or liquid), surfactants, co-surfactants and APIs (drugs). SLNs are composed of solid lipids and surfactants but NLCs are also composed of liquid lipids and oils. Lipids have solid form both at room and body temperature and could be chosen as purified triglycerides, glyceride mixtures and waxes. Surface surfactants increase and improve the stability and cellular permeability and consequently absorption.^{152,153}

They collectively offer the benefits of other colloidal DDSs (e.g. liposomes and polymeric NPs) and avoid their

drawbacks;¹⁵⁴ enhanced dissolution rate, bioavailability, tissue distribution, encapsulation rate, absorption, stability of drug in body fluids, no unpleasant taste (exists with oral preparations), lower toxicity, no organic solvents usage, large-scale production possibility, sterility, reduced first-pass effect and controlled and sustained tissue-targeted delivery with various routes of administration: oral, parenteral, nasal, rectal, ophthalmic, etc.¹⁵⁵

Due to some drawbacks of SLNs such as low drug loading capacity, unpredictable gelation tendency, and drug expulsion after polymorphic transition during storage,¹⁵²⁻¹⁵⁶ NLCs were introduced and synthesized as the improved version of lipid-based nanocarriers with numerous methods exploited to formulate and prepare them.¹⁵⁵⁻¹⁵⁷ Hence, the final nanoparticle characterization must comply with the dynamic processes and such a characterization is the real challenge to represent the highest quality for the product,¹⁵⁷ storage drug expulsion, unpredictable gelation and their co-encapsulation with nano- and micro-sized structures owing to high lipid concentration and surfactants might influence the in-vivo fate of nanoparticles.

The major criteria for the characterization of nanoparticles as efficient and safe DDSs could be addressed as: size, encapsulation efficiency, structure, co-existence of material, surface morphology and functionalization, and minimum drug dose, which arguably influence directly the bioavailability, absorption and distribution of the encapsulated drug (Table 4).

Types of solid lipid nanoparticles

Type 1: Drug molecules/APIs are dispersed either in the lipid core or as amorphous clusters in “the homogenous matrix model”, which offers controlled release features. In order to design and fabricate this type, appropriate concentration of API/lipid ratio must be adjusted using either above-melting point of the lipid or cold methods of high-pressure homogenization (HPH).¹⁶²

Type 2: This type, known as “Drug enriched shell model”, is designed and fabricated with low concentration of API in the melted lipid. Using the hot method of HPH, the lipid phase is precipitated during the cooling phase which leaves a higher concentration of API in the residue of melted lipid leading to the formation of a free-API lipid core being surrounded by an outer shell composed of the saturated API and lipid. This type does not function for sustained release, however, it does for burst release of

API.¹⁶²

Type 3: Hence “Drug enriched core model”, it is designed by solubilizing the drug in the melted lipid up to its saturation solubility. Following cooling of the lipid the drug is super-saturated in the melted lipid and recrystallizes before the lipid does. Further cooling process renders also lipid recrystallization surrounding the prior formed drug-enriched core. This type offers sustained and prolonged drug release.¹⁶²

Types of nanostructured lipid carriers

Imperfect. They are named “imperfect” due to the tiny pores in the solid matrix core which are loaded with API. They are designed by adding and blending solid and liquid lipids (oil) which the co-presence of fatty acids with different chain length (mono-, di- and triacylglycerols) confers an imperfect structure for encapsulating the API.¹⁶³

Amorphous. They are designed blending lipids which don't crystallize after homogenization and cooling process,¹⁶⁴ such as hydroxyl octacosanyl hydroxy stearate, isopropyl myristate and dibutyl adipate, that give them an unorganized amorphous matrix which reduces API repelling of storage and shelf time.¹⁶³

Multiple. The advantage of the higher solubility of lipophilic drugs in liquid lipids than solid lipids can be used for formulating multiple types of NLC. Solid lipids are mixed with oils which are gradually added in higher amounts exceeding their solubility which results in phase separation of tiny particles with the surrounding solid lipid matrix. This type offers controlled drug release without expelling it out of the lipid matrix.¹⁶³

Nevertheless, even the lipid-based DDSs have their own advantages and disadvantages (Table 5).

Mechanisms of drug release by SLNs/NLCs

To improve the benefits and avoid the drawbacks, different criteria have been considered for formulation, design and encapsulation rate of nanomedicines.¹⁸⁵ The encapsulated drug undergoes surface dissolution and degradation of the lipid matrix which results in diffusion of molecules from the matrix into the surrounding tissue.¹⁵⁸ Drug release from SLNs/NLCs is affected by the localization of the drug,¹⁸⁶ which can be loaded both in the core matrix and on the surface, the former results in prolonged and sustain release while the latter burst release (quick early-phase release), subsequently conferring a biphasic release profile

Table 4. Models of drug incorporation for the lipid nanoparticles

Model	Drug Loading Site	Drug Release Pattern
Homogenous matrix of solid solution ^{158,159}	Homogeneous drug dispersion in the lipid matrix of the particles	Diffusion from the solid lipid matrix and/or by degradation of lipid matrix in GI
Drug-enriched shell ^{158,159}	Drug concentration on the outer shell of the nanoparticles	Burst release ¹⁶⁰ modified by varying the formulation conditions: production temperature (preferably cold homogenization) and surfactant concentration ¹⁶¹
Drug-enriched core ^{158,159}	Drug concentration in the core of the nanoparticles	Prolonged drug release ¹⁶¹

Table 5. Advantages and disadvantages of lipid-based drug delivery systems

Characteristics	Biological/Technological aspects	Perspectives	Reference
	Various administration routes	Broad-spectrum drug application, therapy optimization	165
	Biodegradability	Sustained drug release	166
	Controlled drug release	Patients safety, prolonged drug release, in-site drug concentration	167
	Site-specific targeting	Decreased systemic toxicity, targeted therapy	168
	Biocompatibility	No allergic reactions	169
	Increased bioavailability of encapsulated drug	Decreased dose	170
	Decreased adverse effects of toxic drugs	Improved patient safety	171
	Biological barriers penetration	Various administration routes	172
	Reduced dosing frequency	Patients compliance	167
Advantages	Chemical and enzymatic degradation drug protection	Possibility of broad-spectrum administration routes	168
	Physical stability	Improved drug formulation stability	173
	Capacity to encapsulate hydrophilic and hydrophobic drugs	Versatility for different drug groups	174
	Scaled up production	Industrial production possibility	165
	Simple manufacturing	Easy fabrication in labs, low cost	166
	No organic solvents	No toxicity concerns, green chemistry	169
	Co-delivery	Offering combined therapy	166
	Increased drug loading capacity	Decreased formulation dose	173
	Sterilization possibility	Parenteral administration optimization	175
	Small size distribution	Potential alternative for drug delivery	176
	Initial burst effect of encapsulated drug	Patient overdose risk	177
	Low plasma circulation time	Fast reticuloendothelial clearance before in-site deposition	178
	Drug expulsion during storage	Storage and administration stability challenges, industrial-scale limitation	179
Disadvantages	Low drug loading capacity	High requirement of formulation doses	180
	Polydispersity	Undesirable for intravenous administration	181
	Agglomeration	Storage issues	182
	Storage in refrigerated conditions	Transportation issues, expensive storage costs	183
	High operative temperature	Susceptibility of thermolabile drugs	184

starting with an quick release owing to the surface-loaded drug continuing by sustained release of more loaded-drug from the lipid matrix.

The burst release is a feature determined by modifying drug aqueous solubility which is influenced mainly by the surfactant concentration and the temperature via a direct proportion; the higher the last two the higher the burst release. Preparation of SLN/NLC nanoparticles at room temperature has demonstrated no burst release owing to no drug partitioning into water phase and following re-partitioning into lipid phase. Therefore, to decrease the burst release SLNs/NLCs have been prepared without surfactants or surfactants not solubilizing the drug.^{185,187}

***In vivo* fate of SLNs/NLCs after administration**

There are different criteria which determine the *in vivo* fate of nanoparticles: administration route, biological interactions with their environment including distribution, surface adsorption, nanoparticle disaggregation and enzymatic degradation.

Due to the presence of lipids and waxes in the structure

of lipid-based nanoparticles, their fate is highly affected by different pathways and enzymes for their biological interactions, namely as lipases which exist ubiquitously in body and mostly activate by oil/water interface,¹⁸⁸⁻¹⁹⁰ and actively confer various degradation rates to nanoparticles due to their formulation material¹⁹¹⁻¹⁹⁴ the free fatty acids of degradation have been studied by enzymatic test,¹⁹⁵ which demonstrated lesser degradation with long-chain fatty acids of triglycerides and surfactants contained in nanoparticles. Surfactants (e.g. poloxamer 407, poloxamer 188) function either to fasten or postpone the degradation process of nanoparticles, as different surfactants (e.g. cetyl palmitate)¹⁹² have different chain lengths, and this feature could be used in nanoparticle preparation to render a more controlled drug release profile.

So far, there has been few studies,¹⁹⁶ approving whether the presence of food in stomach would affect nanoparticles' *in-vivo* function or not, and this still remains a dilemma to be solved. In one animal study increased bioavailability and blood half-time was reported with oral administration of lipid nanodispersions,¹⁹⁷ and in another study the

increased absorption of nanoparticles into lymph through intraduodenal route was reported.¹⁹⁸

In vivo toxicity evaluation of SLNs/NLCs

Along with their different therapeutic application as DDSs, SLNs and NLCs have been investigated for their in-vivo toxicity/safety profile. Due to their composition of lipids which are physiological components they are generally recognized as safe (GRAS) and better-tolerated nanoparticles than polymeric ones showing lower toxicity,^{199,200} as they are degraded by normal physiological pathways. Nevertheless, the type of lipids and surfactants (emulsifiers) used for their preparation might increase or decrease cell toxicity and even influence encapsulated drug toxicity.^{201,202} Therefore, the toxicity evaluation must include bulk materials, SLNs/NLCs, drug/API itself and drug-encapsulated nanoparticles to analyze thoroughly each component and materials contribution to toxicity. The excipients role for drug encapsulation must be assessed according to the route of administration.²⁰³

Different in-vitro tests, among all cell viability (MTT assay) and oxidative stress, have been exploited to assess the cytotoxicity of nanoparticles; the former functions as the color-change of tetrazolium which is the indication of cell death, and the latter demonstrates DNA damages, elevated amounts of reactive oxygen species, lipid peroxidases and alterations in oxidation/reduction glutathione reactions.²⁰⁴ MTT assay is the most common method used with different dyes such as Neutral red, Trypan blue,²⁰⁴⁻²⁰⁶ and there are in-vitro and in-vivo experiments on cells proving their low toxicity.^{207,208}

SLNs and NLCs co-delivery strategies

Through numerous cancer-related studies, it has been proved that broad-spectrum anti-cancer agents will have many benefits in terms of efficacy over monotherapy. SLNs and NLCs could be a promising carrier for co-delivery of anti-cancer, therapeutic nucleic acids and antibiotics,²⁰⁹ as they are significantly able to enhance the in-vitro and in-vivo therapeutic efficacy of such drugs. Besides such advantage, co-encapsulation of different drugs in one LPDDS might decrease toxicity of the respective anti-cancer drugs and other adverse effects coming with them separately.²¹⁰

Another alternative being offered by co-delivery is RNA interference,²¹¹ especially siRNAs have been exploited for cancer cells to silence the oncogenes expression.²⁰⁹ In this context, miRNAs have been investigated and proved to be efficient to regulate genes associated with tumorigenesis.²¹⁰

In one study,²⁰⁹ cationic SLNs for co-delivery of paclitaxel and human myeloid cell leukemia (MCL1) specific siRNA have been investigated and the final result demonstrated enhanced in-vitro and in-vivo efficacy than administering them separately.

In another similar study,²¹⁰ SLNs were encapsulated for co-delivery of the same active substance with miRNA-34a.

The final result was significant in terms of eliminating lung cancer relapse mostly owing to the synergic efficacy and higher inhibition of specific receptors.

In another study,²¹² the efficacy of Paclitaxel and Verapamil co-loaded SLNs toward breast cancer were investigated to prove the efficiency of verapamil for inhibiting drug efflux transporters (e.g. p-glycoprotein) on multidrug resistance cancer cells. The study demonstrated higher expression downregulation of g-glycoproteins in the specific cancer cells, as well as higher cellular drug uptake and toxicity comparing to Paclitaxel and sole anti-cancer administration.

In another study on antibiotics,²¹³ increased antibacterial activity of Vancomycin was demonstrated. Ion pairing with linoleic acid was exploited for co-delivery and significant effects against *Staphylococcus aureus* infections which could be interpreted owing to the increased lipophilicity, sustained release of antibiotic and synergistic effect. So far different pharmaceutical/biotechnological products have been marketed using lipid-based DDSs (Table 6).

LBDDSs formulations to enhance oral delivery of hydrophobic peptide and protein therapeutics

Although there have been promising achievements with LBDDSs for oral delivery of hydrophobic peptide and protein therapeutics, the hydrophilic peptide and protein delivery still remains a challenge and limited to the in-vitro and in-vivo experiments with no product in the pharmaceutical market.

There have been many studies using these lipid-based scaffolds to prove their potential to be exploited in future studies. there have been numerous studies of Insulin as a hydrophilic peptide encapsulated in micelles, microemulsion NPs and nanocapsules with in-situ and in-vivo experiments on rat with promising results as enhanced permeability, bioavailability and efficacy.²¹⁴⁻²²⁰ In one study SK&F 106760 (a hydrophilic RGD peptide) in the form of microemulsion exploited in in-vivo experiments and demonstrated 50-fold elevated bioavailability.²²¹ In another study Vasopressin was encapsulated as microemulsion in in-situ experiments and resulted in enhanced bioavailability.²²² In one study EGF (a single-chain polypeptide) was encapsulated in microemulsions for in-vivo experiments of gastric ulcer in rats and showed increase efficacy.²²³ In one study on β -lactamase in-vivo experiments resulted in enhanced 2.5-fold bioavailability.²²⁴ N-acetylglucosaminyl and N-acetylmuramyl dipeptide were exploited in one study which demonstrated 10-fold increased bioavailability.²²⁵ In another study Leuprolide acetate was encapsulated in microemulsion for in-vivo experiments and proved increased efficacy.²²⁶ Two experiments of lipid mixtures with Hexarelin and DMP 728 (Cyclic peptide fibrinogen antagonist) as encapsulated drugs with in-situ experiments showed 20-fold and 3-fold intestinal permeability and bioavailability, respectively.^{227,228} In the latter study in dog,

Table 6. List of marketed lipid-based oral pharmaceutical products

Product/Trade name	Drug/Molecule	Nanotechnology/Dosage form	Therapeutic use/Indication	Company/Alliance
Abelcet	Amphotericin B	Nanoliposome/solution	Fungal infections	The Liposome Company Inc
Accutane	Isotretinoin	Emulsion/soft gelatine capsule	Anti-comedogenic	Roche
Agenerases	Amprenavir	Soft gelatine capsule	HIV antiviral	GlaxoSmithKline
ALEC	Dry protein free powder of DPPC-PG	Liposome	Lung diseases in infants	Britannia Pharmaceuticals Ltd
Ambisome	Amphotericin B	Powder	Fungal infections	NeXStar Pharmaceutical Inc
Amphocil	Amphotericin B	Colloidal dispersion	Fungal infections	Sequus Pharmaceutical Inc
Amphotec	Amphotericin B	Nanoliposome/Solution	Fungal infections, leishmaniasis	Sequus Pharmaceutical Inc
Aptivus	tipranavir	Emulsion/soft gelatine capsule	AIDS	Boehringer Ingelheim
Atragen	Tretinoin	Liposome	Acute myeloid leukemia	Aronex Pharmaceuticals Inc
Avodart	Dutasteride	Emulsion	Benign Prostatic Hyperplasia	GSK
Avian retrovirus vaccine	Killed avian retrovirus	Suspension	Chickenpox	Vineland Laboratories
Cipro	Ciprofloxacin	Oral suspension	Antibiotic	Bayer
Convulex	Valproic acid	Soft gelatine capsule	Antiepileptic	Pharmacia
DaunoXome	Daunorubicin citrate	Solution	Kaposi sarcoma in AIDS	NeXStar Pharmaceutical Inc/ Galen Ltd
Depakene	Valproic acid	Emulsion	Epilepsy	Abbott
Depocyt	Cytarabin	Nanoliposome/Solution	Lymphomatous meningitis	Pacira Pharmaceuticals Inc
DepoDur	Morphine	Suspension	Post-surgical pain reliever	Pacira Pharmaceuticals Inc
Doxil	Doxorubicin	Solution	Metastatic ovarian, Kaposi sarcoma in AIDS	Sequus Pharmaceutical Inc
Emend	Aprepitant	Nanosuspensions/Capsule	Antiemetic	Merck-Elan Drug Delivery
Epaxal Berna Vaccine	Inactivated hepatitis-A virions	Suspension	Hepatitis A	Swiss serum & vaccine institute
Estrasorb	estradiol	Topical emulsion	Menopausal therapy	Novavax
Evacet	Doxorubicin	Liposome	Metastatic breast cancer	The liposome company
Fenogal	Fenofibrate	Tablet	Anti hyperliproteinemic	Genus
Fortovase	saquinavir	Spontaneously emulsifying systems/ soft gelatine capsule	HIV antiviral	Roche
Fungizone	Amphotericin B	Solution	Fungal infections	Bristol-Myers Squibb
Gengraf	Cyclosporin A/III	Spontaneously emulsifying systems/ hard gelatine capsule	Immuno-suppressant	Abott
Hectoral	Doxercalciferol	Emulsion	Calcium regulator	Bone care
Juvela	Tocopherol nicotinate	Capsule	Hypertension, hyperlipidemic	Eisai Co.
Kaletra	Lopinavir & Ritonavir	Emulsion/oral solution	HIV antiviral	Abott
Lamprene	Clofazamine	Emulsion	Leprosy	Alliance laboratories/ Geigy
Lipirex	fenofibrate	Hard gelatine capsule	hyperlipidemia or mixed dyslipidemia	Sanofi-Aventis
Marinol	Dronabionol	Emulsion	Anoxeria	Roxane
Megace ES	Megestrol acetate	Nanosuspension	anorexia, cachexia, weight loss in HIV patients	Par Pharmaceuticals- Elan Drug Delivery
MiKasome	Amikacin	Liposome	Bacterial infection	NeXStar Pharmaceutical Inc
Neoral	Cyclosporin A/I	Emulsion	Immunosuppressant	Novartis
Norvir	Ritonavir	Spontaneously emulsifying systems/ soft gelatine capsule	HIV antiviral	Abott
Nyotran	Nystatin	Liposome	Fungal infections	Aronex pharmaceuticals Inc
Panzem NCD	2-Methoxy estradiol	Nanosuspension	anti-proliferative and anti-angiogenic effect	EntreMed Inc.
Prometrium	Progesterone	Emulsion	Endometrial hyperplasia	Solvay
Rapamune	Sirolimus	Nanosuspensions/Tablet	Immunosuppressant	Wyeth Pharmaceuticals – Elan Drug Delivery

Table 6. Continued.

Product/Trade name	Drug/Molecule	Nanotechnology/Dosage form	Therapeutic use/Indication	Company/Alliance
Restandol	Testosterone undecanoate	Capsules	Hormone replacement therapy	Organon laboratories
Rocaltrol	Calcitriol	Emulsion/soft gelatine capsule	Calcium regulator	Roche
Sandimmune Neoral	cyclosporine A/I	Spontaneously emulsifying systems/ soft gelatine capsule	Immunosuppressant	Novartis
Sustiva	Efavirenz	Capsules	HIV antiviral	Bristol-Meyers
Targetin	bexarotene	Soft gelatine capsule	liver cancer	Novartis
Topex-Br	Terbutalinesulphate	Syrup	Asthma	Ozone Pharmaceuticals Ltd
Tricor	Fenofibrate	Nanosuspensions/Tablet	Antihyperlipidemic agent	Abbott Laboratories
Triglide	Fenofibrate	Nanosuspensions/Tablet	Antihyperlipidemic agent	Skye Pharma-First Horizon
Ventus	Prostaglandin-E1	Liposome	Systemic inflammatory disease	The Liposome Company
Vesanoid	tretinoine	Emulsion/soft gelatine capsule	Acne	Roche
VincaXome	Vincristine	Liposome	Solid tumors	NeXstar Pharmaceutical Inc
Zemplar	Paricalcitol	Emulsion	Calcium regulator	Abbott

DuP 532 (an Angiotensin II antagonist) was encapsulated in microemulsion in in-vivo experiments which resulted in 3-fold bioavailability.²²⁸ In three studies in rat and pig, calcitonin was encapsulated in mixed micelles and emulsion for in-situ and in-vivo experiments which demonstrated increased permeability and efficacy and 4-fold hypocalcemia response.²²⁹⁻²³¹ Human growth hormone was encapsulated in in-vivo studies on rabbit and showed 3.3% increased bioavailability.²³²

Regulatory status, commercialization plan and safety information

The status of excipients should be assessed with the regulatory authorities before any pharmaceutical product's introduction into the market²³³ but the expenses of in-vivo toxicity studies are prohibitive for the companies. Such a challenge is happening mainly with the polymeric NPs as there are few of them in the market but lipid NPs owing to their various applications of oils, fats, stabilizer and surfactants have introduced oral and dermal products. The majority of the introduced excipients so far for lipid NPs synthesis are biodegradable, biocompatible and are approved as safe, but some are toxic at high concentrations.²³⁴ In this context the FDA has published guide lists of safe materials and substances (GRAS) and inactive ingredient guide (IIG) for excipients that are approved for exploiting in the pharmaceutical products in the market.²³⁵ These lists explain and provide insights regarding the appropriate excipient concentration for each administration route and the approved inactive ingredients used for a specific route can be used in all the new formulations. This facilitates the process of synthesizing new formulation as the necessary information can be extracted from the GRAS and IIG. Therefore, such excipients are assessed as substances of a drug not individually as from a "scientific point of view" excipients are a major part of the drug formulation. Nevertheless, there are other challenges to consider from a "regulatory point of view"; preclinical and clinical studies

addressing safety issues and in-vivo manifestations of the LBDDSs in terms of clinical therapeutic efficacy. In-vivo immunological and stability findings toward oils and lipid excipients must be reported to provide in-depth information for the regulatory authorities.²³⁶

Besides, factors coming from the biopharmaceuticals are required to be evaluated toward the drug or excipients and this might have paradoxical in-vitro results with in-vivo results due to the physiology of GI tract. Various experiments must be designed and conducted to characterize and recognize the interactions happening among excipients, in-vivo physiological conditions and the drug.²³⁷

In order to understand and characterize the in-vivo fate of drugs encapsulated in LBDDSs, a consortium of academic and industrial scientists has been established (<http://www.lfcsconsortium.org>) which designs experiments to evaluate the function of LBDDSs dispersion and digestion as vital criteria.

Conclusion

Nanotechnology offers promising strategies for enhancing oral bioavailability and therapeutic efficacy of a vast range of drugs; conventional chemical drugs with poor water solubility and biotechnological, peptide and protein therapeutics and biopharmaceuticals. The unique physicochemical features of peptide and protein therapeutics pose challenge for their oral delivery. Hence, their success in site delivery highly depends on technologies and methods to modify these two features not influencing their biological function. In the recent decades numerous DDSs have been introduced and offered by nanotechnology to achieve as high successful delivery as possible and LBDDSs among all has been under investigation owing to their potential for oral delivery of hydrophilic, hydrophobic and lipophilic peptide and protein therapeutics.

LBDDSs enhance solubility and bioavailability of drugs offering strategies such as gastrointestinal lymphatic

transport, altering physiological and biochemical properties of gastrointestinal barriers, elevated solubilization and prolonged gastrointestinal retention. Although, such improvements rely on the encapsulation/loading rate and intrinsic composition of the material used during the fabrication process. Obviously, the choice of materials, such as excipients, will influence the success of delivery route which is determined both by lipid formulation design and peptide/protein molecule emphasizing that each peptide/protein-loaded LBDDS must be designed uniquely. Such material must be in correlation with the drug of choice to achieve the maximum therapeutic efficacy and in-site dose.

Most of the scaffolds described in this review article suggest promising alternatives to overcome gastrointestinal enzymatic degradation and poor membrane penetration. Further systematic studies are required to evaluate their in-vivo efficacy in terms of peptide-/protein-based oral drug delivery. Besides “pharmaco-biotechnological” challenges mentioned in this review such as membrane permeability, protease stability, delivery strategies and increased circulation half-life, there are inevitably several “industrial” challenges as well which finally hamper their industrial scale production and consequently their biomedical translation from lab to pharmaceutical market. “Oral bioavailability” still remains the main challenge of peptide/protein-based drug delivery. These factors could be addressed as materials cost, drug potential market feedback, regulatory status, simple industrial-scale fabrication, financial schemes for required instruments, patient compliance administration and high adaptability to human diverse pharmacokinetics.

Competing Interests

The author declares no conflict of interest.

Ethical Approval

There is none to be disclosed.

Funding

The author declares no funding/support.

References

- Vlieghe P, Lisowski V, Martinez J, Khrestchatskiy M. Synthetic therapeutic peptides: science and market. *Drug Discov Today* 2010;15(1-2):40-56. doi: [10.1016/j.drudis.2009.10.009](https://doi.org/10.1016/j.drudis.2009.10.009)
- Lalatsa A, Schatzlein AG, Uchegbu IF. Strategies to deliver peptide drugs to the brain. *Mol Pharm* 2014;11(4):1081-93. doi: [10.1021/mp400680d](https://doi.org/10.1021/mp400680d)
- Vajo Z, Fawcett J, Duckworth WC. Recombinant DNA technology in the treatment of diabetes: insulin analogs. *Endocr Rev* 2001;22(5):706-17. doi: [10.1210/edrv.22.5.0442](https://doi.org/10.1210/edrv.22.5.0442)
- Takeda A, Cooper K, Bird A, Baxter L, Frampton GK, Gospodarevskaya E, et al. Recombinant human growth hormone for the treatment of growth disorders in children: a systematic review and economic evaluation. *Health Technol Assess* 2010;14(42):1-209. doi: [10.3310/hta14420](https://doi.org/10.3310/hta14420)
- Cutting GR. Modifier genetics: cystic fibrosis. *Annu Rev Genomics Hum Genet* 2005;6:237-60. doi: [10.1146/annurev.genom.6.080604.162254](https://doi.org/10.1146/annurev.genom.6.080604.162254)
- Weatherall DJ. Phenotype-genotype relationships in monogenic disease: lessons from the thalassaemias. *Nat Rev Genet* 2001;2(4):245-55. doi: [10.1038/35066048](https://doi.org/10.1038/35066048)
- Powell JS. Lasting power of new clotting proteins. *Hematology Am Soc Hematol Educ Program* 2014;2014(1):355-63. doi: [10.1182/asheducation-2014.1.355](https://doi.org/10.1182/asheducation-2014.1.355)
- Hirschhorn JN, Lohmueller K, Byrne E, Hirschhorn K. A comprehensive review of genetic association studies. *Genet Med* 2002;4(2):45-61. doi: [10.1097/00125817-200203000-00002](https://doi.org/10.1097/00125817-200203000-00002)
- Savic S, McDermott MF. Clinical genetics in 2014: new monogenic diseases span the immunological disease continuum. *Nat Rev Rheumatol* 2015;11(2):67-8. doi: [10.1038/nrrheum.2014.215](https://doi.org/10.1038/nrrheum.2014.215)
- Hussain N, Jaitley V, Florence AT. Recent advances in the understanding of uptake of microparticulates across the gastrointestinal lymphatics. *Adv Drug Deliv Rev* 2001;50(1-2):107-42. doi: [10.1016/s0169-409x\(01\)00152-1](https://doi.org/10.1016/s0169-409x(01)00152-1)
- Banga AK. Drug delivery today, Pharma Tech 2002, Business Briefing World Market Series, London, 2002, pp. 150–154.
- Wang W. Instability, stabilization, and formulation of liquid protein pharmaceuticals. *Int J Pharm* 1999;185(2):129-88. doi: [10.1016/s0378-5173\(99\)00152-0](https://doi.org/10.1016/s0378-5173(99)00152-0)
- Hillery AM, Lloyd AW, Swarbrick J. *Drug Delivery and Targeting: For Pharmacists and Pharmaceutical Scientists*. 1st ed. London: CRC Press; 2001. doi: [10.1201/b12801](https://doi.org/10.1201/b12801)
- Frokjaer S, Otzen DE. Protein drug stability: a formulation challenge. *Nat Rev Drug Discov* 2005;4(4):298-306. doi: [10.1038/nrd1695](https://doi.org/10.1038/nrd1695)
- Wang W. Protein aggregation and its inhibition in biopharmaceutics. *Int J Pharm* 2005;289(1-2):1-30. doi: [10.1016/j.ijpharm.2004.11.014](https://doi.org/10.1016/j.ijpharm.2004.11.014)
- Rubert Pérez CM, Stephanopoulos N, Sur S, Lee SS, Newcomb C, Stupp SI. The powerful functions of peptide-based bioactive matrices for regenerative medicine. *Ann Biomed Eng* 2015;43(3):501-14. doi: [10.1007/s10439-014-1166-6](https://doi.org/10.1007/s10439-014-1166-6)
- Lax R. The future of peptide development in the pharmaceutical industry. In: *PharManufacturing: The International Peptide Review*. London, UK: World Business Journals, Pharmaceutical Division; 2010.
- Serrano Lopez DR, Lalatsa A. Peptide pills for brain diseases? Reality and future perspectives. *Ther Deliv* 2013;4(4):479-501. doi: [10.4155/tde.13.5](https://doi.org/10.4155/tde.13.5)
- Adessi C, Soto C. Converting a peptide into a drug: strategies to improve stability and bioavailability. *Curr Med Chem* 2002;9(9):963-78. doi: [10.2174/0929867024606731](https://doi.org/10.2174/0929867024606731)
- Adessi C, Soto C. Strategies to improve stability and bioavailability of peptide drugs. *Front Med Chem* 2004;1(1):513-28. doi: [10.2174/1567204043396622](https://doi.org/10.2174/1567204043396622)
- Park K, Kwon IC, Park K. Oral protein delivery: current status and future prospect. *React Funct Polym* 2011;71(3):280-7. doi: [10.1016/j.reactfunctpolym.2010.10.002](https://doi.org/10.1016/j.reactfunctpolym.2010.10.002)
- Donovan MD, Flynn GL, Amidon GL. Absorption of polyethylene glycols 600 through 2000: the molecular weight dependence of gastrointestinal and nasal absorption. *Pharm Res* 1990;7(8):863-8. doi: [10.1023/a:1015921101465](https://doi.org/10.1023/a:1015921101465)
- Ikesue K, Kopečková P, Kopeček J. Degradation of proteins by guinea pig intestinal enzymes. *Int J Pharm* 1993;95(1-3):171-9. doi: [10.1016/0378-5173\(93\)90404-4](https://doi.org/10.1016/0378-5173(93)90404-4)
- Saffran M, Kumar GS, Savariar C, Burnham JC, Williams F, Neckers DC. A new approach to the oral administration of insulin and other peptide drugs. *Science* 1986;233(4768):1081-4. doi: [10.1126/science.3526553](https://doi.org/10.1126/science.3526553)
- Fix JA. Oral controlled release technology for peptides: status and future prospects. *Pharm Res* 1996;13(12):1760-4. doi: [10.1023/a:1016008419367](https://doi.org/10.1023/a:1016008419367)
- Lee HJ. Protein drug oral delivery: the recent progress. *Arch Pharm Res* 2002;25(5):572-84. doi: [10.1007/bf02976925](https://doi.org/10.1007/bf02976925)

27. Lee VH, Satish DK, George MG, Werner R. Oral route of protein and peptide drug delivery. In: Lee VH, ed. *Peptide and Protein Drug Delivery*. New York: Marcel Dekker; 1991. p. 691-738.
28. Pettit DK, Gombotz WR. The development of site-specific drug-delivery systems for protein and peptide biopharmaceuticals. *Trends Biotechnol* 1998;16(8):343-9. doi: [10.1016/s0167-7799\(98\)01186-x](https://doi.org/10.1016/s0167-7799(98)01186-x)
29. Tauzin B. *Report: Biotechnology Medicines in Development*. Washington, DC: Pharmaceutical Research and Manufacturers Association; 2006.
30. Crommelin D, van Winden E, Mekking A. Delivery of pharmaceutical proteins. In: Aulton ME, ed. *Pharmaceutics: The Science of Dosage form Design*. Edinburgh: Churchill Livingstone; 2001. p. 544-53.
31. Florence AT, Attwood D. *Physicochemical Principles of Pharmacy*. London: Pharmaceutical Press; 2006.
32. Cleland JL, Langer R. Formulation and delivery of proteins and peptides: design and development strategies. In: Cleland JL, Langer R, eds. *Formulation and Delivery of Proteins and Peptides*. Washington, DC: American Chemical Society; 1994. p. 1-19.
33. Almeida AJ, Souto E. Solid lipid nanoparticles as a drug delivery system for peptides and proteins. *Adv Drug Deliv Rev* 2007;59(6):478-90. doi: [10.1016/j.addr.2007.04.007](https://doi.org/10.1016/j.addr.2007.04.007)
34. Wang W. Instability, stabilization, and formulation of liquid protein pharmaceuticals. *Int J Pharm* 1999;185(2):129-88. doi: [10.1016/s0378-5173\(99\)00152-0](https://doi.org/10.1016/s0378-5173(99)00152-0)
35. Metselaar JM, Mastrobattista E, Storm G. Liposomes for intravenous drug targeting: design and applications. *Mini Rev Med Chem* 2002;2(4):319-29. doi: [10.2174/1389557023405873](https://doi.org/10.2174/1389557023405873)
36. Gombotz WR, Pettit DK. Biodegradable polymers for protein and peptide drug delivery. *Bioconjug Chem* 1995;6(4):332-51. doi: [10.1021/bc00034a002](https://doi.org/10.1021/bc00034a002)
37. Packhaeuser CB, Schnieders J, Oster CG, Kissel T. In situ forming parenteral drug delivery systems: an overview. *Eur J Pharm Biopharm* 2004;58(2):445-55. doi: [10.1016/j.ejpb.2004.03.003](https://doi.org/10.1016/j.ejpb.2004.03.003)
38. Kompella UB, Lee VH. Delivery systems for penetration enhancement of peptide and protein drugs: design considerations. *Adv Drug Deliv Rev* 2001;46(1-3):211-45. doi: [10.1016/s0169-409x\(00\)00137-x](https://doi.org/10.1016/s0169-409x(00)00137-x)
39. Prego C, Torres D, Alonso MJ. The potential of chitosan for the oral administration of peptides. *Expert Opin Drug Deliv* 2005;2(5):843-54. doi: [10.1517/17425247.2.5.843](https://doi.org/10.1517/17425247.2.5.843)
40. Ugwoke MI, Agu RU, Verbeke N, Kinget R. Nasal mucoadhesive drug delivery: background, applications, trends and future perspectives. *Adv Drug Deliv Rev* 2005;57(11):1640-65. doi: [10.1016/j.addr.2005.07.009](https://doi.org/10.1016/j.addr.2005.07.009)
41. Alpar HO, Somavrapu S, Atuah KN, Bramwell VW. Biodegradable mucoadhesive particulates for nasal and pulmonary antigen and DNA delivery. *Adv Drug Deliv Rev* 2005;57(3):411-30. doi: [10.1016/j.addr.2004.09.004](https://doi.org/10.1016/j.addr.2004.09.004)
42. Schuetz YB, Naik A, Guy RH, Kalia YN. Emerging strategies for the transdermal delivery of peptide and protein drugs. *Expert Opin Drug Deliv* 2005;2(3):533-48. doi: [10.1517/17425247.2.3.533](https://doi.org/10.1517/17425247.2.3.533)
43. Langer R. Where a pill won't reach. *Sci Am* 2003;288(4):50-7. doi: [10.1038/scientificamerican0403-50](https://doi.org/10.1038/scientificamerican0403-50)
44. Myles ME, Neumann DM, Hill JM. Recent progress in ocular drug delivery for posterior segment disease: emphasis on transscleral iontophoresis. *Adv Drug Deliv Rev* 2005;57(14):2063-79. doi: [10.1016/j.addr.2005.08.006](https://doi.org/10.1016/j.addr.2005.08.006)
45. Smart JD. Buccal drug delivery. *Expert Opin Drug Deliv* 2005;2(3):507-17. doi: [10.1517/17425247.2.3.507](https://doi.org/10.1517/17425247.2.3.507)
46. Crommelin DJ, Storm G, Verrijck R, de Leede L, Jiskoot W, Hennink WE. Shifting paradigms: biopharmaceuticals versus low molecular weight drugs. *Int J Pharm* 2003;266(1-2):3-16. doi: [10.1016/s0378-5173\(03\)00376-4](https://doi.org/10.1016/s0378-5173(03)00376-4)
47. Saltzman WM. *Drug Delivery: Engineering Principles for Drug Therapy*. New York: Oxford University Press; 2001. doi: [10.1093/oso/9780195085891.003.0005](https://doi.org/10.1093/oso/9780195085891.003.0005)
48. Ugwoke MI, Agu RU, Verbeke N, Kinget R. Nasal mucoadhesive drug delivery: background, applications, trends and future perspectives. *Adv Drug Deliv Rev* 2005;57(11):1640-65. doi: [10.1016/j.addr.2005.07.009](https://doi.org/10.1016/j.addr.2005.07.009)
49. Myles ME, Neumann DM, Hill JM. Recent progress in ocular drug delivery for posterior segment disease: emphasis on transscleral iontophoresis. *Adv Drug Deliv Rev* 2005;57(14):2063-79. doi: [10.1016/j.addr.2005.08.006](https://doi.org/10.1016/j.addr.2005.08.006)
50. Smart JD. Buccal drug delivery. *Expert Opin Drug Deliv* 2005;2(3):507-17. doi: [10.1517/17425247.2.3.507](https://doi.org/10.1517/17425247.2.3.507)
51. Mackay M, Phillips J, Hastewell J. Peptide drug delivery: colonic and rectal absorption. *Adv Drug Deliv Rev* 1997;28(2):253-73. doi: [10.1016/s0169-409x\(97\)00076-8](https://doi.org/10.1016/s0169-409x(97)00076-8)
52. Hussain A, Ahsan F. The vagina as a route for systemic drug delivery. *J Control Release* 2005;103(2):301-13. doi: [10.1016/j.jconrel.2004.11.034](https://doi.org/10.1016/j.jconrel.2004.11.034)
53. Schuetz YB, Naik A, Guy RH, Kalia YN. Emerging strategies for the transdermal delivery of peptide and protein drugs. *Expert Opin Drug Deliv* 2005;2(3):533-48. doi: [10.1517/17425247.2.3.533](https://doi.org/10.1517/17425247.2.3.533)
54. Agu RU, Ugwoke MI, Armand M, Kinget R, Verbeke N. The lung as a route for systemic delivery of therapeutic proteins and peptides. *Respir Res* 2001;2(4):198-209. doi: [10.1186/rr58](https://doi.org/10.1186/rr58)
55. Bosquillon C, Pr at V, Vanbever R. Pulmonary delivery of growth hormone using dry powders and visualization of its local fate in rats. *J Control Release* 2004;96(2):233-44. doi: [10.1016/j.jconrel.2004.01.027](https://doi.org/10.1016/j.jconrel.2004.01.027)
56. Ghilzai NM, Desai A. Facing the challenges of transmucosal absorption—buccal, nasal and rectal routes. Pharma Tech 2004, Business Briefing World Market Series, London, 2004, pp. 104–106.
57. Choonara BF, Choonara YE, Kumar P, Bijukumar D, du Toit LC, Pillay V. A review of advanced oral drug delivery technologies facilitating the protection and absorption of protein and peptide molecules. *Biotechnol Adv* 2014;32(7):1269-82. doi: [10.1016/j.biotechadv.2014.07.006](https://doi.org/10.1016/j.biotechadv.2014.07.006)
58. Ensign LM, Cone R, Hanes J. Oral drug delivery with polymeric nanoparticles: the gastrointestinal mucus barriers. *Adv Drug Deliv Rev* 2012;64(6):557-70. doi: [10.1016/j.addr.2011.12.009](https://doi.org/10.1016/j.addr.2011.12.009)
59. des Rieux A, Fievez V, Garinot M, Schneider YJ, Pr at V. Nanoparticles as potential oral delivery systems of proteins and vaccines: a mechanistic approach. *J Control Release* 2006;116(1):1-27. doi: [10.1016/j.jconrel.2006.08.013](https://doi.org/10.1016/j.jconrel.2006.08.013)
60. Prego C, Torres D, Fernandez-Megia E, Novoa-Carballal R, Qui noa E, Alonso MJ. Chitosan-PEG nanocapsules as new carriers for oral peptide delivery. Effect of chitosan pegylation degree. *J Control Release* 2006;111(3):299-308. doi: [10.1016/j.jconrel.2005.12.015](https://doi.org/10.1016/j.jconrel.2005.12.015)
61. Amidon GL, Lennern s H, Shah VP, Crison JR. A theoretical basis for a biopharmaceutical drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability. *Pharm Res* 1995;12(3):413-20. doi: [10.1023/a:1016212804288](https://doi.org/10.1023/a:1016212804288)
62. Lipinski CA. Avoiding investment in doomed drugs, is poor solubility an industry wide problem? *Curr Drugs Dis* 2001;4:17-9.
63. Murakami T, Takano M. Intestinal efflux transporters and drug absorption. *Expert Opin Drug Metab Toxicol* 2008;4(7):923-39. doi: [10.1517/17425255.4.7.923](https://doi.org/10.1517/17425255.4.7.923)

64. Uekama K, Hirayama F, Irie T. Cyclodextrin drug carrier systems. *Chem Rev* 1998;98(5):2045-76. doi: [10.1021/cr970025p](https://doi.org/10.1021/cr970025p)
65. Tazuin B. *Report: Biotechnology Medicines in Development*. Washington, DC: Pharmaceutical Research and Manufacturers Association; 2019.
66. Hillery AM. Drug delivery: the basic concepts. In: Hillery AM, Lloyd AW, Swarbrick J, eds. *Drug Delivery and Targeting for Pharmacists and Pharmaceutical Scientists*. London: Taylor & Francis; 2001. p. 1-48.
67. Frokjaer S, Otzen DE. Protein drug stability: a formulation challenge. *Nat Rev Drug Discov* 2005;4(4):298-306. doi: [10.1038/nrd1695](https://doi.org/10.1038/nrd1695)
68. Zizzari AT, Pliatsika D, Gall FM, Fischer T, Riedl R. New perspectives in oral peptide delivery. *Drug Discov Today*. 2021;26(4):1097-1105. doi:[10.1016/j.drudis.2021.01.020](https://doi.org/10.1016/j.drudis.2021.01.020)
69. Scioli Montoto S, Muraca G, Ruiz ME. Solid Lipid Nanoparticles for Drug Delivery: Pharmacological and Biopharmaceutical Aspects. *Front Mol Biosci*. 2020;7:587997. doi:[10.3389/fmolb.2020.587997](https://doi.org/10.3389/fmolb.2020.587997)
70. Wang W. Instability, stabilization, and formulation of liquid protein pharmaceuticals. *Int J Pharm* 1999;185(2):129-88. doi: [10.1016/s0378-5173\(99\)00152-0](https://doi.org/10.1016/s0378-5173(99)00152-0)
71. Mehrdadi S. Acute bacterial meningitis: diagnosis, treatment and prevention. *J Arch Mil Med* 2019;6(4):e84749. doi: [10.5812/jamm.84749](https://doi.org/10.5812/jamm.84749)
72. Mehrdadi S. Drug delivery of solid lipid nanoparticles (SLNs) and nanostructured lipid carriers (NLCs) to target brain tumors. *Adv Pharm Bull* 2023;13(3):512-20. doi: [10.34172/apb.2023.062](https://doi.org/10.34172/apb.2023.062)
73. Tsomaia N. Peptide therapeutics: targeting the undruggable space. *Eur J Med Chem* 2015;94:459-70. doi: [10.1016/j.ejmech.2015.01.014](https://doi.org/10.1016/j.ejmech.2015.01.014)
74. GlobalData. 2015. Available from: <http://www.globaldata.com>.
75. Tomita M, Hayashi M, Horie T, Ishizawa T, Awazu S. Enhancement of colonic drug absorption by the transcellular permeation route. *Pharm Res* 1988;5(12):786-9. doi: [10.1023/a:1015992819290](https://doi.org/10.1023/a:1015992819290)
76. Pappenheimer JR. Physiological regulation of transepithelial impedance in the intestinal mucosa of rats and hamsters. *J Membr Biol* 1987;100(2):137-48. doi: [10.1007/bf02209146](https://doi.org/10.1007/bf02209146)
77. Tang VW, Goodenough DA. Paracellular ion channel at the tight junction. *Biophys J* 2003;84(3):1660-73. doi: [10.1016/s0006-3495\(03\)74975-3](https://doi.org/10.1016/s0006-3495(03)74975-3)
78. Barry PH. Ionic permeation mechanisms in epithelia: biionic potentials, dilution potentials, conductances, and streaming potentials. *Methods Enzymol* 1989;171:678-715. doi: [10.1016/s0076-6879\(89\)71038-7](https://doi.org/10.1016/s0076-6879(89)71038-7)
79. Powell DW. Barrier function of epithelia. *Am J Physiol* 1981;241(4):G275-88. doi: [10.1152/ajpgi.1981.241.4.G275](https://doi.org/10.1152/ajpgi.1981.241.4.G275)
80. Rubas W, Cromwell ME, Shahrokh Z, Villagran J, Nguyen TN, Wellton M, et al. Flux measurements across Caco-2 monolayers may predict transport in human large intestinal tissue. *J Pharm Sci* 1996;85(2):165-9. doi: [10.1021/js950267](https://doi.org/10.1021/js950267) +
81. Madara JL. Regulation of the movement of solutes across tight junctions. *Annu Rev Physiol* 1998;60:143-59. doi: [10.1146/annurev.physiol.60.1.143](https://doi.org/10.1146/annurev.physiol.60.1.143)
82. Shakweh M, Ponchel G, Fattal E. Particle uptake by Peyer's patches: a pathway for drug and vaccine delivery. *Expert Opin Drug Deliv* 2004;1(1):141-63. doi: [10.1517/17425247.1.1.141](https://doi.org/10.1517/17425247.1.1.141)
83. Russell-Jones GJ. Carrier-mediated transport, oral drug delivery. In: Mathiowitz E, ed. *Encyclopedia of Controlled Drug Delivery*. New York, NY: John Wiley & Sons; 1999. p. 173-84.
84. Barthe L, Woodley J, Houin G. Gastrointestinal absorption of drugs: methods and studies. *Fundam Clin Pharmacol* 1999;13(2):154-68. doi: [10.1111/j.1472-8206.1999.tb00334.x](https://doi.org/10.1111/j.1472-8206.1999.tb00334.x)
85. Bai JP, Amidon GL. Structural specificity of mucosal-cell transport and metabolism of peptide drugs: implication for oral peptide drug delivery. *Pharm Res* 1992;9(8):969-78. doi: [10.1023/a:1015885823793](https://doi.org/10.1023/a:1015885823793)
86. Russell-Jones GJ. The potential use of receptor-mediated endocytosis for oral drug delivery. *Adv Drug Deliv Rev* 1996;20(1):83-97. doi: [10.1016/0169-409x\(95\)00131-p](https://doi.org/10.1016/0169-409x(95)00131-p)
87. Swaan PW. Recent advances in intestinal macromolecular drug delivery via receptor-mediated transport pathways. *Pharm Res* 1998;15(6):826-34. doi: [10.1023/a:1011908128045](https://doi.org/10.1023/a:1011908128045)
88. Charman WN, Porter CJH. Lipophilic prodrugs designed for intestinal lymphatic transport. *Adv Drug Deliv Rev* 1996;19(2):149-69. doi: [10.1016/0169-409x\(95\)00105-g](https://doi.org/10.1016/0169-409x(95)00105-g)
89. Florence AT. Issues in oral nanoparticle drug carrier uptake and targeting. *J Drug Target* 2004;12(2):65-70. doi: [10.1080/10611860410001693706](https://doi.org/10.1080/10611860410001693706)
90. Burton PS, Conradi RA, Hilgers AR. (B) Mechanisms of peptide and protein absorption: (2) transcellular mechanism of peptide and protein absorption: passive aspects. *Adv Drug Deliv Rev* 1991;7(3):365-85. doi: [10.1016/0169-409x\(91\)90014-4](https://doi.org/10.1016/0169-409x(91)90014-4)
91. Giannasca PJ, Giannasca KT, Leichtner AM, Neutra MR. Human intestinal M cells display the sialyl Lewis A antigen. *Infect Immun* 1999;67(2):946-53. doi: [10.1128/iai.67.2.946-953.1999](https://doi.org/10.1128/iai.67.2.946-953.1999)
92. Gebert A, Rothkötter HJ, Pabst R. M cells in Peyer's patches of the intestine. In: Jeon KW, ed. *International Review of Cytology*. Academic Press; 1996. p. 91-159. doi: [10.1016/s0074-7696\(08\)61346-7](https://doi.org/10.1016/s0074-7696(08)61346-7)
93. Frey A, Neutra MR. Targeting of mucosal vaccines to Peyer's patch M cells. *Behring Inst Mitt* 1997(98):376-89.
94. Clark MA, Hirst BH, Jepson MA. Lectin-mediated mucosal delivery of drugs and microparticles. *Adv Drug Deliv Rev* 2000;43(2-3):207-23. doi: [10.1016/s0169-409x\(00\)00070-3](https://doi.org/10.1016/s0169-409x(00)00070-3)
95. Buda A, Sands C, Jepson MA. Use of fluorescence imaging to investigate the structure and function of intestinal M cells. *Adv Drug Deliv Rev* 2005;57(1):123-34. doi: [10.1016/j.addr.2004.07.014](https://doi.org/10.1016/j.addr.2004.07.014)
96. Jani PU, Florence AT, McCarthy DE. Further histological evidence of the gastrointestinal absorption of polystyrene nanospheres in the rat. *Int J Pharm* 1992;84(3):245-52. doi: [10.1016/0378-5173\(92\)90162-u](https://doi.org/10.1016/0378-5173(92)90162-u)
97. Kataoka K, Tabata J, Yamamoto M, Toyota T. The association of gap junctions with large particles in the crypt epithelium of the rat small intestine. *Arch Histol Cytol* 1989;52(2):81-6. doi: [10.1679/aohc.52.81](https://doi.org/10.1679/aohc.52.81)
98. Lavelle EC, Sharif S, Thomas NW, Holland J, Davis SS. The importance of gastrointestinal uptake of particles in the design of oral delivery systems. *Adv Drug Deliv Rev* 1995;18(1):5-22. doi: [10.1016/0169-409x\(95\)00048-c](https://doi.org/10.1016/0169-409x(95)00048-c)
99. O'Hagan DT. Intestinal translocation of particulates—implications for drug and antigen delivery. *Adv Drug Deliv Rev* 1990;5(3):265-85. doi: [10.1016/0169-409x\(90\)90020-s](https://doi.org/10.1016/0169-409x(90)90020-s)
100. Hebden JM, Wilson CG, Spiller RC, Gilchrist PJ, Blackshaw E, Frier ME, et al. Regional differences in quinine absorption from the undisturbed human colon assessed using a timed release delivery system. *Pharm Res* 1999;16(7):1087-92. doi: [10.1023/a:1018948102778](https://doi.org/10.1023/a:1018948102778)
101. Shah D, Shen WC. Transcellular delivery of an insulin-transferrin conjugate in enterocyte-like Caco-2 cells. *J Pharm Sci* 1996;85(12):1306-11. doi: [10.1021/js9601400](https://doi.org/10.1021/js9601400)
102. Chakraborty S, Shukla D, Mishra B, Singh S. Lipid--an emerging platform for oral delivery of drugs with poor bioavailability. *Eur J Pharm Biopharm* 2009;73(1):1-15. doi: [10.1016/j.ejpb.2009.06.001](https://doi.org/10.1016/j.ejpb.2009.06.001)
103. Charman WN, Porter CJ, Mithani S, Dressman JB.

- Physicochemical and physiological mechanisms for the effects of food on drug absorption: the role of lipids and pH. *J Pharm Sci* 1997;86(3):269-82. doi: [10.1021/js960085v](https://doi.org/10.1021/js960085v)
104. Crouse RG. Human pharmacology of griseofulvin: the effect of fat intake on gastrointestinal absorption. *J Invest Dermatol* 1961;37:529-33. doi: [10.1038/jid.1961.154](https://doi.org/10.1038/jid.1961.154)
 105. Hörter D, Dressman JB. Influence of physicochemical properties on dissolution of drugs in the gastrointestinal tract. *Adv Drug Deliv Rev* 2001;46(1-3):75-87. doi: [10.1016/s0169-409x\(00\)00130-7](https://doi.org/10.1016/s0169-409x(00)00130-7)
 106. Wagner D, Spahn-Langguth H, Hanafy A, Koggel A, Langguth P. Intestinal drug efflux: formulation and food effects. *Adv Drug Deliv Rev* 2001;50 Suppl 1:S13-31. doi: [10.1016/s0169-409x\(01\)00183-1](https://doi.org/10.1016/s0169-409x(01)00183-1)
 107. Touitou E, Barry BW. *Enhancement in Drug Delivery*. Florida: CRC Press; 2006.
 108. Liversidge GG, Cundy KC. Particle size reduction for improvement of oral bioavailability of hydrophobic drugs: I. Absolute oral bioavailability of nanocrystalline danazol in beagle dogs. *Int J Pharm* 1995;125(1):91-7. doi: [10.1016/0378-5173\(95\)00122-y](https://doi.org/10.1016/0378-5173(95)00122-y)
 109. Charman WN. Lipids, lipophilic drugs, and oral drug delivery—some emerging concepts. *J Pharm Sci* 2000;89(8):967-78. doi: [10.1002/1520-6017\(200008\)89:8<967::aid-jps1>3.0.co;2-r](https://doi.org/10.1002/1520-6017(200008)89:8<967::aid-jps1>3.0.co;2-r)
 110. Charman WN, Porter CJ, Mithani S, Dressman JB. Physicochemical and physiological mechanisms for the effects of food on drug absorption: the role of lipids and pH. *J Pharm Sci* 1997;86(3):269-82. doi: [10.1021/js960085v](https://doi.org/10.1021/js960085v)
 111. Porter CJ, Charman WN. Intestinal lymphatic drug transport: an update. *Adv Drug Deliv Rev* 2001;50(1-2):61-80. doi: [10.1016/s0169-409x\(01\)00151-x](https://doi.org/10.1016/s0169-409x(01)00151-x)
 112. Charman SA, Charman WN, Rogge MC, Wilson TD, Dutko FJ, Pouton CW. Self-emulsifying drug delivery systems: formulation and biopharmaceutic evaluation of an investigational lipophilic compound. *Pharm Res* 1992;9(1):87-93. doi: [10.1023/a:1018987928936](https://doi.org/10.1023/a:1018987928936)
 113. Li C, Fleisher D, Li L, Schwier JR, Sweetana SA, Vasudevan V, et al. Regional-dependent intestinal absorption and meal composition effects on systemic availability of LY303366, a lipopeptide antifungal agent, in dogs. *J Pharm Sci* 2001;90(1):47-57. doi: [10.1002/1520-6017\(200101\)90:1<47::aid-jps6>3.0.co;2-2](https://doi.org/10.1002/1520-6017(200101)90:1<47::aid-jps6>3.0.co;2-2)
 114. Martinez M, Amidon G, Clarke L, Jones WW, Mitra A, Riviere J. Applying the biopharmaceutics classification system to veterinary pharmaceutical products. Part II. Physiological considerations. *Adv Drug Deliv Rev* 2002;54(6):825-50. doi: [10.1016/s0169-409x\(02\)00071-6](https://doi.org/10.1016/s0169-409x(02)00071-6)
 115. Sanjula B, Shah FM, Javed A, Alka A. Effect of poloxamer 188 on lymphatic uptake of carvedilol-loaded solid lipid nanoparticles for bioavailability enhancement. *J Drug Target* 2009;17(3):249-56. doi: [10.1080/10611860902718672](https://doi.org/10.1080/10611860902718672)
 116. Trevaskis NL, Charman WN, Porter CJ. Lipid-based delivery systems and intestinal lymphatic drug transport: a mechanistic update. *Adv Drug Deliv Rev* 2008;60(6):702-16. doi: [10.1016/j.addr.2007.09.007](https://doi.org/10.1016/j.addr.2007.09.007)
 117. Khoo SM, Shackelford DM, Porter CJ, Edwards GA, Charman WN. Intestinal lymphatic transport of halofantrine occurs after oral administration of a unit-dose lipid-based formulation to fasted dogs. *Pharm Res* 2003;20(9):1460-5. doi: [10.1023/a:1025718513246](https://doi.org/10.1023/a:1025718513246)
 118. Wang W. Oral protein drug delivery. *J Drug Target* 1996;4(4):195-232. doi: [10.3109/10611869608995624](https://doi.org/10.3109/10611869608995624)
 119. Patel G, Misra A. Oral delivery of proteins and peptides: concepts and applications. In: Misra A, ed. *Challenges in Delivery of Therapeutic Genomics and Proteomics*. London: Elsevier; 2011. p. 481-529. doi: [10.1016/b978-0-12-384964-9.00010-4](https://doi.org/10.1016/b978-0-12-384964-9.00010-4)
 120. Woodley JF. Enzymatic barriers for GI peptide and protein delivery. *Crit Rev Ther Drug Carrier Syst* 1994;11(2-3):61-95.
 121. Langguth P, Bohner V, Heizmann J, Merkle HP, Wolfram S, Amidon GL, et al. The challenge of proteolytic enzymes in intestinal peptide delivery. *J Control Release* 1997;46(1-2):39-57. doi: [10.1016/s0168-3659\(96\)01586-6](https://doi.org/10.1016/s0168-3659(96)01586-6)
 122. Mersny R. Challenges for the Oral Delivery of Proteins and Peptides: Theoretical and Practical Approaches to Their Delivery. In: Capsugel Symposia Series. Greenwood, SC; 1991. p. 452.
 123. Ganong WF. Regulation of gastrointestinal function. In: Ganong WF, ed. *Review of Medical Physiology*. 12th ed. Lange Medical Publications; 1983. p. 394-420.
 124. Mersny R. Challenges for the Oral Delivery of Proteins and Peptides: Theoretical and Practical Approaches to Their Delivery. In: Capsugel Library Symposia series. 1991. p. 45-52.
 125. Hayakawa E, Lee VH. Aminopeptidase activity in the jejunal and ileal Peyer's patches of the albino rabbit. *Pharm Res* 1992;9(4):535-40. doi: [10.1023/a:1015800615674](https://doi.org/10.1023/a:1015800615674)
 126. Skillman JJ, Gould SA, Chung RS, Silen W. The gastric mucosal barrier: clinical and experimental studies in critically ill and normal man, and in the rabbit. *Ann Surg* 1970;172(4):564-84. doi: [10.1097/00000658-197010000-00004](https://doi.org/10.1097/00000658-197010000-00004)
 127. Meyer RA, McGinley D, Posalaky Z. The gastric mucosal barrier: structure of intercellular junctions in the dog. *J Ultrastruct Res* 1984;86(2):192-201. doi: [10.1016/s0022-5320\(84\)80058-1](https://doi.org/10.1016/s0022-5320(84)80058-1)
 128. MacAdam A. The effect of gastro-intestinal mucus on drug absorption. *Adv Drug Deliv Rev* 1993;11(3):201-20. doi: [10.1016/0169-409x\(93\)90010-2](https://doi.org/10.1016/0169-409x(93)90010-2)
 129. Phelps CF. Biosynthesis of mucus glycoprotein. *Br Med Bull* 1978;34(1):43-8. doi: [10.1093/oxfordjournals.bmb.a071456](https://doi.org/10.1093/oxfordjournals.bmb.a071456)
 130. Allen A, Garner A. Mucus and bicarbonate secretion in the stomach and their possible role in mucosal protection. *Gut* 1980;21(3):249-62. doi: [10.1136/gut.21.3.249](https://doi.org/10.1136/gut.21.3.249)
 131. Kompella UB, Lee VH. Delivery systems for penetration enhancement of peptide and protein drugs: design considerations. *Adv Drug Deliv Rev* 2001;46(1-3):211-45. doi: [10.1016/s0169-409x\(00\)00137-x](https://doi.org/10.1016/s0169-409x(00)00137-x)
 132. Kerss S, Allen A, Garner A. A simple method for measuring thickness of the mucus gel layer adherent to rat, frog and human gastric mucosa: influence of feeding, prostaglandin, N-acetylcysteine and other agents. *Clin Sci (Lond)* 1982;63(2):187-95. doi: [10.1042/cs0630187](https://doi.org/10.1042/cs0630187)
 133. Pankhurst QA, Connolly J, Jones SK, Dobson J. Applications of magnetic nanoparticles in biomedicine. *J Phys D Appl Phys* 2003;36(13):R167. doi: [10.1088/0022-3727/36/13/201](https://doi.org/10.1088/0022-3727/36/13/201)
 134. Kreuter J. Nanoparticulate systems in drug delivery and targeting. *J Drug Target* 1995;3(3):171-3. doi: [10.3109/10611869509015940](https://doi.org/10.3109/10611869509015940)
 135. Jabir NR, Tabrez S, Ashraf GM, Shakil S, Damanhour GA, Kamal MA. Nanotechnology-based approaches in anticancer research. *Int J Nanomedicine* 2012;7:4391-408. doi: [10.2147/ijn.s33838](https://doi.org/10.2147/ijn.s33838)
 136. Li SD, Huang L. Pharmacokinetics and biodistribution of nanoparticles. *Mol Pharm* 2008;5(4):496-504. doi: [10.1021/mp800049w](https://doi.org/10.1021/mp800049w)
 137. Pattni BS, Chupin VV, Torchilin VP. New developments in liposomal drug delivery. *Chem Rev* 2015;115(19):10938-66. doi: [10.1021/acs.chemrev.5b00046](https://doi.org/10.1021/acs.chemrev.5b00046)
 138. Sawant RR, Torchilin VP. Challenges in development of targeted liposomal therapeutics. *AAPS J* 2012;14(2):303-15. doi: [10.1208/s12248-012-9330-0](https://doi.org/10.1208/s12248-012-9330-0)
 139. Vieira DB, Gamarra LF. Getting into the brain: liposome-based strategies for effective drug delivery across the blood-brain barrier. *Int J Nanomedicine* 2016;11:5381-414. doi: [10.2147/](https://doi.org/10.2147/)

- ijn.s117210
140. Craparo EF, Bondi ML, Pitarresi G, Cavallaro G. Nanoparticulate systems for drug delivery and targeting to the central nervous system. *CNS Neurosci Ther* 2011;17(6):670-7. doi: [10.1111/j.1755-5949.2010.00199.x](https://doi.org/10.1111/j.1755-5949.2010.00199.x)
 141. Yang H. Nanoparticle-mediated brain-specific drug delivery, imaging, and diagnosis. *Pharm Res* 2010;27(9):1759-71. doi: [10.1007/s11095-010-0141-7](https://doi.org/10.1007/s11095-010-0141-7)
 142. Chang HI, Yeh MK. Clinical development of liposome-based drugs: formulation, characterization, and therapeutic efficacy. *Int J Nanomedicine* 2012;7:49-60. doi: [10.2147/ijn.s26766](https://doi.org/10.2147/ijn.s26766)
 143. Chen W, Li H, Liu Z, Yuan W. Lipopolyplex for therapeutic gene delivery and its application for the treatment of Parkinson's disease. *Front Aging Neurosci* 2016;8:68. doi: [10.3389/fnagi.2016.00068](https://doi.org/10.3389/fnagi.2016.00068)
 144. Huang H, Cruz W, Chen J, Zheng G. Learning from biology: synthetic lipoproteins for drug delivery. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2015;7(3):298-314. doi: [10.1002/wnan.1308](https://doi.org/10.1002/wnan.1308)
 145. Ag Selecı D, Selecı M, Walter JG, Stahl F, Scheper T. Niosomes as nanoparticulate drug carriers: fundamentals and recent applications. *J Nanomater* 2016;2016:7372306. doi: [10.1155/2016/7372306](https://doi.org/10.1155/2016/7372306)
 146. Moghassemi S, Hadjizadeh A. Nano-niosomes as nanoscale drug delivery systems: an illustrated review. *J Control Release* 2014;185:22-36. doi: [10.1016/j.jconrel.2014.04.015](https://doi.org/10.1016/j.jconrel.2014.04.015)
 147. Chaurasia S, Dogra SS. Transfersomes: novel approach for intranasal delivery. *Eur J Pharm Med Res* 2017;4(3):192-203.
 148. Ezzati Nazhad Dolatabadi J, Omidı Y. Solid lipid-based nanocarriers as efficient targeted drug and gene delivery systems. *TrAC Trends Anal Chem* 2016;77:100-8. doi: [10.1016/j.trac.2015.12.016](https://doi.org/10.1016/j.trac.2015.12.016)
 149. Beloqui A, Solinıs M, Rodrıguez-Gascın A, Almeida AJ, Prıt V. Nanostructured lipid carriers: promising drug delivery systems for future clinics. *Nanomedicine* 2016;12(1):143-61. doi: [10.1016/j.nano.2015.09.004](https://doi.org/10.1016/j.nano.2015.09.004)
 150. Eldem T, Speiser P, Hincal A. Optimization of spray-dried and -congealed lipid micropellets and characterization of their surface morphology by scanning electron microscopy. *Pharm Res* 1991;8(1):47-54. doi: [10.1023/a:1015874121860](https://doi.org/10.1023/a:1015874121860)
 151. Speiser P. *Lipidnanopellets als Tragersystem fur Arzneimittel zur peroralen Anwendung*. European Patent EP 0167825, 1990.
 152. Wei L, Yang Y, Shi K, Wu J, Zhao W, Mo J. Preparation and characterization of loperamide-loaded dynasan 114 solid lipid nanoparticles for increased oral absorption in the treatment of diarrhea. *Front Pharmacol* 2016;7:332. doi: [10.3389/fphar.2016.00332](https://doi.org/10.3389/fphar.2016.00332)
 153. Subedi RK, Kang KW, Choi HK. Preparation and characterization of solid lipid nanoparticles loaded with doxorubicin. *Eur J Pharm Sci* 2009;37(3-4):508-13. doi: [10.1016/j.ejps.2009.04.008](https://doi.org/10.1016/j.ejps.2009.04.008)
 154. Muller RH, Runge SA. Solid lipid nanoparticles (SLN) for controlled drug delivery. In: Benita S, ed. *Submicron Emulsions in Drug Targeting and Delivery*. Amsterdam: Harwood Academic Publishers; 1998. p. 219-34.
 155. uner M, Yener G. Importance of solid lipid nanoparticles (SLN) in various administration routes and future perspectives. *Int J Nanomedicine* 2007;2(3):289-300.
 156. Mukherjee S, Ray S, Thakur RS. Solid lipid nanoparticles: a modern formulation approach in drug delivery system. *Indian J Pharm Sci* 2009;71(4):349-58. doi: [10.4103/0250-474x.57282](https://doi.org/10.4103/0250-474x.57282)
 157. Muller RH, Mader K, Gohla S. Solid lipid nanoparticles (SLN) for controlled drug delivery - a review of the state of the art. *Eur J Pharm Biopharm* 2000;50(1):161-77. doi: [10.1016/s0939-6411\(00\)00087-4](https://doi.org/10.1016/s0939-6411(00)00087-4)
 158. Mehnert W, Mader K. Solid lipid nanoparticles: production, characterization and applications. *Adv Drug Deliv Rev* 2001;47(2-3):165-96. doi: [10.1016/s0169-409x\(01\)00105-3](https://doi.org/10.1016/s0169-409x(01)00105-3)
 159. Pandey S, Shaikh F, Gupta A, Tripathi P, Yadav JS. A Recent Update: Solid Lipid Nanoparticles for Effective Drug Delivery. *Adv Pharm Bull.* 2022;12(1):17-33. doi:[10.34172/apb.2022.007](https://doi.org/10.34172/apb.2022.007)
 160. Muller RH, Mehnert W, Lucks J. Solid lipid nanoparticles (SLN)-an alternative colloidal carrier system for controlled drug delivery. *Eur J Pharm Biopharm* 1995;41(1):62-9.
 161. zur Muhlen A, Mehnert W. Drug release and release mechanism of prednisolone loaded solid lipid nanoparticles. *Pharmazie* 1998;53(8):552-5.
 162. Souto EB, Muller RH. Lipid nanoparticles: effect on bioavailability and pharmacokinetic changes. In: Schafer-Korting M, ed. *Drug Delivery*. Berlin, Heidelberg: Springer; 2010. p. 115-41. doi: [10.1007/978-3-642-00477-3_4](https://doi.org/10.1007/978-3-642-00477-3_4)
 163. Shah R, Eldridge D, Palombo E, Harding I. *Lipid Nanoparticles: Production, Characterization and Stability*. New York, NY: Springer International Publishing; 2015.
 164. Gaba B, Fazil M, Ali A, Baboota S, Sahni JK, Ali J. Nanostructured lipid (NLCs) carriers as a bioavailability enhancement tool for oral administration. *Drug Deliv* 2015;22(6):691-700. doi: [10.3109/10717544.2014.898110](https://doi.org/10.3109/10717544.2014.898110)
 165. Shegokar R, Singh KK, Muller RH. Production & stability of stavudine solid lipid nanoparticles--from lab to industrial scale. *Int J Pharm* 2011;416(2):461-70. doi: [10.1016/j.ijpharm.2010.08.014](https://doi.org/10.1016/j.ijpharm.2010.08.014)
 166. Yu YH, Kim E, Park DE, Shim G, Lee S, Kim YB, et al. Cationic solid lipid nanoparticles for co-delivery of paclitaxel and siRNA. *Eur J Pharm Biopharm* 2012;80(2):268-73. doi: [10.1016/j.ejpb.2011.11.002](https://doi.org/10.1016/j.ejpb.2011.11.002)
 167. Xie S, Zhu L, Dong Z, Wang X, Wang Y, Li X, et al. Preparation, characterization and pharmacokinetics of enrofloxacin-loaded solid lipid nanoparticles: influences of fatty acids. *Colloids Surf B Biointerfaces* 2011;83(2):382-7. doi: [10.1016/j.colsurfb.2010.12.014](https://doi.org/10.1016/j.colsurfb.2010.12.014)
 168. Gastaldi L, Battaglia L, Peira E, Chirio D, Muntoni E, Solazzi I, et al. Solid lipid nanoparticles as vehicles of drugs to the brain: current state of the art. *Eur J Pharm Biopharm* 2014;87(3):433-44. doi: [10.1016/j.ejpb.2014.05.004](https://doi.org/10.1016/j.ejpb.2014.05.004)
 169. Silva AC, Gonzalez-Mira E, Garcıa ML, Egea MA, Fonseca J, Silva R, et al. Preparation, characterization and biocompatibility studies on risperidone-loaded solid lipid nanoparticles (SLN): high pressure homogenization versus ultrasound. *Colloids Surf B Biointerfaces* 2011;86(1):158-65. doi: [10.1016/j.colsurfb.2011.03.035](https://doi.org/10.1016/j.colsurfb.2011.03.035)
 170. Dwivedi P, Khatik R, Khandelwal K, Taneja I, Raju KS, Wahajuddin, et al. Pharmacokinetics study of arteether loaded solid lipid nanoparticles: an improved oral bioavailability in rats. *Int J Pharm* 2014;466(1-2):321-7. doi: [10.1016/j.ijpharm.2014.03.036](https://doi.org/10.1016/j.ijpharm.2014.03.036)
 171. Raza K, Singh B, Singal P, Wadhwa S, Katare OP. Systematically optimized biocompatible isotretinoin-loaded solid lipid nanoparticles (SLNs) for topical treatment of acne. *Colloids Surf B Biointerfaces* 2013;105:67-74. doi: [10.1016/j.colsurfb.2012.12.043](https://doi.org/10.1016/j.colsurfb.2012.12.043)
 172. Ravi PR, Vats R, Dalal V, Murthy AN. A hybrid design to optimize preparation of lopinavir loaded solid lipid nanoparticles and comparative pharmacokinetic evaluation with marketed lopinavir/ritonavir coformulation. *J Pharm Pharmacol* 2014;66(7):912-26. doi: [10.1111/jphp.12217](https://doi.org/10.1111/jphp.12217)
 173. Kheradmandnia S, Vasheghani-Farahani E, Nosrati M, Atyabi F. Preparation and characterization of ketoprofen-loaded solid lipid nanoparticles made from beeswax and carnauba wax. *Nanomedicine* 2010;6(6):753-9. doi: [10.1016/j.nano.2010.06.003](https://doi.org/10.1016/j.nano.2010.06.003)

174. Madan J, Pandey RS, Jain V, Katare OP, Chandra R, Katyal A. Poly (ethylene)-glycol conjugated solid lipid nanoparticles of noscapine improve biological half-life, brain delivery and efficacy in glioblastoma cells. *Nanomedicine* 2013;9(4):492-503. doi: [10.1016/j.nano.2012.10.003](https://doi.org/10.1016/j.nano.2012.10.003)
175. Mehnert W, Mäder K. Solid lipid nanoparticles: production, characterization and applications. *Adv Drug Deliv Rev* 2001;47(2-3):165-96. doi: [10.1016/s0169-409x\(01\)00105-3](https://doi.org/10.1016/s0169-409x(01)00105-3)
176. Wang S, Chen T, Chen R, Hu Y, Chen M, Wang Y. Emodin loaded solid lipid nanoparticles: preparation, characterization and antitumor activity studies. *Int J Pharm* 2012;430(1-2):238-46. doi: [10.1016/j.ijpharm.2012.03.027](https://doi.org/10.1016/j.ijpharm.2012.03.027)
177. Kuo YC, Wang CC. Cationic solid lipid nanoparticles with primary and quaternary amines for release of saquinavir and biocompatibility with endothelia. *Colloids Surf B Biointerfaces* 2013;101:101-5. doi: [10.1016/j.colsurfb.2012.06.002](https://doi.org/10.1016/j.colsurfb.2012.06.002)
178. Cai S, Yang Q, Bagby TR, Forrest ML. Lymphatic drug delivery using engineered liposomes and solid lipid nanoparticles. *Adv Drug Deliv Rev* 2011;63(10-11):901-8. doi: [10.1016/j.addr.2011.05.017](https://doi.org/10.1016/j.addr.2011.05.017)
179. Nafee N, Husari A, Maurer CK, Lu C, de Rossi C, Steinbach A, et al. Antibiotic-free nanotherapeutics: ultra-small, mucus-penetrating solid lipid nanoparticles enhance the pulmonary delivery and anti-virulence efficacy of novel quorum sensing inhibitors. *J Control Release* 2014;192:131-40. doi: [10.1016/j.jconrel.2014.06.055](https://doi.org/10.1016/j.jconrel.2014.06.055)
180. Almeida AJ, Souto E. Solid lipid nanoparticles as a drug delivery system for peptides and proteins. *Adv Drug Deliv Rev* 2007;59(6):478-90. doi: [10.1016/j.addr.2007.04.007](https://doi.org/10.1016/j.addr.2007.04.007)
181. Shah RM, Malherbe F, Eldridge D, Palombo EA, Harding IH. Physicochemical characterization of solid lipid nanoparticles (SLNs) prepared by a novel microemulsion technique. *J Colloid Interface Sci* 2014;428:286-94. doi: [10.1016/j.jcis.2014.04.057](https://doi.org/10.1016/j.jcis.2014.04.057)
182. Soares S, Fonte P, Costa A, Andrade J, Seabra V, Ferreira D, et al. Effect of freeze-drying, cryoprotectants and storage conditions on the stability of secondary structure of insulin-loaded solid lipid nanoparticles. *Int J Pharm* 2013;456(2):370-81. doi: [10.1016/j.ijpharm.2013.08.076](https://doi.org/10.1016/j.ijpharm.2013.08.076)
183. Kalhapure RS, Mocktar C, Sikwal DR, Sonawane SJ, Kathiravan MK, Skelton A, et al. Ion pairing with linoleic acid simultaneously enhances encapsulation efficiency and antibacterial activity of vancomycin in solid lipid nanoparticles. *Colloids Surf B Biointerfaces* 2014;117:303-11. doi: [10.1016/j.colsurfb.2014.02.045](https://doi.org/10.1016/j.colsurfb.2014.02.045)
184. Hao J, Wang F, Wang X, Zhang D, Bi Y, Gao Y, et al. Development and optimization of baicalin-loaded solid lipid nanoparticles prepared by coacervation method using central composite design. *Eur J Pharm Sci* 2012;47(2):497-505. doi: [10.1016/j.ejps.2012.07.006](https://doi.org/10.1016/j.ejps.2012.07.006)
185. Pardeshi C, Rajput P, Belgamwar V, Tekade A, Patil G, Chaudhary K, et al. [Solid lipid based nanocarriers: an overview]. *Acta Pharm* 2013;62(4):433-72. doi: [10.2478/v10007-012-0040-z](https://doi.org/10.2478/v10007-012-0040-z)
186. Muchow M, Maincent P, Müller RH. Lipid nanoparticles with a solid matrix (SLN, NLC, LDC) for oral drug delivery. *Drug Dev Ind Pharm* 2008;34(12):1394-405. doi: [10.1080/03639040802130061](https://doi.org/10.1080/03639040802130061)
187. Yun Y, Cho YW, Park K. Nanoparticles for oral delivery: targeted nanoparticles with peptidic ligands for oral protein delivery. *Adv Drug Deliv Rev*. 2013;65(6):822-832. doi:[10.1016/j.addr.2012.10.007](https://doi.org/10.1016/j.addr.2012.10.007)
188. Borgström B. Importance of phospholipids, pancreatic phospholipase A2, and fatty acid for the digestion of dietary fat: in vitro experiments with the porcine enzymes. *Gastroenterology* 1980;78(5 Pt 1):954-62.
189. Borgström B, Donnér J. The polar interactions between pancreatic lipase, colipase and the triglyceride substrate. *FEBS Lett* 1977;83(1):23-6. doi: [10.1016/0014-5793\(77\)80633-9](https://doi.org/10.1016/0014-5793(77)80633-9)
190. Scow RO, Olivecrona T. Effect of albumin on products formed from chylomicron triacylglycerol by lipoprotein lipase in vitro. *Biochim Biophys Acta* 1977;487(3):472-86. doi: [10.1016/0005-2760\(77\)90217-x](https://doi.org/10.1016/0005-2760(77)90217-x)
191. Olbrich C, Müller RH. Enzymatic degradation of SLN-effect of surfactant and surfactant mixtures. *Int J Pharm* 1999;180(1):31-9. doi: [10.1016/s0378-5173\(98\)00404-9](https://doi.org/10.1016/s0378-5173(98)00404-9)
192. Müller RH, Olbrich C. Solid lipid nanoparticles: phagocytic uptake, in vitro cytotoxicity and in vitro biodegradation, 2nd communication. *Pharm Ind* 1999;61(6):564-9.
193. Olbrich C, Mehnert W, Müller RH. In vitro degradation properties of solid lipid nanoparticles. in: Proc. 2nd World Meeting APGI/APV; Paris; 1998. p. 577-8.
194. Müller RH, Rühl D, Runge SA. Biodegradation of solid lipid nanoparticles as a function of lipase incubation time. *Int J Pharm* 1996;144(1):115-21. doi: [10.1016/s0378-5173\(96\)04731-x](https://doi.org/10.1016/s0378-5173(96)04731-x)
195. Olbrich C, Mehnert W, Müller RH. In vitro degradation properties of solid lipid nanoparticles. in: Proc. 2nd World Meeting APGI/APV; Paris; 1998. p. 627-8.
196. Yang S, Zhu J, Lu Y, Liang B, Yang C. Body distribution of camptothecin solid lipid nanoparticles after oral administration. *Pharm Res* 1999;16(5):751-7. doi: [10.1023/a:1018888927852](https://doi.org/10.1023/a:1018888927852)
197. Penkler L, Müller RH, Runge SA. *Pharmaceutical Cyclosporin Formulation with Improved Biopharmaceutical Properties, Improved Physical Quality and Greater Stability, and Method for Producing Said Formulation*. WO 99/56733, 1999.
198. Bargoni A, Cavalli R, Caputo O, Fundarò A, Gasco MR, Zara GP. Solid lipid nanoparticles in lymph and plasma after duodenal administration to rats. *Pharm Res* 1998;15(5):745-50. doi: [10.1023/a:1011975120776](https://doi.org/10.1023/a:1011975120776)
199. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65(1-2):55-63. doi: [10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
200. Müller RH, Maaßen S, Weyhers H, Specht F, Lucks JS. Cytotoxicity of magnetite-loaded polylactide, polylactide/glycolide particles and solid lipid nanoparticles. *Int J Pharm* 1996;138(1):85-94. doi: [10.1016/0378-5173\(96\)04539-5](https://doi.org/10.1016/0378-5173(96)04539-5)
201. Marcato PD, Durán N. Cytotoxicity and genotoxicity of solid lipid nanoparticles. In: Durán N, Guterres SS, Alves OL, eds. *Nanotoxicology: Materials, Methodologies, and Assessments*. New York, NY: Springer; 2014. p. 229-44. doi: [10.1007/978-1-4614-8993-1_10](https://doi.org/10.1007/978-1-4614-8993-1_10)
202. Almeida H, Amaral MH, Lobão P, Silva AC, Lobo JM. Applications of polymeric and lipid nanoparticles in ophthalmic pharmaceutical formulations: present and future considerations. *J Pharm Pharm Sci* 2014;17(3):278-93. doi: [10.18433/j3dp43](https://doi.org/10.18433/j3dp43)
203. Sandeep Kalepu, Mohanvarma Manthina, Veerabhadhraswamy Padavala. Oral lipid-based drug delivery systems – an overview. *Acta Pharmaceutica Sinica B*. 2013; 3 (6): 361-372. doi:[10.1016/j.apsb.2013.10.001](https://doi.org/10.1016/j.apsb.2013.10.001)
204. Doktorovova S, Souto EB, Silva AM. Nanotoxicology applied to solid lipid nanoparticles and nanostructured lipid carriers - a systematic review of in vitro data. *Eur J Pharm Biopharm* 2014;87(1):1-18. doi: [10.1016/j.ejpb.2014.02.005](https://doi.org/10.1016/j.ejpb.2014.02.005)
205. Fortunov S, Peneva P, Andonova V, Kassarova M. Toxicity assessment of drug delivery nanocarriers. *Sci Technol* 2016;6(1):262-6.
206. Hillegass JM, Shukla A, Lathrop SA, MacPherson MB, Fukagawa NK, Mossman BT. Assessing nanotoxicity in cells in vitro. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2010;2(3):219-31. doi: [10.1002/wnan.54](https://doi.org/10.1002/wnan.54)
207. Maaßen S, Schwarz C, Mehnert W, Lucks JS, Yunis-Specht

- F, Muller BW, et al. Comparison of cytotoxicity between polyester nanoparticles and solid lipid nanoparticles. *Proc Int Symp Control Rel Bioact Mater* 1993;20:490-1.
208. Müller RH, Maassen S, Weyhers H, Mehnert W. Phagocytic uptake and cytotoxicity of solid lipid nanoparticles (SLN) sterically stabilized with poloxamine 908 and poloxamer 407. *J Drug Target* 1996;4(3):161-70. doi: [10.3109/10611869609015973](https://doi.org/10.3109/10611869609015973)
209. Yu YH, Kim E, Park DE, Shim G, Lee S, Kim YB, et al. Cationic solid lipid nanoparticles for co-delivery of paclitaxel and siRNA. *Eur J Pharm Biopharm* 2012;80(2):268-73. doi: [10.1016/j.ejpb.2011.11.002](https://doi.org/10.1016/j.ejpb.2011.11.002)
210. Shi S, Han L, Deng L, Zhang Y, Shen H, Gong T, et al. Dual drugs (microRNA-34a and paclitaxel)-loaded functional solid lipid nanoparticles for synergistic cancer cell suppression. *J Control Release* 2014;194:228-37. doi: [10.1016/j.jconrel.2014.09.005](https://doi.org/10.1016/j.jconrel.2014.09.005)
211. Jin J, Bae KH, Yang H, Lee SJ, Kim H, Kim Y, et al. In vivo specific delivery of c-Met siRNA to glioblastoma using cationic solid lipid nanoparticles. *Bioconjug Chem* 2011;22(12):2568-72. doi: [10.1021/bc200406n](https://doi.org/10.1021/bc200406n)
212. Baek JS, Cho CW. Controlled release and reversal of multidrug resistance by co-encapsulation of paclitaxel and verapamil in solid lipid nanoparticles. *Int J Pharm* 2015;478(2):617-24. doi: [10.1016/j.ijpharm.2014.12.018](https://doi.org/10.1016/j.ijpharm.2014.12.018)
213. Kalhapure RS, Mocktar C, Sikwal DR, Sonawane SJ, Kathiravan MK, Skelton A, et al. Ion pairing with linoleic acid simultaneously enhances encapsulation efficiency and antibacterial activity of vancomycin in solid lipid nanoparticles. *Colloids Surf B Biointerfaces* 2014;117:303-11. doi: [10.1016/j.colsurfb.2014.02.045](https://doi.org/10.1016/j.colsurfb.2014.02.045)
214. Morishita M, Morishita I, Takayama K, Machida Y, Nagai T. Site-dependent effect of aprotinin, sodium caprate, Na₂EDTA and sodium glycocholate on intestinal absorption of insulin. *Biol Pharm Bull* 1993;16(1):68-72. doi: [10.1248/bpb.16.68](https://doi.org/10.1248/bpb.16.68)
215. Lane ME, O'Driscoll C M, Corrigan OL. Quantitative estimation of the effects of bile salt surfactant systems on insulin stability and permeability in the rat intestine using a mass balance model. *J Pharm Pharmacol* 2005;57(2):169-75. doi: [10.1211/0022357055434](https://doi.org/10.1211/0022357055434)
216. Sharma G, Wilson K, van der Walle CF, Sattar N, Petrie JR, Ravi Kumar MN. Microemulsions for oral delivery of insulin: design, development and evaluation in streptozotocin induced diabetic rats. *Eur J Pharm Biopharm* 2010;76(2):159-69. doi: [10.1016/j.ejpb.2010.07.002](https://doi.org/10.1016/j.ejpb.2010.07.002)
217. Cilek A, Celebi N, Tirnaksiz F, Tay A. A lecithin-based microemulsion of rh-insulin with aprotinin for oral administration: Investigation of hypoglycemic effects in non-diabetic and STZ-induced diabetic rats. *Int J Pharm* 2005;298(1):176-85. doi: [10.1016/j.ijpharm.2005.04.016](https://doi.org/10.1016/j.ijpharm.2005.04.016)
218. Elsayed A, Remawi MA, Qinna N, Farouk A, Badwan A. Formulation and characterization of an oily-based system for oral delivery of insulin. *Eur J Pharm Biopharm* 2009;73(2):269-79. doi: [10.1016/j.ejpb.2009.06.004](https://doi.org/10.1016/j.ejpb.2009.06.004)
219. Toorisaka E, Ono H, Arimori K, Kamiya N, Goto M. Hypoglycemic effect of surfactant-coated insulin solubilized in a novel solid-in-oil-in-water (S/O/W) emulsion. *Int J Pharm* 2003;252(1-2):271-4. doi: [10.1016/s0378-5173\(02\)00674-9](https://doi.org/10.1016/s0378-5173(02)00674-9)
220. Watnasirichaikul S, Rades T, Tucker IG, Davies NM. In-vitro release and oral bioactivity of insulin in diabetic rats using nanocapsules dispersed in biocompatible microemulsion. *J Pharm Pharmacol* 2002;54(4):473-80. doi: [10.1211/0022357021778736](https://doi.org/10.1211/0022357021778736)
221. Constantinides PP, Scalart JP, Lancaster C, Marcello J, Marks G, Ellens H, et al. Formulation and intestinal absorption enhancement evaluation of water-in-oil microemulsions incorporating medium-chain glycerides. *Pharm Res* 1994;11(10):1385-90. doi: [10.1023/a:1018927402875](https://doi.org/10.1023/a:1018927402875)
222. Ritschel WA. Microemulsions for improved peptide absorption from the gastrointestinal tract. *Methods Find Exp Clin Pharmacol* 1991;13(3):205-20.
223. Celebi N, Türkyilmaz A, Gönül B, Ozogul C. Effects of epidermal growth factor microemulsion formulation on the healing of stress-induced gastric ulcers in rats. *J Control Release* 2002;83(2):197-210. doi: [10.1016/s0168-3659\(02\)00198-0](https://doi.org/10.1016/s0168-3659(02)00198-0)
224. Rao SV, Yajurvedi K, Shao J. Self-nanoemulsifying drug delivery system (SNEDDS) for oral delivery of protein drugs: III. In vivo oral absorption study. *Int J Pharm* 2008;362(1-2):16-9. doi: [10.1016/j.ijpharm.2008.05.015](https://doi.org/10.1016/j.ijpharm.2008.05.015)
225. Lyons KC, Charman WN, Miller R, Porter CJ. Factors limiting the oral bioavailability of N-acetylglucosaminyl-N-acetylmuramyl dipeptide (GMDP) and enhancement of absorption in rats by delivery in a water-in-oil microemulsion. *Int J Pharm* 2000;199(1):17-28. doi: [10.1016/s0378-5173\(00\)00349-5](https://doi.org/10.1016/s0378-5173(00)00349-5)
226. Zheng JY, Fulu MY. Decrease of genital organ weights and plasma testosterone levels in rats following oral administration of leuprolide microemulsion. *Int J Pharm* 2006;307(2):209-15. doi: [10.1016/j.ijpharm.2005.10.007](https://doi.org/10.1016/j.ijpharm.2005.10.007)
227. Fagerholm U, Sjöström B, Sroka-Markovic J, Wijk A, Svensson M, Lennernäs H. The effect of a drug-delivery system consisting of soybean phosphatidyl choline and medium-chain monoacylglycerol on the intestinal permeability of hexarelin in the rat. *J Pharm Pharmacol* 1998;50(5):467-73. doi: [10.1111/j.2042-7158.1998.tb06187.x](https://doi.org/10.1111/j.2042-7158.1998.tb06187.x)
228. Aungst BJ, Saitoh H, Burcham DL, Huang SM, Mousa SA, Hussain MA. Enhancement of the intestinal absorption of peptides and nonpeptides. *J Control Release* 1996;41(1):19-31. doi: [10.1016/0168-3659\(96\)01353-3](https://doi.org/10.1016/0168-3659(96)01353-3)
229. Hastewell J, Lynch S, Williamson I, Fox R, Mackay M. Absorption of human calcitonin across the rat colon in vivo. *Clin Sci (Lond)* 1992;82(5):589-94. doi: [10.1042/cs0820589](https://doi.org/10.1042/cs0820589)
230. New R, Littlewood G, Guard P, Browning I, Hotten P. Intestinal delivery of calcitonin in pig. *Int J Pharm* 1997;156(1):1-8. doi: [10.1016/s0378-5173\(97\)00130-0](https://doi.org/10.1016/s0378-5173(97)00130-0)
231. Fan Y, Li X, Zhou Y, Fan C, Wang X, Huang Y, et al. Improved intestinal delivery of salmon calcitonin by water-in-oil microemulsions. *Int J Pharm* 2011;416(1):323-30. doi: [10.1016/j.ijpharm.2011.06.029](https://doi.org/10.1016/j.ijpharm.2011.06.029)
232. Yoshiura H, Tahara Y, Hashida M, Kamiya N, Hirata A, Fujii T, et al. Design and in vivo evaluation of solid-in-oil suspension for oral delivery of human growth hormone. *Biochem Eng J* 2008;41(2):106-10. doi: [10.1016/j.bej.2008.04.001](https://doi.org/10.1016/j.bej.2008.04.001)
233. FDA. 21 CFR Part 182, Substances Generally Recognized as Safe. FDA; 2010.
234. U.S. Department of Health and Human Services, Food and Drug Administration. Guidance for Industry: Nonclinical Studies for the Safety Evaluation of Pharmaceutical Excipients. Office of Training and Communication, Division of Drug Information, HFD-240, Center for Drug Evaluation and Research, Food and Drug Administration, or Office of Communication, Training, and Manufacturers Assistance, HFM-40, Center for Biologics Evaluation and Research, Food and Drug Administration. May 2005. Available from: <http://www.fda.gov/cder/guidance/5544fnl.pdf>.
235. U.S. Food and Drug Administration. Inactive Ingredients Search for Approved Drug Products. Division of Labeling and Program Support, Office of Generic Drugs, Center for Drug Evaluation and Research, Food and Drug Administration; 2007. Available from: <http://www.accessdata.fda.gov/scripts/cder/iig/index.cfm>.
236. Maincent P. The regulatory environment: the challenges for lipid-based formulation. *Bull Tech Gattefossé* 2007;100:47-9.
237. Chen ML. Lipid excipients and delivery systems for pharmaceutical development: a regulatory perspective. *Adv Drug Deliv Rev* 2008;60(6):768-77. doi: [10.1016/j.addr.2007.09.010](https://doi.org/10.1016/j.addr.2007.09.010)