

## Essential Guide of Analysis Methods Applied to Silver Complexes with Antibacterial Quinolones

Aura Rusu<sup>1</sup>, Gabriel Hancu<sup>1\*</sup>, Silvia Imre<sup>2</sup>

<sup>1</sup> Faculty of Pharmacy, Pharmaceutical Chemistry Department, University of Medicine and Pharmacy of TîrguMureş, Tîrgu Mureş, Romania.

<sup>2</sup> Faculty of Pharmacy, Department of Analytical Chemistry and Drug Analysis, University of Medicine and Pharmacy of TîrguMureş, Tîrgu Mureş, Romania.

### Article info

#### Article History:

Received: 24 February 2018

Revised: 11 May 2018

Accepted: 24 May 2018

ePublished: 19 June 2018

#### Keywords:

- 4-quinolones
- Fluoroquinolones
- Analysis methods
- Organometallic compounds
- Silver

### Abstract

To describe the chemical structure and characterize physico-chemical properties of organometallic complexes it is necessary to use a complex set of analysis methods. Thus, this review has been compiled as a relevant guide which includes the most commonly used methods of analysis in the study of silver complexes with antibacterial quinolones, compounds with promising biological potential. This selection of analysis methods puts on balance the obtained data and the accessibility of the experimental approach. The steps to follow in order to obtain reliable structural information about organometallic complexes of silver, particularly the silver complexes of antibacterial quinolones, are established and presented in the review.

### Introduction

The potential pharmacological effects of the organometallic complexes can be considered appealing and fascinating. An excellent example for this is the antibacterial activity of metal complexes with quinolones.<sup>1,2</sup> Increased biological effect can be attributed to the interaction of metals with an active organic molecule, and also to the synergism of the antibacterial effect of the complex components.<sup>3,4</sup> The metal complexes with antibacterial quinolones were summarized in previously published reviews.<sup>5-8</sup> Moreover, organometallic complexes show other potential pharmacological effects including anti-inflammatory, anti-cancer, anti-oxidant, anti-parasitic, and even insulin-mimetic properties.<sup>3,4,8,9</sup> Today, the increased bacterial resistance is a worldwide menace; consequently, the discovery of new compounds with an efficient antibacterial effect is a high priority in modern pharmaceutical research. Thereby, organometallic complexes can be a valuable alternative in the fight against bacterial resistance.<sup>10,11</sup> The study of organometallic complexes also includes the development of efficient methods for their analysis. The current paper presents a review of several analytical methods applied in the analysis of silver complexes with antibacterial quinolones.<sup>12-14</sup>

Analysis methods are used for confirmation of the chemical structure of the obtained organometallic complexes. Thus, the most critical aspect is the determination of sites from quinolone structure where the

silver chelation takes place. Quinolone representatives may be coordinated to the silver ion through the 3-carboxyl and 4-pyridone groups.<sup>15,16</sup> However, most of the quinolones may be coordinated through N-heterocycle from 7 positions.<sup>1,2,17</sup> Physico-chemical properties, stability and biological effects of the complexes may vary according to coordination site of silver ion.<sup>12</sup>

### Characterization of quinolone-metal complexes

The antibacterial quinolones (4-quinolones) are a class of synthetic antibiotics whose representatives from the first generations have been used primarily for the treatment of urinary tract infections. Newer generations comprise fluorinated derivatives (fluoroquinolones) with broad-spectrum activity. Antibacterial quinolones compounds present a particular mechanism of action by inhibition of DNA replication. Targets for antibacterial quinolones are two essential enzymes involved in DNA replication, DNA gyrase (or topoisomerase II) and topoisomerase IV.<sup>18,19</sup>

Quinolones are amphoteric compounds with the chemical structure having a carboxyl group at the position 3 and an N-heterocycle (most frequently piperazine) at the position 7. Quinolones interact with metal ions through coordinated interactions that lead to the formation of metal complexes. Consequently, quinolones may participate in the chelation with metal ions as unidentate, bidentate and bridging ligands (Figure 1). A large number of articles focused on

\*Corresponding author: Gabriel Hancu, Tel: +40 265 215551 / 267 or 167, Fax: +40 265 210407, Email: gabriel.hancu@umftgm.ro

©2018 The Authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.


metal complexes of antibacterial quinolones and their biological effects have been published and reviewed.<sup>5-8,12</sup>

Among antibacterial quinolones, metal complexes with promising therapeutic effects are the silver complexes.<sup>20-</sup>

<sup>23</sup> Reports regarding several silver complexes have been developed recently to increase antibacterial activity of ligands or acquire new biological effects.<sup>1,2,14,16,24</sup> The particular coordination of antibacterial quinolones in

metal complexes with one silver ion bridge and two silver ions bridge have been previously described.<sup>12</sup>

Obtaining methods of silver complexes with quinolones are relatively simple. Several factors are essential for the future structure of the complex such as silver: ligand ratio, solvents, pH of the reaction mixture, temperature and time. These particular conditions have been described and commented in one of our previous review.<sup>12</sup>



**Figure 1.** The general structure of antibacterial quinolones and the most common modes of complexation with metal ions (R7 is a heterocycle in various derivatives).

### Analysis methods

The ability of quinolones to coordinate metal ions has been harnessed firstly, in the analyses of pharmaceuticals, biological and food samples. Analysis protocols that have been published comprise different methods such as spectrophotometry,<sup>25-30</sup> spectrofluorimetry,<sup>31-34</sup> chemiluminescence,<sup>35,36</sup> atomic absorption spectrometry,<sup>28</sup> flow injection and high-performance liquid chromatography.<sup>37-39</sup>

To determine the chemical composition and chemical structure of organometallic complexes, a broad category of methods can be used. After synthesis, elemental analysis (C, H, N) is used to elucidate molecular formula and to describe silver complexes.<sup>40</sup>

The most often applied analysis methods are the spectroscopic ones which provide information regarding the chemical structure and complexation model.

### Flame Atomic Absorption Spectrometry (FAAS)

FAAS was successfully used to quantify the silver content in silver complex with quinolones.<sup>1,14</sup> This method is applicable when the sample is in solution or readily solubilized; also, the technique is remarkably free from external interferences.<sup>41-44</sup> However, FAAS is characterized by low sensitivity, is highly expensive and requires extra abilities for the operator.<sup>45</sup> Consequently, the accessibility of FAAS methods is limited.

### Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES)

ICP-AES has been used to analyze a wide variety of sample types.<sup>41</sup> ICP proved to be a suitable method for the determination of silver percentage in the complexes with quinolones.<sup>24</sup>

ICP-AES replaced to some extent the FAAS technique, but it is also characterized by low sensitivity due to high excitation temperature. Therefore most samples are entirely atomized, and some chemical and spectroscopic interferences may occur. This method requires liquid samples preparation identically to those used for FAAS.<sup>41,46</sup>

### Fourier-transform infrared spectroscopy (FT-IR)

FT-IR is a versatile technique, which can register spectra from samples in liquid, solid or gaseous state.<sup>47</sup> FT-IR is a useful analytical method for identification and elucidation of chemical interaction in complexes, by comparing the IR spectrum of the complex to ligand IR spectrum. The essential FT-IR band assignments for antibacterial quinolones are characteristic to  $\nu(\text{C}=\text{O})_{\text{carb}}$  (carboxyl),  $\nu(\text{C}=\text{O})_{\text{pyr}}$  (pyridone), and  $\nu(\text{N}\cdots\text{H})$ , N atoms from heterocycle (R7).<sup>48</sup> The recorded FT-IR data disclose that the 4-pyridone and 3-carboxyl groups are implicated in the coordination of silver. Also, registered FT-IR data could be emphasized in studies if the N

atoms from heterocycle moiety (R7) are involved in complexation of silver.

The advantages of the FT-IR method are the capacity to record spectra with high signal-to-noise ratios (SNR), possibility to increase the number of scans to improve SNR, and wave number precision. However, the disadvantages of FT-IR technique are well known which are mainly represented by artifacts such as carbon dioxide and water vapor peaks.<sup>49</sup>

#### **Nuclear Magnetic Resonance Spectroscopy (NMR)**

NMR is the most frequently used analytical method for the elucidation of the chemical structure.<sup>50,51</sup> In metal complexes of quinolones analysis, <sup>1</sup>H NMR and <sup>13</sup>C NMR spectroscopy techniques are used to indicate the chelation of the metal ion to the ligand by the observed chemical shifts in the metal complex spectrum comparative to that of ligand. The significant shifts occur as a result of the implication of some moieties in the coordination process (e.g., nitrogen atoms from piperazinyl substituent).<sup>1,14</sup>

Though, NMR has a lower sensitivity than the MS technique due to amounts of samples of several micrograms. The significant disadvantages of this analytical method are the high cost of deuterated solvents, the price of the spectrometer, and the complexity of the software.<sup>51</sup>

#### **Mass spectrometry (MS)**

MS is an important analytical method used to obtain qualitative and quantitative information about compounds after their conversion to ions.<sup>52</sup> The different types of ionization sources available today allow characterization of a broad spectrum of compounds, most often based on an electrospray process.<sup>53,54</sup> However, MS technique is less encountered in the molecular weights determinations of silver complexes of antibacterial quinolones despite the facts that the method has a high sensitivity and requires a short time of analysis.<sup>55</sup> The coordination of silver ions has lower stability when the silver complexes are analyzed by Electrospray Ionisation Mass Spectrometry (ESI-MS) technique. Another disadvantage is related to their weak solubility in solvents compatible with the MS interface.<sup>14,23</sup>

#### **Ultraviolet-visible spectroscopy (UV-VIS)**

The UV-VIS electronic absorption spectra is a useful and straightforward spectroscopic method for studying organometallic complexes. Mainly, this technique is an easy-to-use, robust, reliable and cheap method which is used to check the identity of drugs. The UV-VIS spectra of parent compounds are compared and evaluated with the UV-VIS spectra of silver complexes.<sup>56</sup>

The electronic absorption spectrum is of two types such as *d-d* spectrum and charge transfer spectrum. The *d-d* spectrum deals with the electronic transitions within the *d*-orbitals. In the charge – transfer spectrum, electronic transitions occur from metal to ligand or vice-versa.<sup>57</sup>

Thus, the electronic spectra of silver complexes measured in the solid state could indicate distinct differences by comparison to the spectra of the parent compound.<sup>14,24,57,58</sup> For example, the hypsochromic and hyperchromic shifts occur in the main band of levofloxacin spectrum, and a new band occurs in the spectrum of synthesized silver complexes (at 440-450 nm).<sup>14</sup>

#### **Fluorescence spectroscopy**

Antibacterial quinolones present fluorescence properties which are exploited in analytical determination.<sup>59-61</sup> Fluorescence spectroscopy analysis could be applied for the identification of fluorescent properties of quinolones and their metal complexes and to study the equilibrium constants of complexes.<sup>21,24</sup>

As an example, the fluorescent emission intensity of complex Ag(H-norfloxacin)<sub>2</sub>(NO<sub>3</sub>) in the solid state was larger than that of free H-norfloxacin (about two times). The coordination of norfloxacin to silver ion probably increases the ligand conformational rigidity; thereby this diminishes the nonradiative decay of the ( $\pi - \pi^*$ ) excited state of the ligand.<sup>21</sup> Instead, the fluorescent emission intensity of polymer Ag<sub>4</sub>(Hciprofloxacin)<sub>2</sub>(ciprofloxacin)<sub>2</sub>(NO<sub>3</sub>)]·4H<sub>2</sub>O decreases significantly probably due to weak Ag–Ag interactions, leading to a reduced luminescence intensity.<sup>22</sup> Competitive studies by fluorescence spectroscopy method using 3,8-diamino-5-ethyl-6-phenyl-phenanthridinium bromide (EB) also show the capability of silver complexes to displace EB from the DNA-intercalator EB.<sup>62,63</sup> Silver complexes can quench the EB–DNA fluorescence. Thus, these experimental data prove that EB was in direct competition with silver complexes at the same binding site for bind to DNA through intercalation.<sup>24,64-66</sup> Fluorescence spectroscopy is as an adequate, sensitive, reliable and relatively cheap technique and could be useful for analysis of silver complexes and their effects.<sup>67,68</sup>

#### **Single-crystal X-ray diffraction (XRD)**

The most powerful method of analysis of metal complexes is XRD, used to determine the arrangement of atoms of a crystalline solid in three-dimensional space and analysis of the interatomic distances.<sup>69,70</sup> Meanwhile with the development of the importance of crystallography field the program SHELXL has been improved and became more performant to validate and archive crystal structures. Also, SHELXL is accessible to academics. SHELXTL and SHELXS 97 are improved version of the software.<sup>71</sup> Both SHELXTL and SHELXS 97 refined by full-matrix least-squares based on  $F^2$ (SHELXL 97)<sup>72</sup> were used in the analysis of some silver complexes of antibacterial quinolones.<sup>16,17,22,24</sup> XRD is a fast (under 20 minutes) and powerful technique for identification of unknown compounds which needs minimal sample preparation, wide availability of units, and relatively uncomplicated data interpretation. However, XRD presents some limitation related to the

standard reference database, homogeneity of the sample, detection limits, and peak overlay.<sup>73</sup> Unfortunately, these analysis methods seem to be less accessible to researchers due to the costs of equipment.

### **Thermal analysis techniques**

The thermal methods complete analysis of silver complexes by revealing their specific thermophysical properties. This category includes Differential Scanning Calorimetry (DSC), Thermogravimetric analysis (TGA), Derivative thermo-gravimetric (DTG) analysis, and Differential thermal analysis (DTA) which gather information about the stability of the complexes (decomposition temperature, kinetics of decomposition, water content).<sup>74</sup> If the complexation process occurs, the silver complexes will present notable differences of thermal curves from the ligand (melting point, number of decomposition phases, exothermic or endothermic peaks).<sup>1,14,16,23</sup>

DTA and DSC methods offer similar qualitative data. However, DTA can be used at higher temperatures than DSC, but more reliable quantitative information is obtained by DSC.<sup>74</sup> The precision of these methods is limited due to the dependence of the DTA signals from experimental conditions. DSC is the appropriate thermal method to investigate structure transitions in polymers due to the high sensitivity to any change in the sample or crucible and heating rate. DSC and TGA are both sensitive to the heating rate and sample masses. TGA provides reliability related to mass changes, but due to small amounts of samples, the non-homogeneous materials cannot be tested.<sup>75</sup>

### **Density functional theory (DFT)**

DFT is an increasingly used computational technique suitable to predict the properties of molecules: molecular structures, atomization energies, vibrational frequencies, electric and magnetic properties, ionization energies, reaction paths etc.<sup>76,77</sup>

The fully optimized geometries of the parent compound and metal complex are presented graphically. Selected bond angles, total energy, charge density, and total dipole moment are also essential data in the characterization of metal complexes. The most common density functionals in DFT model chemistry for some silver complex with levofloxacin, B3LYP (Becke's three-parameter exchange with Lee, Yang, and Parr correlation functional) for levofloxacin and MWB28 for silver.<sup>14</sup> The only limitation of the DFT method is the selection of the density functionals. Nowadays, there are no systematic principles of selecting the density functionals. The most published density functionals have been derived by comparison with experimental data.<sup>78</sup>

### **Conductivity measurements**

Conductivity measurements provide data about electrolytic or non-electrolytic nature of complexes.<sup>79,80</sup> Generally, the silver complexes of antibacterial

quinolones were soluble in aprotic solvents such as dimethylsulfoxide (DMSO), and dimethylformamide (DMF) and conductivity values are lower than an electrolyte 1:1.<sup>14,79,80</sup>

Conductivity measurement is an extensive and useful method, with high reliability, sensitivity, and relatively low cost. Also, the measure of conductivity is a rapid and inexpensive technique to determine the ionic strength of a solution. However, it is a non-specific technique due to inability to distinguish between different types of ions.<sup>81</sup> Based on previously published data we choose to summarise some relevant methods suitable for studies of silver complexes with fluoroquinolones (Table 1).

### **Discussion**

Correlation and combination of the data obtained from several methods of analysis is a requirement to characterize the chemical structure of a certain organometallic complex. In the preliminary analysis, it is necessary to collect data related to physical properties (appearance, solubility, conductivity, stability) followed by the establishment of the molecular formula that can be determined by elemental analysis combined with FAAS or ICP. To determine the silver content, both FAAS and ICP-AES methods can be used successfully. In a metal complex of an antibacterial quinolone, the polarity of a metal ion is reduced as a consequence to the partial sharing of positive charge with the ligand donor groups and of the overlap with the ligand orbitals. The delocalization of  $\pi$  electrons is increased over the whole chelate ring. Thereby, the lipophilic nature of metal ion is increased. This modification enhances the passage of the complex through the lipid membranes and cells penetration.<sup>4</sup>

The XRD technique is not a very accessible method for researchers, consequently to determine the complexity of the metal ions various combinations of spectroscopic, thermal and computational methods may be used.

Among the spectroscopic methods, FT-IR and MS provide the most unequivocal evidence of silver ion chelation with quinolone ligands. Also, NMR, UV VIS and fluorescence spectroscopy complete and reinforce the data obtained through FT-IR and MS techniques. The DSC seems to be the most appropriate thermal analysis method to assess the behavior of complexes at temperature change in comparison to parent compounds. Nowadays, DTF is a common computational technique suitable to predict the properties of molecules and fulfill the requirements of silver complexes with quinolones characterization.

Obtaining the most accurate chemical structure of silver complexes is closely related to its biological properties, such as the identification of pharmacophores,<sup>82</sup> interaction with DNA and antibacterial activity,<sup>83,84</sup> blocking of topoisomerase II enzymes (inhibiting the DNA repair activity) and possible anti-cancer, anti-fungal and anti-parasitic effects.<sup>2,4,18</sup>

**Table 1.** Analysis methods suitable for silver complexes with antibacterial quinolones derivatives.

Analysis method	Purpose	Mode	Complexes	References
Elemental analysis	Determination of %C, %H și %N	Comparison found vs. calculated %C, %H și %N of complexes	Most of the synthesized silver complexes.	
Atomic absorption spectroscopy (AAS)	Determination of % Ag	Comparison found vs. calculated %Ag	Ag(MXF)(H-IMZ)]·2.5H <sub>2</sub> O	1
Flame atomic absorption spectroscopy (FAAS)	Determination of % Ag	Comparison found vs. calculated %Ag	(LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	14
Inductively coupled plasma (ICP) spectroscopy	Determination of % Ag	Comparison found vs. calculated %Ag	[Ag <sub>2</sub> (PIP) <sub>2</sub> ] <sub>2</sub> ·8H <sub>2</sub> O	24
FT-IR spectroscopy	Identification and elucidation of chemical structure	Comparison that of ligand IR spectrum, in the range of 400–4000 cm <sup>-1</sup> by preparing sample pellets with KBr	Most of the synthesized silver complexes.	
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectrometry	Elucidation of the chemical structure	Indicating the coordination of the fluoroquinolone to the silver ion	Ag(MXF)(H-IMZ)]·2.5H <sub>2</sub> O (LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	1, 14
Mass spectrometry	Identification and elucidation of chemical structure	Comparison that of ligand mass spectrum AEI MS 30 mass spectrometer at 70 eV	[Ag <sub>2</sub> (NFX) <sub>2</sub> ](NO <sub>3</sub> ) <sub>2</sub> (LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	14,23
Electronic spectroscopy	Identification and elucidation of chemical structure	Comparison that of ligand UV-VIS spectrum	[Ag(NLX) <sub>2</sub> ] Ag <sub>4</sub> (H-CPF) <sub>2</sub> (CPF) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O [Ag <sub>2</sub> (NFX) <sub>2</sub> ](NO <sub>3</sub> ) <sub>2</sub> Ag(MXF)(H-IMZ)]·2.5H <sub>2</sub> O	1,16,22,23
Fluorescence spectroscopy	Identification of fluorescent properties of complexes	Comparison that of ligand fluorescence spectrum	Ag <sub>4</sub> (H-CPF) <sub>2</sub> (CPF) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O [Ag <sub>2</sub> (PIP) <sub>2</sub> ] <sub>2</sub> ·8H <sub>2</sub> O	22,24
X-ray crystallography/Single-crystal X-ray diffraction (XRD)	Determination of atom arrangements in crystalline solid in three-dimensional space Analysis the interatomic distances	SHELXS 97 refined by full-matrix least-squares based on F <sup>2</sup> (SHELXL 97) SHELXTL	Ag <sub>4</sub> (H-CPF) <sub>2</sub> (CPF) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O [Ag <sub>2</sub> (H-ENX) <sub>4</sub> ](NO <sub>3</sub> ) <sub>2</sub> [Ag(NLX) <sub>2</sub> ] [Ag <sub>2</sub> (PIP) <sub>2</sub> ] <sub>2</sub> ·8H <sub>2</sub> O	16,17,22,24
DTF calculations	Show full geometry optimizations of the ligand and silver complexes	DFT method in Gaussian09	(LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	14
Differential scanning calorimetry (DSC) analysis	Characterisation of the thermophysical properties	DSC curves: (10°C min <sup>-1</sup> temperature increase rate from 40 to 400°C, weight of the samples 3 mg).	(LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	14
TGA-DTG-DTA	Gathering information about the stability of the complex (decomposition temperature, kinetics of decomposition, water content)	- TGA: from ambient temperature to 800°C, nitrogen atmosphere - TGA: an oxygen atmosphere (10°C min <sup>-1</sup> temperature increase rate from ambient temperature to 1000°C) - TGA, DTG, and DTA: a dynamic nitrogen atmosphere(20 mL min <sup>-1</sup> ), 10°C min <sup>-1</sup> temperature increase rate from 25 up to 800°C	[Ag <sub>2</sub> (NFX) <sub>2</sub> ](NO <sub>3</sub> ) <sub>2</sub> Ag(MXF)(H-IMZ)]·2.5H <sub>2</sub> O [Ag(NLX) <sub>2</sub> ]	1,16,23
Molar conductivity measurements	Indicates if the compounds are of electrolytic or non-electrolytic nature	10 <sup>-3</sup> M solution of complexes in DMSO	[Ag(PFX)(H-IMZ)]·2H <sub>2</sub> O (LVF) <sub>2</sub> Ag(NO <sub>3</sub> ) (LVF) <sub>2</sub> Ag(NO <sub>3</sub> )(CH <sub>3</sub> OH) (LVF)Ag(C <sub>6</sub> H <sub>6</sub> O <sub>7</sub> )·3H <sub>2</sub> O	2,14
Kinetic data	The activation energy and the size of the ion allow a particular approach of the ligand.	Coats–Redfern and Horowitz–Metzger methods	[Ag <sub>2</sub> (NFX) <sub>2</sub> ](NO <sub>3</sub> ) <sub>2</sub>	23
Measurement of equilibrium constants of complexes	Calculation of the stability constant, K	Fluorescence measurements	Ag(H-NFX) <sub>2</sub> (NO <sub>3</sub> )	21

Abbreviations: CPF – ciprofloxacin, ENX – enoxacin, IMZ – imidazole, LVF – levofloxacin, MXF–moxifloxacin, NLX – nalidixic acid, NFX – norfloxacin, PFX – pefloxacin, PIP – pipemidic acid, SHELXS 97, SHELXL 97- software names (programs for the refinement of crystal structures from diffraction data), Gaussian 09 - computational chemistry program, full-matrix least-squares based on F<sup>2</sup> – refinement method

## Conclusion

The complexation of silver ions with antibacterial quinolones represents a research area of increasing importance, taking into consideration the acute and current problem of bacterial resistance to many marketed drugs. Pharmaceutical properties of antibacterial quinolones can be optimized by obtaining silver complexes with enhanced antibacterial activity or other biological properties. Characterization of metal complexes with derivatives from the quinolone class can be accomplished through a series of analytical methods. The selection of analytical methods is a compromise between the consistency of the corroborated data and the accessibility of the experimental approach. This review describes the main steps to follow for obtaining reliable structural information regarding silver complexes with quinolones and comprises a synthesis of classical and modern analytical methods. The most accurate characterization of an organometallic compound is a prerequisite for determining and understanding the mechanisms of biological effects.

## Ethical Issues

Not applicable.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## References

- Soayed AA, Refaat HM, Noor El-Din DA. Metal complexes of moxifloxacin-imidazole mixed ligands: Characterization and biological studies. *Inorganica Chim Acta* 2013;406:230-40. doi: 10.1016/j.ica.2013.04.040
- Soayed AA, Refaat HM, Noor El-Din DA. Characterization and biological activity of Pefloxacin-imidazole mixed ligands complexes. *Inorganica Chim Acta* 2014;421:59-66. doi: 10.1016/j.ica.2014.05.020
- Akinremi CA, Obaleye JA, Amolegbe SA, Adediji JF, Bamigboye MO. Biological activities of some Fluoroquinolones-metal complexes. *Int J Med Biomed Res* 2012;1(1):24-34. doi: 10.14194/ijmbr.115
- Uivarosi V. Metal complexes of quinolone antibiotics and their applications: an update. *Molecules* 2013;18(9):11153-97. doi: 10.3390/molecules180911153
- Turel I. The Interactions of Metal Ions with Quinolone Antibacterial Agents. *Coord Chem Rev* 2002;232:27-47. doi: 10.1016/S0010-8545(02)00027-9
- Lipunova GN, Nosova EV, Charushin VN. Metallocomplexes of fluoroquinolonecarboxylic acids. *Russ J Gen Chem* 2009;79(12):2753-66. doi: 10.1134/S1070363209120342
- Serafin A, Stanczak A. The Complexes of Metal Ions with Fluoroquinolones. *Russ J Coord Chem* 2009;35(2):81-95. doi: 10.1134/S1070328409020018
- Singh R, Debnath A, Masram DT, Rathore D. Synthesis and Biological Activities of Selected Quinolone-Metal Complexes. *Res J Chem Sci* 2013;3(6):83-94.
- Psomas G, Kessissoglou DP. Quinolones and non-steroidal anti-inflammatory drugs interacting with copper(II), nickel(II), cobalt(II) and zinc(II): structural features, biological evaluation and perspectives. *Dalton Trans* 2013;42(18):6252-76. doi: 10.1039/c3dt50268f
- Feio MJ, Sousa I, Ferreira M, Cunha-Silva L, Saraiva RG, Queirós C, et al. Fluoroquinolone-metal complexes: a route to counteract bacterial resistance? *J Inorg Biochem* 2014;138:129-43. doi: 10.1016/j.jinorgbio.2014.05.007
- Rusu A, Hancu G, Uivarosi V. Fluoroquinolone pollution of food, water and soil, and bacterial resistance. *Environ Chem Lett* 2015;13(1):21-36. doi: 10.1007/s10311-014-0481-3
- Rusu A, Hancu G, Munteanu AC, Uivarosi V. Development perspectives of silver complexes with antibacterial quinolones: Successful or not? *J Organomet Chem* 2017;839:19-30. doi: 10.1016/j.jorganchem.2017.02.023
- Rusu A, Hancu G, Tóth G, Toma F, Mare AD, Man A, et al. Synthesis, characterization and microbiological activity evaluation of two silver complexes with norfloxacin. *Farmacia* 2016;64(6):922-32.
- Rusu A, Hancu G, Tóth G, Vancea S, Toma F, Mare AD, et al. New silver complexes with levofloxacin: Synthesis, characterization and microbiological studies. *J Mol Struct* 2016;1123:384-93. doi: 10.1016/j.molstruc.2016.07.035
- El-Gamel NE. Silver(I) complexes as precursors to produce silver nanowires: structure characterization, antimicrobial activity and cell viability. *Dalton Trans* 2013;42(27):9884-92. doi: 10.1039/C3DT33092C
- Debnath A, Mogha NK, Masram DT. Metal complex of the first-generation quinolone antimicrobial drug nalidixic acid: structure and its biological evaluation. *Appl Biochem Biotechnol* 2015;175(5):2659-67. doi: 10.1007/s12010-014-1450-9
- Zhong DC, Chen ZF, Liu YC, Luo XJ, Barta C, Liang H. Syntheses, crystal structures of Ni(II), Ag(I)-enoxacin complexes, and their antibacterial activity. *J Coord Chem* 2010;63(18):3146-54. doi: 10.1080/00958972.2010.507811
- Aldred KJ, Kerns RJ, Osheroff N. Mechanism of Quinolone Action and Resistance. *Biochemistry* 2014;53(10):1565-74. doi: 10.1021/bi5000564
- Fàbrega A, Madurga S, Giralt E, Vila J. Mechanism of action of and resistance to quinolones. *Microb Biotechnol* 2009;2(1):40-61. doi: 10.1111/j.1751-7915.2008.00063.x
- Baenziger NC, Fox CL, Modak SL. The structure of silver pefloxacin, an antibiotic related to nalidixic acid. *Acta Cryst* 1986;C42:1505-9. doi: 10.1107/S0108270186091679

21. Li YX, Chen ZF, Xiong RG, Xue Z, Ju HX, You XZ. A mononuclear complex of norfloxacin with silver(I) and its properties. *Inorg Chem Commun* 2003;6(7):819-22. doi: 10.1016/S1387-7003(03)00115-1
22. Chen ZF, Yu LC, Zhong DC, Liang H, Zhu XH, Zhou ZY. An unprecedented 1D ladder-like silver(I) coordination polymer with ciprofloxacin. *Inorg Chem Commun* 2006;9(8):839-43. doi: 10.1016/j.inoche.2006.04.034
23. Refat MS. Synthesis and characterization of norfloxacin-transition metal complexes (group 11, IB): Spectroscopic, thermal, kinetic measurements and biological activity. *Spectrochim Acta A Mol Biomol Spectrosc* 2007;68(5):1393-405. doi: 10.1016/j.saa.2006.12.078
24. Li MT, Sun JW, Sha JQ, Wu HB, Zhang EL, Zheng TY. An unprecedented Ag-pipemidic acid complex with helical structure: Synthesis, structure and interaction with CT-DNA. *J Mol Struct* 2013;1045:29-34. doi: 10.1016/j.molstruc.2013.04.028
25. Fratini L, Schapoval EES. Ciprofloxacin determination by visible light spectrophotometry using iron(III)nitrate. *Int J Pharm* 1996;127(2):279-82. doi: 10.1016/0378-5173(95)04290-3
26. Eboka CJ, Aigbavboa SO, Akerele JO. Colorimetric determination of the fluoroquinolones. *J Antimicrob Chemother* 1997;39(5):639-41. doi: 10.1093/jac/39.5.639
27. Rizk M, Belal F, Ibrahim F, Ahmed S, Sheribah ZA. Derivative spectrophotometric analysis of 4-quinolone antibacterials in formulations and spiked biological fluids by their Cu(II) complexes. *J AOAC Int* 2001;84(2):368-75.
28. El-Brashy AM, El-Sayed Metwally M, El-Sepai FA. Spectrophotometric and Atomic Absorption Spectroscopic Determination of Some Fluoroquinolone Antibacterials by Ion-pair Complex Formation with Bismuth (III) Tetraiodide. *J Chin Chem Soc* 2005;52(2):253-62. doi: 10.1002/jccs.200500038
29. Cazedey ECL, Bonfilio R, Araújo MB, Salgado HRN. A First-Derivative Spectrophotometric Method for the Determination of Ciprofloxacin Hydrochloride in Ophthalmic Solution. *Phys Chem* 2012;2(6):116-22. doi: 10.5923/j.pc.20120206.06
30. Shrivastava DR, Shrivastava A, Patel R. A review of spectrophotometric determination of antibiotic norfloxacin. *Int J Pharm Sci Res* 2017;8(9):3619-29. doi: 10.13040/IJPSR.0975-8232.8(9).3619-29
31. Tan H, Zhang Z, Chen Y. Detection of mercury ions (Hg<sup>2+</sup>) in urine using a terbium chelate fluorescent probe. *Sens Actuators B Chem* 2011;156(1):120-5. doi: 10.1016/j.snb.2011.03.081
32. Han Y, Wu X, Yang J, Sun S. The fluorescence characteristic of the yttrium-norfloxacin system and its analytical application. *J Pharm Biomed Anal* 2005;38(3):528-31. doi: 10.1016/j.jpba.2005.01.033
33. Kaur K, Saini S, Singh B, Malik AK. Highly sensitive synchronous fluorescence measurement of danofloxacin in pharmaceutical and milk samples using aluminium (III) enhanced fluorescence. *J Fluoresc* 2012;22(5):1407-13. doi: 10.1007/s10895-012-1079-4
34. Tao S, Li X, Wang C, Meng C. A Dual-Emission Amphiphile /Dye Modified Mesoporous Silica as Fluorescent Sensor for the Detection of Fe<sup>3+</sup>, Cr<sup>3+</sup> and Al<sup>3+</sup>. *ChemistrySelect* 2016;1(12):3208-14. doi: 10.1002/slct.201600672
35. Attia MS, Youssef AO, Essawy AA, Abdel-Mottaleba MSA. A highly luminescent complexes of Eu(III) and Tb(III) with norfloxacin and gatifloxacin doped in sol-gel matrix: A comparable approach of using silica doped Tb(III) and Eu(III) as optical sensor. *J Lumin* 2012;132(10):2741-6. doi: 10.1016/j.jlumin.2012.05.012
36. Murillo Pulgarín JA, Alañón Molina A, Jiménez García E. Simplex optimization of the variables influencing the determination of pefloxacin by time-resolved chemiluminescence. *Spectrochim Acta A Mol Biomol Spectrosc* 2018;193:117-24. doi: 10.1016/j.saa.2017.12.002
37. Soledad García M, Isabel Albero M, Sánchez-Pedreño C, Salem Abuherba M. Flow injection spectrophotometric determination of ofloxacin in pharmaceuticals and urine. *Eur J Pharm Biopharm* 2005;61(1-2):87-93. doi: 10.1016/j.ejpb.2005.03.007
38. Al-Momani IF, Haj-Hussein AT, Tahtamouni AN. Flow Injection Spectrophotometric and Chromatographic Determination of Ciprofloxacin and Norfloxacin in Pharmaceutical Formulations. *J Flow Injection Anal* 2008;25(2):151-5.
39. Takeda N, Gotoh M, Matsuoka T. Rapid screening method for quinolone residues in livestock and fishery products using immobilised metal chelate affinity chromatographic clean-up and liquid chromatography-fluorescence detection. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 2011;28(9):1168-74. doi: 10.1080/19440049.2011.587028
40. Ma TS. Elemental Analysis, Organic Compounds. In: Meyers RA, editor. *Encyclopedia of Physical Science and Technology*. 3rd ed. Orlando: Academic Press; 2003. PP. 393-405.
41. Ebdon L, Evans EH, Fisher AS, Hill SJ. An Introduction to Analytical Atomic Spectrometry. New York: John Wiley&Sons; 1998.
42. Sun JM, Xu P, Han XT, Sun HW. Determination of trace silver by flame atomic absorption spectrometry using chitosan enriching method. *Guang Pu Xue Yu Guang Pu Fen Xi* 2005;25(2):290-2.
43. Shakerian F, Haji Shabani AM, Dadfarnia S, Shabani M. Flame Atomic Absorption Spectrometric Determination of Trace Amounts of Silver after Solid-Phase Extraction with 2-Mercaptobenzothiazole Immobilized on

- Microcrystalline Naphthalene. *Journal of Chemistry* 2013;2013:6. doi: 10.1155/2013/465825
44. Bianchin JN, Martendal E, Carasek E. Determination of Trace Silver in Water Samples by Online Column Preconcentration Flame Atomic Absorption Spectrometry Using Termite Digestion Product. *J Autom Methods Manag Chem* 2011;2011:839365. doi: 10.1155/2011/839365
45. Karimia MA, Mohammadi SZ, Mohadesi A, Hatefi-Mehrjardi A, Mazloum-Ardakani M, Sotudehnia Korani L, et al. Determination of silver(I) by flame atomic absorption spectrometry after separation/preconcentration using modified magnetite nanoparticles. *Sci Iran* 2011;18(3):790-6. doi: 10.1016/j.scient.2011.06.008
46. Bings NH, von Niessen JO, Schaper JN. Liquid sample introduction in inductively coupled plasma atomic emission and mass spectrometry — Critical review. *Spectrochim Acta Part B At Spectrosc* 2014;100:14-37. doi: 10.1016/j.sab.2014.08.011
47. Stuart B. Infrared Spectroscopy: Fundamentals and Applications. New York: Wiley; 2004.
48. Dorofeev VL. Infrared spectra and the structure of drugs of the fluoroquinolone group. *Pharm Chem J* 2004;38(12):693-7. doi: 10.1007/s11094-005-0063-6
49. Smith BC. Fundamentals of Fourier Transform Infrared Spectroscopy. 2nd ed. New York: Taylor & Francis; 2016.
50. Macomber RS. A Complete Introduction to Modern NMR Spectroscopy. New York: Wiley Interscience Publication; 1998.
51. Elipe MVS. Advantages and disadvantages of nuclear magnetic resonance spectroscopy as a hyphenated technique. *Anal Chim Acta* 2003;497(1-2):1-25. doi: 10.1016/j.aca.2003.08.048
52. Schalley CA. Modern Mass Spectrometry. Topics in Current Chemistry, Volume 225. New York: Springer-Verlag; 2003.
53. Ho CS, Lam CW, Chan MH, Cheung RC, Law LK, Lit LC, et al. Electrospray Ionisation Mass Spectrometry: Principles and Clinical Applications. *Clin Biochem Rev* 2003;24(1):3-12.
54. Tang Q, Chen F, Xin X. Study of the Fragmentation Patterns of Nine Fluoroquinolones by Tandem Mass Spectrometry. *Anal Lett* 2012;45(1):43-50. doi: 10.1080/00032719.2011.565448
55. de Hoffmann E, Stroobant V. Mass Spectrometry: Principles and Applications. 3rd ed. Chichester, UK: John Wiley & Sons Ltd; 2007.
56. Watson DG. Pharmaceutical Analysis, A Textbook for Pharmacy Students and Pharmaceutical Chemists. 4th ed. London, UK: Churchill Livingstone; 2016.
57. Lever ABP. Inorganic Electronic Spectroscopy Volume 33 of Studies in physical and theoretical chemistry. New York: Elsevier; 1997.
58. Sadeek AS. Synthesis, thermogravimetric analysis, infrared, electronic and mass spectra of Mn(II), Co(II) and Fe(III) norfloxacin complexes. *J Mol Struct* 2005;753(1-3):1-12. doi: 10.1016/j.molstruc.2005.06.011
59. Zuyun H, Houping H, Ruxiu C, Yun'e Z. Study on the Fluorescence Properties of Some Fluoroquinolones in Various Media. *Wuhan Univ J Nat Sci* 1997;2(3):353-8. doi: 10.1007/BF02829921
60. Vilches AP, Nieto MJ, Mazzieri MR, Manzo RH. Structure-Fluorescence Relationships in Antimicrobial Fluoroquinolones (AMFQs). *Molecules* 2000;5(3):398-400. doi: 10.3390/50300398
61. Park HR, Kim TH, Bark KM. Physicochemical properties of quinolone antibiotics in various environments. *Eur J Med Chem* 2002;37(6):443-60. doi: 10.1016/S0223-5234(02)01361-2
62. Heller DP, Greenstock CL. Fluorescence lifetime analysis of DNA intercalated ethidium bromide and quenching by free dye. *Biophys Chem* 1994;50(3):305-12. doi: 10.1016/0301-4622(93)E0101-A
63. Vardevanyan PO, Antonyan AP, Parsadanyan MA, Davtyan HG, Karapetyan AT. The binding of ethidium bromide with DNA: interaction with single- and double-stranded structures. *Exp Mol Med* 2003;35(6):527-33. doi: 10.1038/emm.2003.68
64. Huang WY, Li J, Kong SL, Wang ZC, Zhu HL. Cu(II) and Co(II) Ternary Complexes of Quinolone Antimicrobial Drug Enoxacin and Levofloxacin: Structure and Biological Evaluation. *RSC Adv* 2014;4(66):35193-204. doi: 10.1039/C4RA05812G
65. Tarushi A, Raptopoulou CP, Psycharis V, Terzis A, Psomas G, Kessissoglou DP. Zinc(II) complexes of the second-generation quinolone antibacterial drug enrofloxacin: Structure and DNA or albumin interaction. *Bioorg Med Chem* 2010;18(7):2678-85. doi: 10.1016/j.bmc.2010.02.021
66. Skyrianou KC, Psycharis V, Raptopoulou CP, Kessissoglou DP, Psomas G. Nickel-quinolones interaction. Part 4. Structure and biological evaluation of nickel(II)-enrofloxacin complexes compared to zinc(II) analogues. *J Inorg Biochem* 2011;105(1):63-74. doi: 10.1016/j.jinorgbio.2010.09.007
67. Settle F. Handbook of Instrumental Techniques for Analytical Chemistry. New Jersey: Prentices Hall; 1997.
68. René AJ. Principle and Applications of Fluorescence Spectroscopy. Oxford: Blacwell Publishing; 2008.
69. Ni TW, Tofanelli MA, Ackerson CA. Chapter 5 - Structure Determination by Single Crystal X-Ray Crystallography. In: Tsukuda T, Häkkinen H, editors. *Frontiers of Nanoscience, Volume 9, Protected Metal Clusters: From Fundamentals to Applications*. Amsterdam: Elsevier; 2015. PP. 103-25.
70. Hansford GM. Phase-Targeted X-Ray Diffraction. *J Appl Crystallogr* 2016;49(Pt 5):1561-71. doi: 10.1107/S1600576716011936
71. Sheldrick GM. Crystal structure refinement with SHELXL. *Acta Crystallogr C Struct Chem* 2015;71(Pt 1):3-8. doi: 10.1107/S2053229614024218

72. Sheldrick GM, Schneider TR. SHELXL: High-resolution refinement. *Methods Enzymol* 1997;277:319-43.
73. Bunaciu AA, Udriștioiu EG, Aboul-Enein HY. X-ray diffraction: instrumentation and applications. *Crit Rev Anal Chem* 2015;45(4):289-99. doi: 10.1080/10408347.2014.949616
74. Brown ME. Introduction to Thermal Analysis Techniques and Applications. New York: Kluwer Academic Publishers; 2004.
75. Feist M. Thermal Analysis: Basics, Applications, and Benefit. *ChemTexts* 2015;1:8. doi: 10.1007/s40828-015-0008-y
76. Mujahid M, Kia AF, Duff B, Egan DA, Devereux M, McClean S, et al. Spectroscopic Studies, DFT Calculations, and Cytotoxic Activity of Novel Silver(I) Complexes of Hydroxy Ortho-Substituted-Nitro-2H-Chromen-2-One Ligands and a Phenanthroline Adduct. *J Inorg Biochem* 2015;153:103-13. doi: 10.1016/j.jinorgbio.2015.10.007
77. Yang Y, Gao H. Comparison of DFT Methods for Molecular Structure and Vibration Spectra of Ofloxacin Calculations. *Spectrochim Acta A Mol Biomol Spectrosc* 2012;85(1):303-9. doi: 10.1016/j.saa.2011.10.019
78. Tsipis AC. DFT flavor of coordination chemistry. *Coord Chem Rev* 2014;272:1-29. doi: 10.1016/j.ccr.2014.02.023
79. Geary WJ. The use of conductivity measurements in organic solvents for the characterisation of coordination compounds. *Coord Chem Rev* 1971;7(1):81-115. doi: 10.1016/S0010-8545(00)80009-0
80. Vieira LM, de Almeida MV, Lourenço MC, Bezerra FA, Fontes AP. Synthesis and antitubercular activity of palladium and platinum complexes with fluoroquinolones. *Eur J Med Chem* 2009;44(10):4107-11. doi: 10.1016/j.ejmech.2009.05.001
81. Conductivity Theory and Practice. Villeurbane Cedex. France: Radiomete Analytical SAS; 2004. [cited 2018 May 9]. Available from: [http://www.analytical-chemistry.uoc.gr/files/items/6/618/agwgimometria\\_2.pdf](http://www.analytical-chemistry.uoc.gr/files/items/6/618/agwgimometria_2.pdf)
82. Mehrotra R, Shukla SN, Gaur P, Dubey A. Identification of Pharmacophore in Bioactive Metal Complexes: Synthesis, Spectroscopic Characterization and Application. *Eur J Med Chem* 2012;50:149-53. doi: 10.1016/j.ejmech.2012.01.049
83. Patel MN, Chhasatia MR, Gandhi DS. Interaction of Drug Based Binuclear Mixed-Ligand Complexes with DNA. *Bioorg Med Chem* 2009;17(15):5648-55. doi: 10.1016/j.bmc.2009.06.023
84. Patel MN, Gandhi DS, Parmar PA. DNA Interaction and In-Vitro Antibacterial Studies of Fluoroquinolone Based Platinum(II) Complexes. *Inorg Chem Commun* 2012;15:248-51. doi: 10.1016/j.inoche.2011.10.037